

GROUP 4A INSECTICIDE

GROUP 4 7 12 FUNGICIDES

PULL HERE TO OPEN ►



Insecticide with Fungicides

A seed treatment product for protection against damage from listed insects, seed-borne diseases, and seedling diseases on dried shelled beans and soybean.

Active Ingredients:

Thiamethoxam ¹	20.80%
Mefenoxam ²	3.13%
Fludioxonil ³	1.04%
Sedaxane ⁴	1.04%

Other Ingredients

73.99%

Total:

100.00%

¹CAS No. 153719-23-4

²CAS No. 70630-17-0 and CAS No. 69516-34-3

³CAS No. 131341-86-1

⁴CAS No. 874967-67-6

CruiserMaxx Vibrance is a flowable concentrate for seed treatment containing 1.99 lb thiamethoxam, 0.30 lb mefenoxam, 0.10 lb fludioxonil and 0.10 lb sedaxane per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See additional precautionary statements and directions for use in booklet.

EPA Reg. No. 100-1508 EPA Est. 100-NE-001

SCP 1508A-L1B 0916

4071490

Product ID **55590**

15 gallons
Net Contents

®

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS
<p>HAZARDS TO HUMANS AND DOMESTIC ANIMALS</p> <p style="text-align: center;">CAUTION</p> <p>Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.</p> <p>PERSONAL PROTECTIVE EQUIPMENT (PPE) Applicators and other handlers must wear:</p> <ul style="list-style-type: none"> • Long-sleeved shirt and long pants • Chemical-resistant gloves made of: barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, polyvinyl chloride (PVC) ≥14 mils, Viton® ≥14 mils • Shoes plus socks <p>Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.</p> <p>ENGINEERING CONTROLS</p> <p>When handlers use closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.</p>
<p>USER SAFETY RECOMMENDATIONS</p> <p>Users should:</p> <ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
<p>ENVIRONMENTAL HAZARDS</p> <p>This product is toxic to wildlife, freshwater and estuarine/marine fish, oysters, and shrimp and highly toxic to aquatic invertebrates. Runoff may be hazardous to aquatic organisms in neighboring areas. Exposed treated seed may be hazardous to wildlife. Do not contaminate water when disposing of equipment washwater or rinsate.</p>

continued...

PRECAUTIONARY STATEMENTS (*continued*)

Pollinator Precautions

Thiamethoxam is highly toxic to bees, and effects are possible as a result of exposure to translocated residues in blooming crops.

Groundwater Advisory

Mefenoxam is known to leach through soil into ground water under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow, and may result in groundwater contamination.

Fludioxonil has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow, and may result in groundwater contamination.

Thiamethoxam has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow, and may result in groundwater contamination.

Physical and Chemical Hazards

Do not use, pour, spill or store near heat or open flame. Do not store near or use with oxidizing agents.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For use in commercial seed treatment facilities. Use is also permitted as an end-use seed treatment on agricultural establishments at planting, or immediately before planting, as specified in the Specific Crop Use Directions. This product is to be used in liquid or slurry treaters only.

RESTRICTIONS

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Maximum usage when applying both metalaxyl- and mefenoxam-containing products to the same crop within the same season: Do not apply more than the maximum seasonal total for the active ingredient as stated on the label of the product containing the lowest seasonal total on that crop.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves: barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, polyvinyl chloride (PVC) ≥14 mils, Viton ≥14 mils
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR INSECT AND/OR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

Treatment of highly mechanically scarred or damaged seed or seed known to be of low vigor and poor quality may result in reduced germination and/or reduction of seed and seedling vigor. Treat a quantity of seed using equipment similar to that planned for treating the total seed lot. Prior to treatment, conduct germination tests on a portion of seed before committing the total seed lot to a selected seed treatment.

Due to seed quality, crop or variety sensitivity, and seed storage conditions beyond the control of Syngenta, no claims are made to guarantee the germination of seed or propagating material for all crop seed when treated with CruiserMaxx Vibrance.

USE INFORMATION

CruiserMaxx Vibrance is a seed treatment product containing the active ingredients thiamethoxam (insecticide) and fludioxonil, mefenoxam and sedaxane (fungicides). CruiserMaxx Vibrance protects against damage from listed early-season insects, soil-borne and seed-borne diseases on dried shelled beans, and soybean.

Thiamethoxam is a systemic seed treatment insecticide belonging to the neonicotinoid class of chemistry. Thiamethoxam protects against listed chewing and sucking insects through contact and ingestion.

Mefenoxam fungicide is active against *Pythium*, *Phytophthora* and systemic downy mildew.

Fludioxonil fungicide is active against *Fusarium* and *Rhizoctonia*, and suppresses seed-borne *Sclerotinia* and *Phomopsis* species.

Sedaxane fungicide is active against seed decay, seedling blight and damping-off caused by *Rhizoctonia* species.

RESISTANCE MANAGEMENT

GROUP	4A	INSECTICIDE
-------	----	-------------

CruiserMaxx Vibrance contains thiamethoxam, a Group 4A insecticide. Thiamethoxam is a systemic insecticide belonging to the neonicotinoid class of chemistry which includes nicotinic acetylcholine receptor (nAChR) agonists.

Insect populations may contain individuals naturally resistant to Group 4A insecticides and if used repeatedly in the same fields, then resistant members may eventually dominate the population. Because resistance development cannot be predicted, use sound resistance management strategies established for the crop and use area.

Base seed treatment on an integrated pest management program that includes field sanitation, historical information related to pesticide use, careful selection of pest-tolerant crop varieties, scouting, and management practices which optimize populations of natural enemies of insect pests such as within-field refugia (untreated areas). Sound management programs also consider cultural and biological control practices.

In order to maintain susceptibility to this class of chemistry:

- Use products at their full, specified doses.
- Use appropriate, well-maintained equipment. Use specified water volumes and apply at optimal temperatures in order to obtain optimal treatment.
- When rate ranges are given, use the higher rate within the listed rate range when insect pressure is expected to be high.
- Avoid using a single active ingredient or mode of action (same insecticide group) exclusively for season long control of insect species with more than one generation per crop season.
- For insect species with successive or overlapping generations, use a treatment window approach. A treatment window is a period of time defined by the stage of crop development and the biology of the pests of concern. Within the treatment window, depending on the length of residual activity, single or consecutive applications may be made using seed, in-furrow, or foliar treatments unless otherwise excluded by product labels. Do not exceed the maximum amount of this insecticide's mode of action allowed per growing season.
- Following a treatment window of this insecticide's mode of action, rotate to a treatment window of effective products with a different mode of action before making additional applications of this insecticide.

If resistance to this product develops in your area, this product or other products with a similar mode of action may not provide adequate control. If poor performance cannot be attributed to improper application or weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for the crop and use area.

Syngenta encourages responsible product stewardship to ensure effective long term control of the insect pests on this label.

For additional information on Insect Resistance Management:

- Contact Syngenta representatives at 1-800-334-9481
- Contact your local Cooperative Extension Service specialist, pest control advisor, or certified crop advisor
- Visit the Insecticide Resistance Action Committee (IRAC) on the web at: <http://www.irc-online.org>

GROUP	4	7	12	FUNGICIDES
-------	---	---	----	------------

CruiserMaxx Vibrance contains mefenoxam, a Group 4 fungicide; sedaxane, a Group 7 fungicide; and fludioxonil, a Group 12 fungicide. Mefenoxam belongs to the phenylamide class of chemistry which interferes with fungal RNA synthesis. Sedaxane is a succinate dehydrogenase inhibitor (SDHI) and belongs to the carboxamide class of chemistry which disrupts cellular respiration and energy generation. Fludioxonil belongs to the phenylpyrrole class of chemistry which interferes with osmotic signal transduction.

Fungal populations may contain individuals naturally resistant to Group 4, 7, or 12 fungicides and if used repeatedly in the same fields, then resistant members may eventually dominate the population. Because resistance development cannot be predicted, the use of this product should conform to sound resistance management strategies such as alternation with fungicides with a different mode of action and/or tank mixes established for the crop and use area.

Use should be based on an IPM program that includes field sanitation, scouting, historical information related to pesticide use, and crop rotation. The IPM program should also consider cultural, biological, and other chemical control practices.

Syngenta encourages responsible product stewardship to ensure effective long term control of the fungal diseases on this label.

For additional information on Fungicide Resistance Management:

- Contact Syngenta representatives at 1-800-334-9481
- Contact your local extension specialist or certified crop advisor
- Visit the Fungicide Resistance Action Committee (FRAC) on the web at: <http://www.frac.info>

MIXING PROCEDURES

Important: Always re-circulate CruiserMaxx Vibrance thoroughly before using.

The typical density of CruiserMaxx Vibrance is 9.56 pounds per gallon. Consult the manufacturer of the application equipment you plan to use for suitability for this application and for instructions on operation and calibration of the equipment. Follow the manufacturer application instructions for the seed treatment equipment being used.

Apply CruiserMaxx Vibrance as a water-based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of disease control. Thoroughly mix the specified amount of CruiserMaxx Vibrance into the required amount of water or liquid inoculant for the slurry treater and dilution rate to be used.

Certain crops require addition of inoculants when the seed is treated or planted. CruiserMaxx Vibrance is compatible with several liquid inoculant products. Consult the maker of the inoculant product and a Syngenta representative for directions before applying CruiserMaxx Vibrance with inoculants.

The total application volume must be sufficient to provide desired level of coverage. Dilution is typically done with water or liquid inoculants. The minimum slurry volume to achieve adequate coverage is 4.0 fluid ounces per 100 pounds of seed. More diluent may be required to obtain complete coverage.

Continuous agitation or mixing of the slurry mixture is necessary to prevent settling out of the solution.

Allow seed to dry before bagging.

Follow planter manufacturer specifications for use of talc or other hopper box additives at planting. Seed must be completely dry before adding to planter.

Use an EPA approved dye/colorant that imparts an unnatural color to the seed as required in 40CFR 153.155(c).

CROP ROTATION RESTRICTIONS

- In the event of crop failure or harvest of a crop grown from seed treated with CruiserMaxx Vibrance, the field may be replanted according to the following schedule:

Plantback Interval Table

Immediate Plantback	Minimum 30-Day Plantback Interval
Canola	Alfalfa
Cereal Grains: Barley, Corn (Field, Pop, Sweet), Oats, Rye, Sorghum, Triticale, and Wheat	Cereal Grains Crop Group 15
Cotton	Cucurbit Vegetables Crop Group 9
Dried Shelled Pea and Bean Crop Subgroup 6C	Fruiting Vegetables Crop Group 8
Oilseeds: Borage, Crambe, Flax Seed, and Mustard Seed	Head and Stem Brassica Crop Subgroup 5A
Potato	Leafy Brassica Greens Crop Subgroup 5B
Soybean	Leafy Vegetables Crop Group 4
Sugarbeet	Legume Vegetables (Succulent or Dried) Crop Group 6
	Mint: Peppermint and Spearmint
	Onion, Dry Bulb
	Peanut
	Root Vegetables Crop Subgroup 1A

Plantback Interval Table (continued)

Immediate Plantback	Minimum 30-Day Plantback Interval
	Safflower Strawberry Sunflower Tobacco Tuberous and Corm Vegetables (Except Potato) Crop Subgroup 1D

- For any other crops the minimum plant back interval is 120 days from the date the seeds treated with CruiserMaxx Vibrance were planted. A cover crop other than the crops listed above that is planted for erosion control or soil improvement may be planted sooner than the 120 day interval; however, the crop may not be grazed or harvested for food or feed.

SEED BAG LABEL REQUIREMENTS

The Federal Seed Act requires that bags containing treated seeds shall be labeled with the following statements:

- This seed has been treated with thiamethoxam insecticide and fludioxonil, mefenoxam, and sedaxane fungicides.
- Do not use for feed, food, or oil purposes.
- User is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.
- Use an EPA approved dye/colorant that imparts an unnatural color to the seed as required in 40CFR 153.155(c).

In addition, the U.S. Environmental Protection Agency requires the following statements on bags containing seeds treated with CruiserMaxx Vibrance:

- **Ground Water Advisory:** Mefenoxam is known to leach through soil into ground water under certain conditions as a result of agricultural use. Fludioxonil and Thiamethoxam have properties and characteristics associated with chemicals detected in ground water. Mefenoxam, Fludioxonil, and Thiamethoxam may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.
- **Pollinator Precautions:** Thiamethoxam is highly toxic to bees, and effects are possible as a result of exposure to translocated residues in blooming crops.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in the ethanol by-products that are used in agronomic practice.
- Do not allow children, pets, or livestock to have access to treated seed.
- Store away from feeds and foodstuffs.
- Wear long-sleeved shirt, long pants and chemical resistant gloves when handling treated seed.

- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Dispose of all excess treated seed. Leftover treated seed may be doublesown around the headland or buried away from water sources in accordance with local requirements.
- Do not contaminate water bodies when disposing of planting equipment wash waters.
- Dispose of seed packaging in accordance with local requirements.
- In the event of crop failure or harvest of a crop grown from seed treated with CruiserMaxx Vibrance, the field may be replanted according to the following schedule:

Plantback Interval Table

Immediate Plantback	Minimum 30-Day Plantback Interval
Canola Cereal Grains: Barley, Corn (Field, Pop, Sweet), Oat, Rye, Sorghum, Triticale, and Wheat Cotton Dried Shelled Pea and Bean Crop Subgroup 6C Oilseeds: Borage, Crambe, Flax Seed, Mustard Seed Potato Soybean Sugarbeet	Alfalfa Cereal Grains Crop Group 15 Cucurbit Vegetables Crop Group 9 Fruiting Vegetables Crop Group 8 Head and Stem Brassica Crop Subgroup 5A Leafy Brassica Greens Crop Subgroup 5B Leafy Vegetables Crop Group 4 Legume Vegetables (Succulent or Dried) Crop Group 6 Mint: Peppermint and Spearmint Onion, Dry Bulb

Plantback Interval Table (continued)

Immediate Plantback	Minimum 30-Day Plantback Interval
	Peanut Root Vegetables Crop Subgroup 1A Safflower Strawberry Sunflower Tobacco Tuberous and Corm Vegetables (Except Potato) Crop Subgroup 1D

- For any other crops the minimum plant back interval is 120 days from the date the seeds treated with CruiserMaxx Vibrance were planted. A cover crop other than the crops listed above that is planted for erosion control or soil improvement may be planted sooner than the 120 day interval; however, the crop may not be grazed or harvested for food or feed.
- For soybeans: Do not use at a rate that will result in more than 0.266 lb thiamethoxam per acre (120 grams ai/A) per calendar year, regardless of type of application (seed treatment and/or foliar). Do not use at a rate that will result in more than 0.083 lb thiamethoxam per acre (37.8 grams ai/A) per calendar year as a seed treatment application. Do not apply more than 0.004 lb (2.0 g) of fludioxonil per acre per calendar year as a seed treatment. Do not apply more than 0.013 lb (5.7 g) of mefenoxam per acre per calendar year as a seed treatment.
- For dried shelled peas and beans (except soybeans) crop subgroup 6C: Do not use at a rate that will result in more than 0.083 lb thiamethoxam per acre (37.8 grams ai/A) per calendar year. Do not apply more than 0.004 lb (2.0 g) of fludioxonil per acre per calendar year as a seed treatment. Do not apply more than 0.013 lb (5.7 g) of mefenoxam per acre per calendar year as a seed treatment.
- This seed has been treated with 0.0756 mg ai thiamethoxam per seed.
- With the exception of soybeans, do not make any soil or foliar application of products containing thiamethoxam to crops grown from seed treated with a neonicotinoid insecticide. For soybeans, do not apply a neonicotinoid insecticide within 45 days of planting seed treated with CruiserMaxx Vibrance.
- The maximum number of applications per season is 2.

CROP USE DIRECTIONS

When applied according to the **CRUISERMAXX VIBRANCE RATE TABLE**, CruiserMaxx Vibrance provides early season protection against injury by aphids, bean leaf beetle, grape colaspis, leaf miners, leaf hoppers, Mexican bean beetle, seed corn maggot, three-cornered alfalfa hopper, thrips, white grubs, and wireworm.

CruiserMaxx Vibrance provides protection against damping-off and seed borne rots due to Pythium, Phytophthora, Fusarium, Rhizoctonia species and early season Phytophthora root rot as well as Anthracnose on dry beans, and soybean caused by seed-borne *Colletotrichum* spp. CruiserMaxx Vibrance also suppresses seed-borne Sclerotinia and Phomopsis species. Additional Apron XL® may be necessary for high levels of *Phytophthora* or *Pythium*. Refer to the table below to determine the appropriate use rate of Apron XL, if needed. Read and follow all label direction for Apron XL use.

It is the pesticide user's responsibility to ensure that all products in a tank mix are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products of the mixture (for example, first aid from one product, REI from another).

CruiserMaxx Vibrance Rate Table

Crop	Use rate (fl oz per 100 lb seed)		Grams active ingredient per 100 kg seed	
	Cruiser Maxx Vibrance	Apron XL	Cruiser Maxx Vibrance	Apron XL
Dried Shelled Beans including: Bean <i>Lupinus</i> species, including: grain sweet white white sweet lupin Bean, <i>Phaseolus</i> species, including: field bean kidney bean lima bean, dry navy bean pinto bean tepary bean wax bean yellow bean Bean, <i>Vigna</i> species, including: adzuki bean blackeyed pea catjang cowpea Crowder pea moth bean mung bean rice bean southern pea urd bean	3.22	0.32	Thiamethoxam 50 gm	Mefenoxam 7.5 gm
			Mefenoxam 7.5 gm	
			Fludioxonil 2.5 gm	
			Sedaxane 2.5 gm	
			Total = 62.5 gm	

CruiserMaxx Vibrance Rate Table (continued)

Crop	Use rate (fl oz per 100 lb seed)		Grams active ingredient per 100 kg seed	
	Cruiser Maxx Vibrance	Apron XL	Cruiser Maxx Vibrance	Apron XL
Broad Bean (fava bean, dry)	3.22	0.32	Thiamethoxam 50 gm	Mefenoxam 7.5 gm
Guar			Mefenoxam 7.5 gm	
Lablab Bean			Fludioxonil 2.5 gm Sedaxane 2.5 gm Total = 62.5 gm	

CruiserMaxx Vibrance Rate Table

Crop	Rate of CruiserMaxx Vibrance		
	fl oz per 100 lb seed ¹ or fl oz per 140,000 seeds ¹	grams ai per 100 kg seed	mg ai per seed
Soybean, including soybean, vegetable	3.22 fl oz or 1.50 fl oz per 140,000 seeds	Thiamethoxam 50 gm Mefenoxam 7.5 gm Fludioxonil 2.5 gm Sedaxane 2.5 gm Total = 62.5 gm	Total of all active ingredients = 0.0945 mg

¹The mg ai per seed, fl oz CruiserMaxx Vibrance per 100 lb seed, and fl oz CruiserMaxx Vibrance per 140,000 seeds rates are based on 3,000 seeds per pound.

STORED GRAIN PROTECTION

When treated according to the directions for post-planting protection against listed pests, CruiserMaxx Vibrance will also provide protection during post treatment storage of the seed listed on this label against damage from the following stored grain insects: Indian Meal Moth (*Plodia interpunctella*), Rice Weevil (*Sitophilus oryza*), Red Flour Beetle (*Tribolium castaneum*), and Lesser Grain Borer (*Rhizopertha dominica*).

If the seed to be treated has existing infestations of stored grain insects, fumigate the seed with a registered product approved for such use prior to treating with CruiserMaxx Vibrance and bagging.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in a cool, dry place. Storage for extended periods above 90°F is not recommended.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

Pesticide Disposal

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling [less than or equal to 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling [greater than 5 gallons – mini-bulk]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling [greater than 5 gallons – bulk]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.

Apron XL[®], CruiserMaxx[®], Vibrance[®], the ALLIANCE FRAME 
the SYNGENTA Logo and the PURPOSE ICON 
are Trademarks of a Syngenta Group Company

Viton[®] is a trademark of E. I. du Pont de Nemours & Company, Inc.

©2016 Syngenta

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1508A-L1B 0916
4071490

GROUP 4A INSECTICIDE

GROUP 4 7 12 FUNGICIDES



Insecticide with Fungicides

A seed treatment product for protection against damage from listed insects, seed-borne diseases, and seedling diseases on dried shelled beans and soybean.

Active Ingredients:	
Thiamethoxam ¹	20.80%
Mefenoxam ²	3.13%
Fludioxonil ³	1.04%
Sedaxane ⁴	1.04%
Other Ingredients	73.99%
Total:	100.00%

¹CAS No. 153719-23-4

²CAS No. 70630-17-0 and CAS No. 69516-34-3

³CAS No. 131341-86-1

⁴CAS No. 874967-67-6

CruiserMaxx Vibrance is a flowable concentrate for seed treatment containing 1.99 lb thiamethoxam, 0.30 lb mefenoxam, 0.10 lb fludioxonil and 0.10 lb sedaxane per gallon.

See additional precautionary statements and directions for use in booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-1508

EPA Est. 100-NE-001

Product ID **55590**

CruiserMaxx®, Vibrance® and the SYNGENTA Logo are trademarks of a Syngenta Group Company.

©2016 Syngenta

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1508A-L1B 0916
4071490

15 gallons
Net Contents

KEEP OUT OF REACH OF CHILDREN. CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

FIRST AID

If **swallowed**: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **HOT LINE NUMBER**: For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), call **1-800-888-8372**.

ENVIRONMENTAL HAZARDS: This product is toxic to wildlife, freshwater and estuarine/marine fish, oysters, and shrimp and highly toxic to aquatic invertebrates. Runoff may be hazardous to aquatic organisms in neighboring areas. Exposed treated seed may be hazardous to wildlife. Do not contaminate water when disposing of equipment washwater or rinsate.

Pollinator Precautions: Thiamethoxam is highly toxic to bees, and effects are possible as a result of exposure to translocated residues in blooming crops.

Groundwater Advisory: Mefenoxam is known to leach through soil into ground water under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow, and may result in groundwater contamination.

Fludioxonil has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow, and may result in groundwater contamination.

Thiamethoxam has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow, and may result in groundwater contamination.

Physical and Chemical Hazards: Do not use, pour, spill or store near heat or open flame. Do not store near or use with oxidizing agents.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place. Storage for extended periods above 90°F is not recommended.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling: Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.

