

---

**SUPPLEMENTAL LABELING**

---

**Syngenta Crop Protection, LLC**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300

**SCP 1571B-S1 1116**

<b>GROUP</b>	<b>U15</b>	<b>FUNGICIDE</b>
--------------	------------	------------------

**Orondis® Gold 200**

Fungicide

**This supplemental label expires on 11/14/2019 and must not be used or distributed after this date.**

Active Ingredient:

Oxathiapiprolin (CAS No. 1003318-67-9)..... 18.7%

Other Ingredients:..... 81.3%

Total:..... 100.0%

**KEEP OUT OF REACH OF CHILDREN.**

**EPA Reg. No. 100-1571**

**All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed. Before using Orondis as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

The Syngenta logo, featuring the word "syngenta" in a lowercase, sans-serif font with a registered trademark symbol. A stylized leaf icon is positioned above the letter "g".

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### CROP USE DIRECTIONS

#### Tuberous and Corm Vegetables, Crop Subgroup 1C

Crops (including all cultivars, varieties, and/or hybrids of these)			
Arrachaca	Chayote (root)	Sweet potato	
Arrowroot	Chufa	Tanier	
Artichoke, Chinese	Dasheen (taro)	Turmeric	
Artichoke, Jerusalem	Ginger	Yam bean	
Canna, edible	Leren	Yam, true	
Cassava, bitter and sweet	Potato		
Target Disease	Rate	Application Timing	Use Directions
Pink rot ( <i>Phytophthora erythroseptica</i> )	4.8 fl oz/A	Apply as an in-furrow application at planting.	Apply as a 6- to 8-inch band directly over the seed pieces in-furrow, and then close furrows.
<b>Resistance Management:</b>			
<ul style="list-style-type: none"> <li>Do not follow soil applications of oxathiapiprolin-containing products with foliar applications of oxathiapiprolin-containing products. Use either soil applications or foliar applications but not both for disease control.</li> </ul>			
USE RESTRICTIONS			
<ol style="list-style-type: none"> <li>Refer to <b>Use Restrictions</b> section of the booklet for additional product use restrictions.</li> <li><b>Maximum Annual Rate:</b> <b>Soil:</b> 19.2 fl oz/A/year (0.25 lb ai/A/year of oxathiapiprolin-containing products)</li> <li>Do not use on Tuberous and Corm Vegetables in California</li> <li><b>Pre-harvest Interval (PHI):</b> 5 days</li> </ol>			

Orondis® and the Syngenta logo are trademarks of a Syngenta Group Company

©2016 Syngenta