

# Supplemental Labeling



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Sapphire®

EPA Reg. No. 62719-547

This supplemental label expires on May 6, 2021 and must not be used or distributed after this date.

### Use of Sapphire on Noncropland Areas

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Sapphire® specialty herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Sapphire according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Sapphire.

### Directions for Use

Refer to product label for Precautions and Restrictions, Mixing Instructions and Application Directions.

### Noncropland Sites

Use Sapphire specialty herbicide for postemergence control of annual and perennial broadleaf weeds in non-cropland including: airports, tops and outer banks of irrigation ditches/canals, military lands, non-irrigation ditch banks, roadsides, unimproved rough turf, vacant lots including: restoration sites, campgrounds, parks, prairie management, trails and trailheads, recreation areas, wildlife openings and wildlife habitat and management areas. For postemergence application and control of susceptible annual and perennial broadleaf weeds, apply Sapphire with a non-ionic surfactant at 0.25% v/v.

For total vegetation control on a site where no vegetation is desired, tank mixing this product with other herbicides may be necessary depending on the weeds on the site and their stage of growth at application.

Control existing vegetation by cultivation, hand weeding, or use of a postemergence herbicide in a tank mix with Sapphire.

Weeds Controlled or Suppressed	Rate	
	Fl oz/acre	Fl oz/1000 sq ft
Chickweed (common) Chickweed (mouseear) Clover, hop Clover, red Clover, white Dandelion, common Dollarweed English-lawn daisy False dandelion Florida betony Ivy Ground lespedeza, common medic, black Plaintain, broadleaf Plantain, buckhorn Parsley-piert Virginia buttonweed	13-24	0.30 – 0.55
Kyllinga, green (suppression only) <sup>1</sup> Nutsedge, purple <sup>1</sup> Nutsedge, yellow <sup>1</sup>	16-24	0.37 – 0.55

<sup>1</sup>Re-treatment may be necessary.

#### Specific Use Restrictions

- Refer to Mixing Directions section for tank mixing precautions. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.
- Do not apply more than 2.3 pints (36 fl oz) of Sapphire (0.09 lb ai penoxsulam) per acre per year.
- Do not apply more than a total of 24 fl. oz. (0.06 lb ai penoxsulam) of Sapphire per acre in a single application.
- Do not graze livestock or feed the forage cut from areas treated with Sapphire.
- For postemergence control of susceptible annual and perennial broadleaf weeds, apply Sapphire with a non-ionic surfactant at 0.25% v/v.
- Do not make more than 2 application(s) per year.
- **Retreatment Interval:** 28 days

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R354-006

EPA accepted: 03/06/18

Replaces R354-003