

**RESTRICTED USE PESTICIDE  
Due to Acute Toxicity**

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

**syngenta.**

**INTRA**

**Section 24(c) Special Local Need Label**

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF TEXAS**

**Gramoxone® SL 2.0  
Control of Volunteer Cotton in Corn**

**EPA Reg. No. 100-1431  
EPA SLN No. TX-130009**

**This label expires and must not be distributed or used in accordance with this  
SLN registration after December 31, 2021**



**DANGER / POISON  
PELIGRO / VENENO**

**DIRECTIONS FOR USE**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

Crop	Use Pattern	Rate Per Acre	Precautions, Restrictions and Comments
<b>FIELD CORN POPCORN SWEETCORN SEED CORN</b>	Postemergence Directed Spray (including hooded or shielded)	1.0-2.0 pt/A	<ul style="list-style-type: none"> <li>• Make no more than 3 applications per year for all postemergent directed sprays</li> <li>• Minimum total spray volume: 10 gal/A</li> <li>• Apply when cotton is actively growing</li> <li>• Do not apply when weather conditions favor drift</li> <li>• Severe damage and/or complete kill can occur if spray contacts corn plants</li> </ul> <p><b>HOODED OR SHIELDED SPRAYERS</b></p> <ul style="list-style-type: none"> <li>• To avoid excessive crop phytotoxicity, use a hooded or shielded sprayer with skids or wheels on the spray boom to maintain spray height.</li> <li>• Apply by directing spray between the rows and using hooded or shielded sprayers to prevent spray contact with crop plants.</li> </ul> <p><b>DIRECTED SPRAY WITHOUT HOODED OR SHIELDED SPRAYERS</b></p> <ul style="list-style-type: none"> <li>• Apply when corn is at least 10" tall with nozzles arranged to spray no higher than the lower 3" of corn stalks.</li> <li>• Corn plants shorter than 10" may be injured and not recover (corn height measured from soil surface to top of whorl).</li> <li>• For corn greater than 20" tall, arrange the nozzles to spray no higher than the lower 1/3 of the corn stalks.</li> <li>• Corn foliage sprayed will be injured, but the crop will recover and develop normally.</li> </ul>
<b>FIELD CORN POPCORN SEED CORN</b>	Harvest Aid Broadcast	1.0-2.0 pt/A	<ul style="list-style-type: none"> <li>• Make ONE (1) application at least 7 days prior to harvest.</li> <li>• Minimum total spray volume: Ground 20 gal/A; Air 5 gal/A</li> <li>• Apply after the corn is mature after the black layer has formed at the base of the kernels (this indicates maturity). Consult your local agricultural authority for help in identifying the black layer.</li> </ul>

**Additional Precautions, Restrictions and Comments:**

- Apply between cotyledon and 6 leaf stage of the volunteer cotton plants. Cotton plants larger than the 6 leaf stage may not be completely killed.
- Use higher rates as size of volunteer cotton plant increases
- Add a nonionic surfactant containing at least 75% surface active ingredient at 0.25% v/v
- Drought stressed cotton plants can be difficult to kill and desiccation may not be complete.

24(c) Registrant:  
 Syngenta Crop Protection, LLC  
 P. O. Box 18300  
 Greensboro, NC 27419-8300

Label Code: TX1431117CA0317