



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, NC 27709
1-866-99BAYER (1-866-992-2937)

SERENADE OPTIMUM®

Reduced application rates recommended for SERENADE OPTIMUM on Fruiting Vegetables

EPA Reg. No. 264-1160

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE IN THE STATES OF FLORIDA AND GEORGIA ONLY

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

Reduced application rates recommended for SERENADE OPTIMUM on Fruiting Vegetables.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation. For use only as described on this label. Not for isolation or deformulation. Do not culture.

SERENADE OPTIMUM may be tank-mixed with other registered pesticides to enhance plant disease control. This product cannot be mixed with any product with a prohibition against such mixing. When tank-mixing SERENADE OPTIMUM with other registered pesticides, always read and follow all use directions, restrictions, and precautions of both SERENADE OPTIMUM and the tank-mix partner(s). Use of the resulting tank mix must be in accordance with the more restrictive label limitations and precautions. Do not exceed label dosage rates.

Crops	Disease	Rate oz/acre	Application Instructions
Fruiting Vegetables Pepper Tomato Eggplant Ground Cherry Tomatillo Okra and other fruiting vegetables	Bacterial Spot <i>Xanthomonas</i> spp. Target Spot <i>Corynespora</i> <i>Cassicola</i>	8 to 12	Begin application soon after emergence or transplant and when environmental conditions are conducive to disease development. Continue applications on a 2- to 7-day interval or as needed. When conditions are conducive to rapid disease development, for improved control, use SERENADE OPTIMUM in a tank mix program with copper-based bactericides registered for control of Bacterial Spot at labeled rates.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information, Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).