



**Bayer CropScience LP**  
**P.O. Box 12014**  
**2 T.W. Alexander Drive**  
**Research Triangle Park, NC 27709**  
**1-866-99BAYER (1-866-992-2937)**

## SERENADE OPTIMUM®

SERENADE OPTIMUM is recommended to suppress Citrus Canker on Citrus

**EPA Reg. No. 264-1160**

# PRODUCT BULLETIN

### FOR DISTRIBUTION AND USE IN THE STATE OF FLORIDA ONLY

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

### **SERENADE OPTIMUM is recommended to suppress Citrus Canker on Citrus**

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the applicable directions, restrictions and precautionary statements plus all the other information appearing on the EPA-registered product label.

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation. For use only as described on this label. Not for isolation or deformulation. Do not culture.

SERENADE OPTIMUM may be tank-mixed with other registered pesticides to enhance plant disease control. This product cannot be mixed with any product with a prohibition against such mixing. When tank-mixing SERENADE OPTIMUM with other registered pesticides, always read and follow all use directions, restrictions, and precautions of both SERENADE OPTIMUM and the tank-mix partner(s). Use of the resulting tank mix must be in accordance with the more restrictive label limitations and precautions. Do not exceed label dosage rates.

Crops	Disease	Rate oz/acre	Application Instructions
<b>Citrus</b> Orange Grapefruit Lemon Tangerine Tangelo Pummelo and other citrus crops	<b>Citrus Canker</b> <i>Xanthomonas citri</i> subsp. <i>Citri</i> .	8	Begin foliar applications at the onset of new leaf flush, when environmental conditions are conducive to disease development.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information, Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).