

Guarantee[®]

SEAWEED EXTRACT

Guarantee[®] is a versatile liquefied seaweed (*Ascophyllum nodosum*) extract. It can be used for agriculture, hydroponics, landscape, gardens and other growing applications.

Application Guidelines

Formulated for foliar or soil application.

Shake or agitate before use. Apply in the early morning or late afternoon for best results. Allow foliar product to dry on plant prior to irrigation. Use spray adjuvant for superior coverage.

Application Rates

Please see back panel for specific crop applications.

General recommended application rates are 1/2 to 1 liter Guarantee[®] per hectare (1 to 2 pints per acre.)

For smaller areas, use 60 ml Guarantee[®] per 3.78 liters of water. (2 oz. Guarantee[®] per gallon of water.)

0.0.1

GUARANTEED ANALYSIS

Soluble Potash (K₂O) 1.00%

Derived from: Seaweed Extract

Mixing Guidelines

Use a minimum of 3.8 liters of water carrier per 93 m² (1 U.S. gallon per 1,000 sq. ft.). Fill mix tank with half of required spray water and turn on agitation. Measure product into tank, add remaining water and other soluble materials.

Compatibility

This product is compatible with most fertilizers and other spray materials. Conduct a jar test as needed.

Storage and Use Precautions

Store in original containers at temperatures above freezing. Mix with other materials only at time of application and in exact amounts required. Do not return previously mixed product to its original container.

Keep out of reach of children. May be harmful if swallowed. Avoid contact with eyes, skin, and clothing. If irritation occurs, seek medical attention.

This product is not intended to be a complete fertilizer or nutrient source and should be used in conjunction with a regular fertilizer program.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

CONDITIONS OF SALE...The Manufacturer warrants only that product conforms to label description. Buyer and user agree to accept all liability associated with handling, use, and disposal of this product.

Not for use in organic crop and organic food production.

Florida: F1762



Ocean Organics[®]

P.O. Box 1448 • Waldoboro, ME 04572
888-312-0106 • www.oceanorganics.com

Corporate Office: Ann Arbor, MI

Net Contents : ■ 9.5 L (2.5 Gal)

Net Weight : 9.6 Kg (21.1 Lb)

Bulk Density: 8.42 lbs/gal at 68° F

■ 56.8 L (15 Gal)

57.3 Kg (126.4 Lb)

■ 208.2 L (55 Gal)

210.1 Kg (481.8 Lb)

■ 1041.0 L (275 Gal)

1050.7 Kg (2409 Lb)

© 2011 Ocean Organics[®] 05/2017

Application Rates and Timing

Recommended application rates are 1 to 4 pints [1/2 to 2 quarts] per acre. See below for specific crop applications

Alfalfa

1. Apply at planting, 1-2 pints/acre
2. Repeat after each cutting or at 2 to 4 week intervals, 1-2 pints/acre

Asparagus

1. New crop at fern growth in spring, 1-2 Pints/Acre
2. Mature crop at fern growth, 1-2 Pints/Acre
3. After cutting, 1-2 Pints/Acre

Blueberries

1. At pre-bloom, 2-4 Pints/Acre
2. At flowering, 2-4 Pints/Acre
3. At petal fall, 2-4 Pints/Acre
4. 30 days after last application, 2-4 Pints/Acre
5. 30 days after last application or 2 weeks prior to harvest, 2-4 Pints/Acre
6. Optional: 2-4 weeks after harvest for over wintering and increased fertile buds, 2-4 Pints/Acre

Cane Berries

1. At pre-bloom, approximately 2 weeks prior to bloom, 2-4 Pints/Acre
2. After petal fall, 2-4 Pints/Acre
3. 4 weeks after last application, 2-4 Pints/Acre
4. 4 weeks after last application, 2-4 Pints/Acre
5. 2 weeks before harvest, 2-4 Pints/Acre

Brassicas (Cauliflower, Broccoli, Cabbage, Kale and Brussels Sprouts)

1. At 4 to 6 leaf stage, 1-2 Pints/Acre
2. 2 weeks after 1st application, 1-2 Pints/Acre
3. At Head Initiation, 1-2 Pints/Acre

Bulb Vegetables

- (Onion, Garlic, Leeks, and Shallots)
1. At Planting, 1-2 pints per acre
 2. Repeat every 2-4 weeks until harvest is complete, 1-2 pints per acre

Citrus

1. Pre-bloom at Pinhead, 2 Pints/Acre
2. Post bloom at 2/3 petal fall, 2 Pints/Acre

3. At summer oil application, 2 Pints/Acre
4. Optional: with fall miticide, 2 Pints/Acre
5. For fresh market varieties: 6 to 8 weeks prior to harvest, 2 Pints/Acre
6. For non-bearing trees: apply at the onset of every flush (approx every 30 days), 2 Pints/Acre

Corn

1. At Planting, 1-2 pints/acre
2. 14" to pretassel, 1-2 pints/acre
3. 36", 1-2 pints/acre

Cotton

1. At planting, 1-2 pints/acre
2. Repeat every 2-3 weeks, 1-2 pints/Acre

Cucurbits (Watermelon, Cantaloupe, Honeydew, Cucumber, Squash, Muskmelon, and Pumpkins)

1. At 3 to 4 leaf stage, 1-2 Pints/Acre
2. At pre-bloom, 1-2 Pints/Acre
3. Every 14 days through harvest, 1-2 Pints/Acre

Grains (Wheat, Triticale)

1. At planting, 1-2 pints per acre
2. Follow with applications at 6", 12" and 18" growth stage, 1-2 pints/acre

Grapes (Wine)

1. At new growth, 1-2 Pints/Acre
2. At pre-bloom to ¼ bloom, 1-2 Pints/Acre
3. 10 days to 2 weeks or at fruit set, 1-2 Pints/Acre
4. 2-3 weeks later, 1-2 Pints/Acre

Grapes (Table, Raisin, Juice)

1. At new growth, 1-4 inch shoot, 1-2 pints/acre
2. Rachis Stretch application at 16" to 24" shoot growth, 2-4 pints/acre
3. Sizing application - include with sizing sprays, 2-4 pints per acre
4. Color: 1-2 pints per acre for 4-5 weeks starting at 20% colorbreak
5. Optional - Continue with weekly applications during the summer months, 1 pint per acre

Hops

1. Apply at start of training in the spring, 1-2 pints/acre
2. Follow with applications every 2-4 weeks until harvest, 1-2 pints/acre

Hydroponic Vegetables

1. 0.05 to 1.0 ounces per 100 gallons of water
- In soilless systems, apply continuously with each fertigation cycle. In closed systems, reapply every 7-14 days.

Leafy Vegetables (Celery, Lettuce, Spinach, Swiss Chard, Endive, Raddicchio, etc)

1. At 3 to 5 leaf stage, 1-2 Pints/Acre
2. Every 2 weeks until harvest, 1-2 Pints/Acre

Legumes - Dry and Succulent (Beans, Garbanzos, Lentils, Peas and Soybeans)

1. In furrow at planting, 2 Pints/Acre
2. At pre-bloom, 3 to 4 leaf stage, 2 Pints/Acre
3. Repeat every 2 to 4 weeks until harvest

Orchard Fruit

1. At pink buds, 1-2 Pints/Acre
2. At petal fall, 1-2 Pints/Acre
3. 2-3 weeks after 2nd spraying, 1-2 Pints/Acre
4. Every 2 weeks prior to harvest, 1-2 Pints/Acre
5. On late Apple Varieties, i.e. Jona Gold, Gala, late season applications 1 month and 2 weeks prior to harvest to improve red fruit color, 3 pints/acre

Pasture (Forage Grasses)

1. At early growth stage, 1-2 Pints/Acre
2. 1-2 weeks after mowing/grazing, 1-2 Pints/Acre

Peanuts

1. At pegging, 1-2 Pints/Acre
2. 2 weeks after 1st application, 1-2 Pints/Acre
3. 1-2 weeks after 2nd application, 1-2 Pints/Acre
4. 1-3 weeks after 3rd application, 1-2 Pints/Acre (May be optional)

Peppers

1. At 6" stage, 1-2 Pints/Acre
2. At pre-bloom, 1-2 Pints/Acre
3. At initial fruiting, 1-2 Pints/Acre
4. 2-3 days after picking, 1-2 Pints/Acre

Potatoes

1. At tuber set, 1-2 Pints/Acre
2. 10-14 days after 1st application, 1-2 Pints/Acre
3. At early bloom, 1 to 2 Pints/Acre

Rice

1. Apply 1-2 Pints/Acre 25-30 days after seeding at tiller initiation
2. Follow with 1-2 Pints/Acre at panicle emergence

Root and Tuber Vegetables (Beets, Carrots, Sweet Potatoes)

1. Apply at planting, 1-2 pints/Acre
2. Repeat every 2-4 weeks until harvest

Soybeans

1. At preplant, 1 to 2 Pints/Acre
2. At 3 to 4 leaf stage, 1 to 2 Pints/Acre
3. 2-3 weeks later, 1 to 2 Pints/Acre

Strawberries

1. 10 days to 2 weeks after emergence, 2-4 Pints/Acre
2. At pre-bloom, 2-4 Pints/Acre
3. At petal fall or fruit set, 2-4 Pints/Acre
4. Every 2-3 weeks through harvest, 2-4 Pints/Acre

Sunflowers

1. Apply 1-2 pints at planting
2. Follow with applications every 2-4 weeks, 1-2 pints/acre

Tomatoes (Fresh and Processing)

1. At Transplant, 1-2 pints/acre
2. Repeat every 2-4 weeks until harvest is complete (4 to 8 oz per fertigation)

Tree Nuts (Pecans, Walnuts, Almonds, Pistachios, etc)

1. At pre-bloom (2 weeks prior to bloom), 2-4 Pint/Acre
2. After petal fall, 2-4 Pint/Acre
3. Every 2-4 weeks during the summer months, 2-4 Pint/Acre
4. Optional: post harvest application 2-4 weeks after harvest, 2-4 Pint/Acre
5. Apply at planting of new tree stock, 1-2 pints/acre

Woody Perennials

1. Apply 2-4 pints/acre in the spring at regrowth
2. Follow with applications every 2-4 weeks