

Supplemental Label



Valent U.S.A. Corporation
1333 N. California Blvd., Ste. 600
Walnut Creek, CA 94596-8025

SELECT[®] 2 EC HERBICIDE

EPA Reg. No. 59639-3

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL AND PAMPHLET. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

FOR POSTEMERGENCE GRASS CONTROL IN:

Crop ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Use Rate Per Acre ⁽²⁾	Crop Oil Concentrate* Rate Per Acre ⁽³⁾	
Beet, Garden	30 days	6-8 fl. oz.	1% v/v in the finished spray volume.	
Carrot	30 days	6-8 fl. oz.	1% v/v in the finished spray volume.	
Canola ⁽⁴⁾	75 days	4-5 fl. oz.	1% v/v in the finished spray volume.	
Celery, Rhubarb	30 days	6-8 fl. oz.	1% v/v in the finished spray volume.	
Clover (for use on clover grown in the states of ID, OR and WA only)	15 days	6-16 fl. oz.	1% v/v in the finished spray volume.	
Cranberry	30 days	6-8 fl. oz.	1% v/v in the finished spray volume.	
Cucurbits, including: Cantaloupes (all) Cucumber Gherkin Honeydew Melon	Muskmelons (all) Pumpkin Squash (all) Watermelon	14 days	6-8 fl. oz.	1% v/v in the finished spray volume.
Flax ⁽⁵⁾	75 days	4-5 fl. oz.	1% v/v in the finished spray volume.	
Fruiting Vegetable, (except Tomato) including: Eggplant	Groundcherry Pepino Peppers (all) Tomatillo	20 days	6-8 fl. oz.	1% v/v in the finished spray volume.
Head and Stem Brassica Vegetables, including:	Broccoli Cabbage Cauliflower Brussels sprouts	30 days	6-8 fl. oz.	1% v/v in the finished spray volume.
Leafy Brassica Greens, ⁽⁶⁾ including: Broccoli Raab Cabbage, Chinese (Bok Choy)	Collards Kale Mustard Greens Mustard Spinach Rape Greens Turnip Greens	14 days	6-8 fl. oz.	1 % v/v in the finished spray volume.
Leaf Lettuce	14 days	6-8 fl. oz.	1% v/v in the finished spray volume.	
Mint ⁽⁶⁾	21 days	6-16 fl. oz.	1 % v/v in the finished spray volume.	
Mustard Seed ⁽⁴⁾	75 days	4-5 fl. oz.	1% v/v in the finished spray volume.	

Crop ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Use Rate Per Acre ⁽²⁾	Crop Oil Concentrate* Rate Per Acre ⁽³⁾
Potato, Sweet Potato , Yam and other tuberous and corn vegetables including: Artichoke Chinese Jerusalem Cassava Bitter Sweet Ginger	30 days	6-16 fl. oz.	1% v/v in the finished spray volume for all crops except potatoes. For potatoes: add crop oil concentrate at the rate of 1 qt./A for ground applications. AMS at the rate of 2.5 - 4.0 lbs./A may be added in potatoes.
Radish	15 days	6-8 fl. oz.	1% v/v in the finished spray volume.
Root Vegetables (except Radish), including: Chicory Ginseng Horseradish Turnip	30 days	6-8 fl. oz.	1% v/v in the finished spray volume.
Spinach⁽⁶⁾	14 days	6-8 fl. oz.	1 % v/v in the finished spray volume.
Strawberry	4 days	6-8 fl. oz.	1% v/v in the finished spray volume.
Sugar Beet	40 days	6-16 fl. oz.	1 qt./A for ground application. AMS at the rate of 2.5 - 4.0 lbs./A may be added.
Sunflower	70 days	6-16 fl. oz.	1 qt./A for ground application. AMS at the rate of 2.5 - 4.0 lbs./A may be added.

* Acceptable crop oil concentrates would be those that contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils.

- (1) SELECT 2 EC Herbicide is not recommended for use on vegetable crops being grown for seed production unless specific use directions are provided.
- (2) Do not apply more than 8 fl. oz./A in a single application, except for crops noted above that allow for rates up to 16 fl. oz./A. For repeat applications make on a minimum of a 14-day interval. For clover and radish do not apply more than 16 fl. oz. (0.25 lb. ai) per acre in a season.
- (3) Do not add nitrogen fertilizer source spray adjuvants to the spray mixture, unless specifically advised otherwise.
- (4) Do not apply SELECT 2 EC Herbicide to canola or mustard seed after the crop has begun bolting. Crop injury may occur when applied during the bloom period. Do not exceed 5 fl. oz. (0.08 lb. ai) per acre in a season.
- (5) Do not apply SELECT 2 EC Herbicide to flax during the bloom period. Crop injury may occur when applied during the bloom period. Do not exceed 5 fl. oz. (0.08 lb. ai) per acre in a season.
- (6) Registration pending in the state of New York – 8/02.

RESTRICTIONS AND LIMITATIONS

Do not apply more than 32 fl. oz. of SELECT 2 EC Herbicide (0.5 lb. ai) per acre per season. Application on Long Island, New York is restricted to no more than 16 fl. oz. of SELECT 2 EC Herbicide (0.25 lb. ai) per acre per season.

RECOMMENDATIONS FOR ANNUAL OR PERENNIAL GRASSES

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Use the high rate under heavy grass weed pressure and/or when grasses are at maximum height.
- For the control of annual grasses in crops other than canola, flax and mustard seed use 6 to 8 fl. oz./A, for canola, flax and mustard seed use 4 to 5 fl. oz./A. Do not exceed 5 fl. oz./A in canola, flax or mustard seed per acre per application.
- For the control of perennial grasses use 8 fl. oz./A, except for those crops listed above that allow for rates up to 16 fl. oz./A.
- Please refer to container label for specific use rates, weeds and weed sizes.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION. PLEASE REFER TO CONTAINER LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Copyright © 2002 by Valent U.S.A. Corporation

Manufactured for
Valent U.S.A. Corporation
1333 N. California Blvd., Suite 600
Walnut Creek, CA 94596-8025
www.valent.com
Form 2002-SEL-0011
Made in U.S.A.