



Bayer CropScience LP  
 P.O. Box 12014  
 2 T.W. Alexander Drive  
 Research Triangle Park, North Carolina 27709  
 1-866-99BAYER (1-866-992-2937)

**VELUM® PRIME**  
**For Suppression of Melanose and Greasy Spot in Citrus**  
**EPA Reg. No. 264-1078**

## **FIFRA 2(ee) Recommendation**

### **FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF FLORIDA AND TEXAS**

**This recommendation is valid until December 31, 2020**

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its label.

<b>Fruit, citrus, group 10-10</b>		
Australian Desert Lime; Australian Finger Lime; Australian Round Lime; Brown River Finger Lime; Calamondin; Citron; Citrus Hybrids; Grapefruit; Japanese Summer Grapefruit; Kumquat; Lemon; Lime; Mediterranean Mandarin; Mount White Lime; New Guinea Wild Lime; Orange, Sour; Orange, Sweet; Pummelo; Russell River Lime; Satsuma Mandarin; Sweet Lime; Tachibana Orange; Tahiti Lime; Tangelo; Tangerine (Mandarin); Tangor; Trifoliate Orange; Uniq Fruit; cultivars, varieties and/or hybrids of these.		
<b>Disease Suppression</b>	<b>Application Rate</b>	<b>Product Instructions</b>
Melanose <i>(Diaporthe citri)</i>	6.5 to 6.84 fl oz/acre	Apply at the critical timings for disease suppression. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 21-day interval.
Greasy spot <i>(Mycosphaerella citri)</i>		
<b>Restrictions:</b>		
<ul style="list-style-type: none"> <li>Do not apply more than 13.7 fl oz of VELUM PRIME (0.446 lbs Fluopyram) per acre per year, regardless of formulation or method of application (soil or foliar).</li> <li>Apply using chemigation equipment.</li> <li>Do not apply VELUM PRIME within 7 days of harvest.</li> <li>To limit the potential for development of disease resistance to this fungicide class, do not make more than 2 sequential applications of VELUM PRIME or any Group 7-containing fungicide before rotating with a fungicide from a different Group.</li> </ul>		

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.  
 For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any crop-protection product, always read and follow label directions.  
 For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*