



WILBUR-ELLIS®
Ideas to Grow With™

GUARANTEED ANALYSIS

Sulfur (S)3.00%
Zinc (Zn)7.00%

DERIVED FROM: Zinc Sulfate complexed with Lignosulfonate.



KEEP OUT OF REACH OF CHILDREN PRECAUTION

Do not ingest. Avoid contact with skin, eyes or clothing.
Avoid breathing dust, vapor or mist.

FIRST AID

In all cases, call a poison control center or doctor for further treatment advice.

IF SWALLOWED, call a poison control center or doctor immediately. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person. **IF ON SKIN**, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. **IF INHALED**, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration. **IF IN EYES**, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.

For chemical spills, leaks, fire or exposure,
call CHEMTREC: (800) 424-9300.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

Store in a safe manner. Store in original container only and keep tightly sealed when not in use. Dispose of unused product and empty containers in accordance with federal, state and local regulations.

GENERAL INFORMATION

FOLI-GRO® ZINC 7% is formulated primarily for foliar applications to prevent or correct zinc deficiencies in a wide range of agronomic and ornamental plants. Its use is suggested as a supplement to a regular, balanced fertilizer program to enhance yields and improve quality. Application of FOLI-GRO ZINC 7% is a means of obtaining a quick response to needed elements. Foliar applications of FOLI-GRO ZINC 7% may be particularly beneficial during periods of peak nutrient demand, for crops grown on soils having poor nutrient availability or to crops suffering from a weakened root system. Avoid foliar nutrient applications to crops during periods of moisture, salinity or heat stress. Applications of foliar nutrients to sensitive crops such as leafy vegetables or soft-skinned fruit during periods of excessive temperatures may cause cosmetic injury.

FOLI-GRO ZINC 7% should be used as part of a comprehensive Total Nutrition System® for optimizing plant growth, development, yield and quality.

DIRECTIONS FOR USE

FOLI-GRO ZINC 7% may be applied as a foliar spray by air or by ground. Maintain agitation to provide uniform distribution within the spray tank. Thorough and even spray coverage is essential for maximum nutrient uptake and crop safety. Always include a high quality wetting agent, such as R-11® or SYL-COAT®, at recommended rates. Optimum rate of application will vary based on a variety of factors including degree of need or deficiency, growth stage, yield potential, environmental conditions, existing soil fertility levels, soil pH, organic matter content, etc.

FOLIAR APPLICATIONS (Ground or Air): Apply the recommended amount of FOLI-GRO ZINC 7% per acre per application throughout the growing season as needed to maintain adequate levels for growth. Severe deficiencies may require more frequent applications. Use sufficient water to ensure good coverage. Do not exceed 1 quart per gallon of water. If applied too concentrated, slight burning of the foliage may occur.

SOIL APPLICATIONS: Use 2-4 quarts of FOLI-GRO ZINC 7% per acre as a preplant or side-dress application. Dilute with water for more uniform application. Always determine compatibility before preparing the application mix.

IRRIGATION SYSTEM APPLICATIONS (Soil, Drip or Sprinkler): FOLI-GRO ZINC 7% may be applied through irrigation systems at a rate of 0.5-1.5 gallons per acre. Before applying with other fertilizers always check compatibility. Contact your local Wilbur-Ellis Company representative for more information.

Fill spray tank 1/3-1/2 with water. With agitator running, add FOLI-GRO ZINC 7%. Agitate while adding remaining water and any other fertilizer. Add pesticide last and

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.html>

continue to agitate. Agitate well prior to spraying if mix has been left standing for a short time. Early correction of a zinc deficiency is desirable. Make the FOLI-GRO ZINC 7% application when the plants are young. Application of FOLI-GRO ZINC 7% will reduce zinc deficiencies occurring during periods of rapid plant growth or physiological stress.

BLUEBERRIES, CANEBERRIES, CRANBERRIES, GRAPES AND STRAWBERRIES: Apply 1-3 quarts per acre prior to bloom. Do not use FOLI-GRO ZINC 7% on fresh market grapes after fruit formation.

CUCURBITS: Apply 2-4 quarts per acre as needed. Allow a minimum of 14 days between applications.

FIELD CROPS: Apply 1-6 pints of FOLI-GRO ZINC 7% per acre depending on the degree of zinc deficiency.

GRASS SEED AND TURF: Apply 1-4 pints of FOLI-GRO ZINC 7% per acre in the normal spray program.

SMALL GRAIN CROPS (Including but not limited to barley, oats, rice, rye, and wheat): Apply 1-6 pints of FOLI-GRO ZINC 7% per acre per application between tillering and boot stage.

TREE CROPS: Apply 1-4 quarts per acre in sufficient spray volume for good cover. When applying by air in less than 20 gallons of water, use no more than 2 quarts per acre. Foliar zinc on plums followed by rain or sprinkler irrigation may cause cosmetic leaf shot-holing. Metals applied with 30 days of harvest may stain or ink fruit.

VEGETABLE CROPS (Including but not limited to carrots, cole crops, cucurbits, lettuce, onions, melons, potatoes, and tomatoes): Apply 1-6 pints FOLI-GRO ZINC 7% per acre per week in the normal spray program. For specific crop recommendations consult with the Wilbur-Ellis representative in your area.

COMPATIBILITY

FOLI-GRO ZINC 7% is compatible with many co-applied nutrient sources as well as plant protection products. Always jar test new combinations for compatibility prior to field mixing.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using the product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of the product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of many different factors including, without limitation, manner of use or application, weather, combination with other products, or crop conditions. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Manufacturer and Seller harmless from any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label. EXCEPT FOR THIS WARRANTY, THE PRODUCT IS FURNISHED "AS-IS," AND NEITHER SELLER NOR MANUFACTURER MAKES ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT; SELLER AND MANUFACTURER SPECIFICALLY DISCLAIM ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Buyer and User accept all risks arising from any use of this product, including without limitation uses contrary to label instructions, under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or Manufacturer.

To the extent permitted by law, neither Manufacturer nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE BUYER OR USER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT, OR, AT THE ELECTION OF MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

These Conditions of Sale and Limitation of Warranty and Liability shall be interpreted in accordance with the laws of the State of California, excluding its conflicts of laws rules, and may not be amended by any oral or written agreement.

WILBUR-ELLIS logo, Ideas to Grow With, FOLI-GRO, R-11, SYL-COAT, and Total Nutrition System are registered trademarks, and FOLI-GRO logo is a trademark of Wilbur-Ellis Company LLC. The OMRI Listed Seal is a registered trademark of OMRI. K-083018

Manufactured by: WILBUR-ELLIS COMPANY LLC
7 E. Washington Ave., Yakima, WA 98903

LOT#

NET CONTENTS: 2.5 GALLONS (9.46 Liters)

WEIGHT PER GALLON: 10.4 POUNDS at 68°F