

Fruit Tree, Vegetable & Ornamental Fungicide

For Control of a Broad Spectrum of Vegetable, Fruit and Ornamental Plant Diseases



Active Ingredient:

Chlorothalonil (tetrachloroisophthalonitrile) 29.6%
 Other Ingredients: 70.4%
 Total: 100.0%

Contains 2.91 Pounds Chlorothalonil Per Gallon

EPA Reg. No. 60063-9-54705

EPA Est. No. 48498-CA-1

Attention: This product contains chlorothalonil which is a chemical known to the State of California to cause cancer.

NET CONTENTS: 1 Pint / 473.16 mL

Manufactured for:

LAWN AND GARDEN PRODUCTS, INC.

P. O. Box 35000 • Fresno, CA 93745 • (559) 499-2100
 www.montereylawnandgarden.com

KEEP OUT OF REACH OF CHILDREN WARNING-AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. • Call a poison control center or doctor for further treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have affected person sip a glass of water if able to swallow. • Do not induce vomiting unless told by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	

Emergency phone numbers	(800) 858-7378 NPIC (human and animal health) (800) 424-9300 CHEMTREC (transportation and spills)
NOTES TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Persons having a temporary allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids.	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING - AVISO: May be fatal if inhaled. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Do not breathe spray mist. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

Personal Protective Equipment (PPE):

Mixers, loaders, applicators and all other handlers must wear:

- Long-sleeved shirt and long pants;
- Shoes plus socks;
- Protective eye wear;
- Chemical resistant gloves (some of the materials that are chemical-resistant to this product are barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyethylene, polyvinyl chloride, or viton; If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart);
- A NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any N, R, P, or HE prefilter.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Keep children and pets off treated area until dry.

Environmental Hazards

This product is toxic to aquatic invertebrates and wildlife. DO NOT apply directly to water or to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. DO NOT contaminate water when disposing of equipment washwater or rinsate.

Chlorothalonil can contaminate surface water through spray drift. DO NOT apply when weather conditions favor drift from treated areas. Under some conditions, it may also have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

Chlorothalonil degradates are known to leach through soil into ground water under certain conditions as a result of label use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

DIRECTIONS FOR USE

General Precautions and Restrictions

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT allow people or pets to enter treated areas until sprays have dried or dusts have settled.

DO NOT apply this product in a way that will contact other persons, or pets, either directly or through drift. Only protected handlers may be in the area during application.

DO NOT apply this product through any type of irrigation system.

Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, recreational park athletic fields, athletic fields located on or next to schools (ie., elementary, middle and high schools), campgrounds, churches, and theme parks.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use in small vegetable or fruit gardens, on plants intended for aesthetic purposes or climatic modification and being grown in exterior plantscapes, ornamental gardens or parks.

READ ENTIRE LABEL AND USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Mixing and Applying

FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE is a liquid concentrate that readily mixes with water and can be used in a tank-type manually operated or power sprayer.

FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE is highly effective for the control of a broad spectrum of vegetable, fruit and ornamental plant diseases when it is used according to the directions on this label. Thorough uniform coverage of plant surfaces is essential for good disease control.

Shake well before using. Mix appropriate amount with water and use at once.

GARDEN VEGETABLE DISEASES

Apply FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE at the rate indicated for each garden crop for the disease to be controlled, by mixing in one gallon of water to cover approximately 200 square feet (10 ft. x 20 ft.) of planted crop. Repeat applications at intervals specified. Use the 7 day schedule during severe disease conditions or wet weather. Do not apply closer to harvest than the indicated **Preharvest Interval (PHI)** for each type of crop. Do not apply more than the specified amount of FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE per 200 sq.ft. of garden area per year for each crop type.

Note: Avoid spraying plants during extremely hot and sunny weather. Do not apply to wilted plants. Do not graze treated areas or feed treated plant parts to livestock.

tsp. = teaspoonful (1/6 fluid ounce)

Crop	PHI (days)	Diseases	Rate Per 200 Sq. Ft.	Application Directions
Bean (Snap)	7	Rust Botrytis blight (gray mold)	2 ¾ tsp. per application 11 tsp. per year	Begin applications during early bloom stage or when disease first threatens and repeat at 7 day intervals.
Beans (Dry)	14	Rust Anthracnose Downy mildew Cercospora leaf spot (blackeye only) Ascochyta blight	2 tsp. per application 8 tsp. per year	Begin applications during early bloom stage and repeat at 7 to 10 day intervals. For use only on beans to be harvested dry with pods removed.
Cabbage Chinese Cabbage (tight-Headed varieties only) Cauliflower Broccoli Chinese Broccoli Brussels Sprouts	7	Alternaria leaf spot Downy mildew Ring spot	2 tsp. per application 15 tsp. per year	Begin applications after transplants are set or when conditions favor disease development. Repeat at 7 to 10 day intervals.
Carrot	0	Cercospora (Early) blight Alternaria (Late) blight	2 tsp. per application 18 tsp. per year	Start applications when disease threatens and repeat at 7 to 10 day intervals.
Celery	7	Cercospora (Early) blight Septoria (Late) blight Basal stalk rot (<i>Rhizoctonia solani</i>)	2 ¾ tsp. per application 22 tsp. per year	Start applications shortly after crop emergence or when transplants are set. Re-apply at 7 day intervals

Corn (sweet)	14	Helminthosporium leaf blights Rust	2 tsp. per application 11 tsp. per year	Begin applications when conditions favor disease development and repeat at 7 day intervals.
Cucurbits: Cucumber Cantaloupe Muskmelon Honeydew melon Watermelon Squash Pumpkin	0	Anthracnose Downy mildew Target spot Cercospora leaf spot Gummy stem blight (black rot) Alternaria leaf blight Scab Powdery mildew <i>Sphaerotheca only</i>)	2 ¾ tsp. per application 19 tsp. per year	Begin applications when plants are in first true leaf stage or when conditions are favorable for disease development. Repeat applications at 7 day intervals. Note: Spraying mature watermelons may result in sunburn of the upper surface of the fruit. DO NOT apply FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE to watermelons when any of the following conditions are present: 1. Intense heat and sunlight; 2. Drought conditions; 3. Poor vine canopy; 4. Other crop and environmental conditions which may be conducive to increased natural sunburn. DO NOT combine FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE with anything except water for application to watermelons unless your prior use has shown the combination to be non-injurious to watermelons under your conditions of use.
Onion (dry bulb), Garlic	7	Botrytis leaf blight or blast Purple blotch	2 ¾ tsp. per application 18 tsp. per year	Apply as follows: Low Disease Hazard & Prior to Infection: 1 tsp. Frequency: 10 days. Low Disease Hazard & Some Disease Present: 2 tsp. Frequency: 7 to 10 days. High Disease Hazard: 2 ¾ tsp. Frequency: 7 days.
Onion (green) bunching), Leek Shallot	14 (green onion, leek, shallot)	Botrytis leaf blight or blast Purple blotch Downy mildew (suppression)	2 ¾ tsp. per application 8 tsp. per year	Begin applications prior to favorable infection periods, and repeat at 7 to 10 day intervals for as long as conditions favor disease. Use the high rate and a 7 day schedule of applications when heavy dew or rain persist. Do not apply more than 3 times per season. If additional disease control is needed before harvest, use another registered fungicide.
Parsnip	10	Alternaria leaf spot Downy mildew Anthracnose Botrytis blight (gray mold) Bottom rot (Rhizoctonia)	2 tsp. per application 7 tsp. per year	Make the first application at the first sign of disease or when conditions are favorable for infection. Continue applications on a 7 to 10 day schedule.
Potato	7	Late blight Early blight Botrytis vine rot	1 ½ tsp. per application 14 tsp. per year	Begin applications at the low rate when vines are first exposed and leaf wetness occurs. Repeat applications at 7 to 10 day intervals. Use the highest rate and shortest interval when plants are rapidly growing and disease conditions are severe.

Tomato	0	<p>FOLIAGE (apply every 7-10 days):</p> <p>Early blight Late blight Gray leaf spot Gray leaf mold Septoria leaf spot Target spot</p> <p>FRUIT (apply every 7-14 days beginning at fruit set):</p> <p>Anthrachnose Alternaria fruit rot (black mold) Botrytis gray mold Late blight fruit rot Rhizoctonia fruit rot</p>	<p>2 ½ tsp. per application</p> <p>18 tsp. per year</p>	<p>Begin applications when dew or rain occur and disease threatens. Use the highest rate and shortest interval specified when disease conditions are severe.</p>
--------	---	--	---	--

Cherry Plum Prune	split		per year	mid to late winter before budswell.
		<p>Shothole Brown rot blossom blight Lacy (russet) scab on prune Cherry leaf spot Scab</p>		<p>Make one application at budbreak or popcorn (pink, red or early white bud). If weather conditions favor disease, make a second application 10 days later (full bloom to petal fall). Apply at shuck split to prevent infections on young fruit. If additional disease control is needed after shuck split and before harvest, use another registered fungicide.</p>

ORNAMENTAL GARDEN PLANTS

Apply FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE at a rate of 2 ¼ teaspoons per gallon, or 3 Tablespoons (1½ fluid ounces) per 4 gallons of water unless other directions are given in the tables below. Apply enough diluted spray to provide thorough coverage of all plant parts that are intended to be protected from disease, generally ranging from ½ to 3 gallons per 1,000 square feet of garden area. Repeat applications at 7 to 14 day intervals until conditions are no longer favorable for disease. During periods when conditions favor severe disease incidence, generally cloudy or wet weather, apply FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE at 7 day intervals. **DO NOT apply more than a total of 37 fluid ounces of FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE (0.84 lbs chlorothalonil active ingredient) per 1,000 sq. ft. per growing season.**

Fruits and other structures which may be borne on treated plants **MUST NOT BE EATEN.**

This product may be used in greenhouses. DO NOT use mist blowers or high pressure spray equipment when making applications of this product in greenhouses.

FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE is recommended for control of fungal diseases referred to by numbers in parentheses following each type of ornamental plant. The user should test for possible phytotoxic responses, using recommended rates on each type of ornamental plant on a small area prior to widespread use. Applications made during bloom may damage flowers and/or fruits.

FRUIT AND NUT TREE DISEASES

Apply this product in sufficient water to obtain uniform coverage of tree canopy, including the undersides of leaves, to the point of spray runoff. Mix the indicated rate of FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE into each 4 gallons of water to be used for each application, generally sufficient to treat 200 square feet.

Do not apply closer to harvest than the indicated **Preharvest Interval (PHI)** for each type of crop.

Note: Avoid spraying plants during extremely hot and sunny weather. **DO NOT apply FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE within one week before or after application of oil or an oil-based pesticide.** Do not graze treated areas or feed treated plant parts to livestock.

tsp. = teaspoonful (1/6 fluid ounce)

Crop	PHI (days)	Diseases	Rate Per 4 Gallons	Application Directions
Blueberry	42	Mummy Berry Anthrachnose	3 ½ tsp. per application 10 ½ tsp. per year	Begin applications at budbreak (green tip). Repeat applications until early bloom at 10 day intervals. DO NOT apply after early bloom, otherwise phytotoxicity may occur to the developing fruit.
Filberts (Hazelnuts)	120	Eastern filbert blight	3 ½ tsp. per application 10 ½ tsp. per year	Begin applications at leaf bud break and repeat at 2 to 4 week intervals. Based on available residue data, use of this product on filberts is restricted to Oregon.
Papaya	14	Alternaria fruit spot Anthrachnose Stem end rot	2 ¼ tsp. per application 8 tsp. per year	Begin treatment when conditions favor development of disease and continue treatments at 14 day intervals until weather conditions no longer favor disease development.
Passion Fruit (Hawaii only)	7	Alternaria fruit and leaf spot (brown spot)	2 tsp. per application 9 tsp. per year	Apply in sufficient water to obtain adequate coverage of fruit and leaves. Begin applications before fruit spots appear (April to July) and re-apply at 14 day intervals until weather conditions no longer favor disease development.
Stone Fruits: Peach Nectarine Apricot	Do not apply after shuck	Leaf curl	3 ¼ tsp. per application 18 ¾ tsp.	Apply at leaf fall in late autumn. When conditions favor high disease levels apply once or twice more in

Ornamentals Recommended For Treatment With FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE

Broadleaf Shrubs and Trees	
<p>Andromeda (<i>Pieris</i>) (4) Ash (<i>Fraxinus</i>) (1) Aspen (1) Azalea (1,2,4) Buckeye, Horsechestnut (1) Camellia (2) Cherry-laurel (1) Crabapple (1,6) Dogwood (1) Eucalyptus (3) Euonymus (1) Firethorn (<i>Pyracantha</i>) (1) Flowering almond (1,2) Flowering cherry (1,2) Flowering peach (1,2) Flowering plum (1,2) Flowering quince (1,2) Hawthorn (1,6)</p>	<p>Holly (1) Lilac (5) Magnolia (1) Maple (1) Mountain laurel (1) Oak (red group only) (1,7) Oregon-grape (<i>Mahonia</i>) (6) Red-tip (<i>Photinia</i>) (1) Poplar (1) Privet (<i>Ligustrum</i>) (1) Rhododendron (1,2,4) Sand cherry (1,2) Sequoia (1) Spirea (1) Sycamore, Planetree (1) Viburnum (5) Walnut (<i>Juglans</i>) (1)</p>

Flowering ^a Plants and Bulbs	
Arabian violet (2) Begonia (1) Carnation (1,2) Chrysanthemum (1,2) Crocus (1) Daffodil (1) Daisy (1) Geranium (1,6) Gladiolus (1,2) Hollyhock (6) Hydrangea (foliage only) (1,6) Iris (1,2)	Lily (1) Marigold (1) Narcissus (1) Pansy (1) Petunia (1,4) Phlox (1) Poinsettia ^b (1) Rose ^c (1) Statice (1) Tulip (1) Zinnia (1,5)

a/ Avoid applications during bloom period on plants where flower injury is unacceptable.

b/ Discontinue applications prior to bract formation; phytotoxicity is possible on the bracts.

c/ Use 1½ Tbs. of FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE per 4 gallons of water.

Foliage Plants	
Aglaonema (1) Areca palm (1) Artemesia (1) Boston fern (<i>Nephrolepis</i>) (1) Dracaena (1) Dumbcane (<i>Dieffenbachia</i>) (1) Fatsia (<i>Aralia</i>) (1) Ficus (1) Florida ruffle fern (1) Leatherleaf fern (1)	Lipstick plant (1) Ming aralia (1) Oyster plant (<i>Rhoeo</i>) (1) Pachysandra ^d (1) Parlor palm (<i>Chamaedorea</i>) (1) Peperomia (1) Philodendron (1,4) Prayer plant (<i>Maranta</i>) (1) Syngonium (1) Zebra plant (<i>Aphelandra</i>) (1)

d/ Use 4 Tbs. of FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE per 4 gallons of water.

Diseases controlled with FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE

1. Leafspots & Foliar Blights:	
Actinopelte leafspot Alternaria leafspot or leaf blight Anthracnose (<i>Gnomonia</i> , <i>Glomerella</i> , <i>Colletotrichum</i> , <i>Discula</i>) blights Black spot (<i>Diplocarpon</i>) Botrytis blights Cephalosporium leafspot Cercospora leafspot Cercosporidium leafspot Shothole (<i>Stigmina</i>) Corynespora stem & leafspots Curvularia leafspot Dactylaria leafspot Didymellina leafspot Drechslera (<i>Bipolaris</i>) leafspots, inkspot	Fabraea (<i>Entomosporium</i>) leafspot Fusarium (<i>Gibberella</i>) leafspot Gloeosporium black leafspot Marssonina leafspot Monilinia blossom blight, twig blight Mycosphaerella ray blight Myrothecium leafspot, brown rot Phyllosticta leafspot Ramularia leafspot Rhizoctonia web blight Scab (<i>Venturia</i>) Septoria leafspot Sphaeropsis leafspot Stagonospora leaf scorch Tan leafspot (<i>Curvularia</i>) Volutella leaf blight

2. Flower Spots & Blights:	
Botrytis flower spot, flower blight Curvularia flower spot Monilinia blossom blight	Ovulinia flower blight Rhizopus blossom blight Sclerotinia flower blight

3. Cylindrocladium stem canker	

4. Phytophthora leaf blight, dieback	

5. Powdery mildews:	
<i>Erysiphe cichoracearum</i> <i>Sphaerotheca fuliginea</i>	<i>Microsphaera</i> spp.

6. Rusts:	
<i>Gymnosporangium</i> spp. <i>Pucciniastrum hydrangeae</i>	<i>Puccinia</i> spp.

7. Taphrina blister	

CONIFER DISEASES

Apply this product in sufficient water to obtain uniform coverage of tree canopy, including the undersides of leaves, to the point of spray runoff. Mix the indicated rate of FRUIT TREE, VEGETABLE & ORNAMENTAL

FUNGICIDE into each 4 gallons of water to be used for each application, generally sufficient to treat 200 square feet.

Do not apply more than 16 fluid ounces (100 teaspoonfuls) of FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE per 1,000 square feet per year on conifers.

Note: Avoid spraying plants during extremely hot and sunny weather. **DO NOT apply FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE within one week before or after application of oil or an oil-based pesticide.** Do not graze treated areas or feed treated plant parts to livestock.

tsp. = teaspoonful (1/6 fluid ounce)

Diseases	Rate Per 4 Gallons	Application Directions
Scleroderris canker (pines) Swiss needlecast Sirococcus tip blight Rhizosphaera needlecast (spruces) Scirrhia brown spot (pines)		Make the first application in spring when new shoot growth is 1/2 to 2 inches in length. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development.
Cyclaneusma and Lophodermium needlecasts (pines)	5 tsp. per application	Apply in early spring prior to budbreak. Repeat applications at approximately 6 to 8 week intervals, until spore release ceases in late fall. Apply monthly during periods of frequent rainfall, and where Lophodermium infections occur during dormancy (Pacific Northwest). During drought periods, applications may be suspended, then resumed upon next occurrence of needle wetness.
Rhabdocline needlecast (Douglas-fir)		Apply at budbreak and repeat at 3 to 4 week intervals until needles are fully elongated and conditions no longer favor disease development. When irregular budbreak occurs, apply weekly until all trees have broken bud, then every 3 to 4 weeks as specified above.
Botrytis seedling blight, Phoma twig blight		Begin applications when seedlings are 4 inches tall and when cool, moist conditions favor disease development. Make additional applications at 7 to 14 day intervals as long as disease favorable conditions persist.
Autoecious needle rust (Weir's cushion rust) (spruces)		Begin applications when 10% of buds have broken and repeat twice thereafter at 7-10 day intervals.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Pesticide Storage: Store in a cool place. Protect from excessive heat.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY AND LIMITATION OF DAMAGES

CONDITIONS OF SALE: To the extent consistent with applicable law, Lawn and Garden Products, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Lawn and Garden Products, Inc. **LAWN AND GARDEN PRODUCTS, INC. DISCLAIMS ALL OTHER WARRANTIES, EXPRESS OR IMPLIED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LAWN AND GARDEN PRODUCTS, INC. SHALL**

NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, AND LAWN AND GARDEN PRODUCTS, INC.'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE. BUYER AND USER ACKNOWLEDGE AND ASSUME ALL RISKS AND LIABILITY RESULTING FROM HANDLING, STORAGE AND USE OF THIS PRODUCT. LAWN AND GARDEN PRODUCTS, INC. DOES NOT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTY, GUARANTEE OR REPRESENTATION CONCERNING THIS PRODUCT.

0704/0810(02)