

**RESTRICTED USE PESTICIDE  
Due to Acute Toxicity**

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.



**Section 24(c) Special Local Need Label**

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF NORTH CAROLINA**

**Gramoxone® SL 2.0  
For Suppression and/or Control of Palmer Amaranth in Peanut**

**EPA Reg. No. 100-1431  
EPA SLN No. NC-120004**

**This label expires and must not be distributed or used in accordance with this SLN  
registration after December 31, 2022**

**KEEP OUT OF REACH OF CHILDREN**



**DANGER / POISON  
PELIGRO / VENENO**

**DIRECTIONS FOR USE**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.
- **FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.**

**Application Method**

Gramoxone SL 2.0 can be applied through recirculating rope wick or carpet roller equipment for suppression or control of glyphosate-resistant Palmer amaranth or to prevent seed production.

Make application at least 6 inches above the peanut canopy, and set application equipment to avoid dripping. Low ground speed enhances coverage and hence weed control. Speeds less than 5 mph are recommended.

Control of large weeds is enhanced if application is made in late afternoon or early evening.

Follow application equipment manufacturer's instructions.

### **Mixing Instructions and Application Rate**

Mix 1 part of Gramoxone SL 2.0 with 1-1.5 parts of water to prepare a 40-50% solution. Add nonionic surfactant containing 75% or more surface-active agent at a minimum of 0.25% v/v (2 pt/100 gal or 0.32 fl oz/gal) of the finished volume.

Adjust equipment to apply up to 2 pt/A of the herbicide-water mixture.

### **Restrictions**

1. Do not apply more than 1 pt/A of Gramoxone SL 2.0 using the recirculating rope wick or carpet roller application method.
2. Do not exceed 0.5 lb ai/A (1 qt/A) total for the entire growing season from all applications on this or any other supplemental or Section 3 Federal label for peanut.
3. Make applications of Gramoxone SL 2.0 as early as possible, and not less than 30 days before harvest to control weeds that may interfere with harvesting operations.
4. Do not allow livestock to graze in treated areas.
5. Do not feed hay or threshings from treated fields to livestock.

---

©2017 Syngenta

Gramoxone® and the Syngenta logo are trademarks of a Syngenta Group Company

24(c) Registrant:  
Syngenta Crop Protection, LLC  
P. O. Box 18300  
Greensboro, NC 27419-8300

Label Code: NC1431033CA0817