

---

## SUPPLEMENTAL LABELING

---

**Syngenta Crop Protection, LLC**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300  
SCP 620A-S3 0118

PROMETRYN	GROUP	<b>5</b>	HERBICIDE
-----------	-------	----------	-----------

### Caparol® 4L

Herbicide

**This supplemental label expires on 01/31/2021 and must not be used or distributed after this date.**

Active Ingredient:

Prometryn: 2,4-bis(isopropylamino)-  
6-(methylthio)-s-triazine .....

44.4%

Other Ingredients:

55.6%

Total:

100.0%

Caparol 4L contains 4 lb active ingredient per gallon.

Shake well before using.

**KEEP OUT OF REACH OF CHILDREN.**

## CAUTION

EPA Reg. No. 100-620

All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Caparol 4L as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**syngenta**

## **DIRECTIONS FOR USE**

### **Winter Weed Control**

#### **Winter and Early Spring Weed Control in AL, AR, LA, MO, MS, TN, and VA**

For control of winter and early spring germinating annual weeds (including henbit, common chickweed, sibara, and Palmer amaranth), apply 1.5-2 pt of Caparol 4L after bedding (e.g., stale seedbed) from November 1 until 14 days before planting cotton. Use the 2.0 pt/A rate for applications made in Nov. or Dec. Use the 1.5 pt/A rate for applications made from Jan. 1 to 30 days before cotton planting. Applications may be made before or after weeds emerge. For control of emerged weeds, preferably less than 2 inches in height, add a suitable and approved crop oil concentrate or surfactant according to its label. In the event weeds exceed 2 inches in height at the time of treatment, apply Caparol 4L in tank mixture with a contact herbicide (e.g., solo glyphosate or Gramoxone SL 2.0). Refer to the label of the contact herbicide for rates of application, additives, and for weed height restrictions at time of application. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

After applying Caparol 4L, do not mechanically till the seedbed prior to the cotton planting process, as this will encourage germination of weed seeds.

Follow with a preemergence herbicide program for cotton. In the event that a subsequent application of Caparol 4L is made, do not exceed the total rate of Caparol 4L that may be applied to a single cotton crop.

Caparol® and the Syngenta logo are trademarks of a Syngenta Group Company

©2018 Syngenta