

Supplemental Labeling



Dow AgroSciences

Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

FIFRA 24(c) Special Local Need Label (SLN)

For Distribution and Use Only in the State of California

For control or suppression of foxglove and lettuce aphid in *Brassica* (cole) leafy vegetables, leafy vegetables (except *Brassica*) and Watercress.

Sequoia™

EPA Reg. No. 62719-623

SLN CA-170004

This label expires and must not be distributed or used in accordance with this SLN registration after March 31, 2022.

Environmental Hazard

This product is highly toxic to bees exposed through contact during spraying and while spray droplets are still wet. Do not apply this product or allow it to drift to blooming crops or weeds while bees or other pollinating insects are actively foraging the treatment area. This product may be toxic to bees exposed to treated foliage for up to 3 hours following application. Toxicity is reduced when spray droplets are dry.

Risk to managed bees and native pollinators from contact with pesticide spray or residues can be minimized when applications are made before 7:00 am or after 7:00 pm local time or when the temperature is below 50° F at the site of application.

Refer to the Directions for Use for crop specific restrictions and additional advisory statements to protect pollinators.

This product is toxic to aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Directions for Use

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This state-specific Section 24(c) labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, and precautions on the EPA registered label for Sequoia, EPA Reg. No. 62719-623 and this label.

Refer to product label for General Use Precautions, Mixing and Application instructions.

Brassica (Cole) Leafy Vegetables (Crop Group 5)¹

¹*Brassica* (cole) leafy vegetables (crop group 5) including broccoli, broccoli raab, Brussels sprouts, cabbage, cauliflower, cavalo, Chinese broccoli (gia lon), Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, rape greens, white flowering broccoli

Pests and Application Rates:

Pests	Sequoia (fl oz/acre)
foxglove and lettuce aphids	1.5 – 2.0

	(0.023 – 0.031 lb ai/acre)
--	----------------------------

Application Timing: Treat in accordance with local economic thresholds. Consult your Dow AgroSciences representative, cooperative extension service, certified crop advisor or state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: Use a higher rate in the rate range for heavy pest populations.

Restrictions:

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 7 days apart.
- Do not exceed a maximum 2 fluid ounces of product per application.
- Do not make more than four applications per crop.
- Do not make more than two consecutive applications per crop.
- Do not apply to crops grown for seed.
- Do not apply more than a total of 8 fl oz of Sequoia (0.124 lb ai of sulfoxaflor) per acre per year.
- If blooming vegetation is present 12 feet out from the downwind edge of the field, a downwind 12-foot on-field buffer must be observed.

Leafy Vegetables (Except *Brassica*) (Crop Group 4)¹ and Watercress

¹Leafy vegetables (except *Brassica*) (crop group 4) including amaranth, arugula, cardoon, celery, celtuce, chervil, Chinese celery, Chinese spinach, corn salad, cos (romaine), dandelion, dock, edible-leaved chrysanthemum, endive (escarole), finocchio, Florence fennel, garden cress, garden purslane, garland chrysanthemum, head lettuce, leaf lettuce, leafy amaranth, New Zealand spinach, orach, parsley, radicchio (red chicory), rhubarb, spinach, sweet anise, sweet fennel, Swiss chard, tampala, upland cress, vine spinach, winter cress, winter purslane, yellow rocket

Pests and Application Rates:

Pests	Sequoia (fl oz/acre)
foxtail and lettuce aphids	1.5 – 2.0 (0.023 – 0.031 lb ai/acre)

Application Timing: Treat in accordance with local economic thresholds. Consult your Dow AgroSciences representative, cooperative extension service, certified crop advisor or state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: Use a higher rate in the rate range for heavy pest populations.

Restrictions:

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 7 days apart.
- Do not exceed a maximum 2 fluid ounces of product per application.
- Do not make more than four applications per crop.
- Do not make more than two consecutive applications per crop.
- Do not apply to crops grown for seed.
- Do not apply more than a total of 8 fl oz of Sequoia (0.124 lb ai of sulfoxaflor) per acre per year.
- If blooming vegetation is present 12 feet out from the downwind edge of the field, a downwind 12-foot on-field buffer must be observed.

Registrant: Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268
1-800-992-5994

®™ Trademark of The Dow Chemical Company (“Dow”) or an affiliated company of Dow