



Bayer CropScience

Bayer CropScience LP
 P.O. Box 12014
 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina 27709
 1-866-99BAYER (1-866-992-2937)

REASON® 500 SC Foliar Fungicide
 EPA Reg. No. 264-695
 plus
SCALA™ brand SC Fungicide
 EPA Reg. No. 264-788

For control of early season potato diseases
 in tank-mix with reduced rates.

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF CALIFORNIA, COLORADO, CONNECTICUT, DELAWARE, FLORIDA, IDAHO, MAINE, MARYLAND, MICHIGAN, MINNESOTA, MONTANA, NEBRASKA, NEW JERSEY, NORTH DAKOTA, OHIO, OKLAHOMA, OREGON, PENNSYLVANIA, RHODE ISLAND, VERMONT, VIRGINIA, WASHINGTON, WEST VIRGINIA, WISCONSIN.

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. It is a violation of federal law to use this product in a manner inconsistent with its label. This product bulletin must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling.

GENERAL DIRECTIONS

<p>Early blight (including strobilurin-resistant strains)</p>	<p>At or before row closure: Apply Reason 500 SC at 4.0 fl oz/acre in tank mix with Scala SC at 7.0 fl oz/acre (preventative late blight rate for Reason 500 SC).</p>
<p>Brown spot Botrytis leaf spot Black dot (suppression) Late blight</p>	<p>After row closure: Apply Reason 500 SC at 5.5-8.2 fl oz/acre in tank mix with Scala SC at 7.0 fl oz/acre. Applications should be made on a 7- to 10-day interval</p>

RESTRICTIONS:

Follow all mix-partner labeling instructions. A maximum of 24.6 fl oz of Reason 500 SC may be applied per growing season. Applications of Reason 500 SC can be made up to 14 days before harvest. Do not apply more than 35 fl oz/acre of Scala SC per crop. Do not apply Scala SC within 7 days of harvest. For resistance management, alternate each QoI-containing application with a non-QoI containing fungicide.

APPLICATION:

Apply using ground equipment, chemigation, or by air. Under conditions favorable for disease development, shorten the spray intervals and/or use higher registered rates for improved control.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937) or

Visit our worldwide web site at www.bayercropscienceus.com

*As with any crop-protection product, always read and follow label directions.
 For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*