
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1533A-S1 1116

Segovis®

GROUP	U15	FUNGICIDE
--------------	------------	------------------

Fungicide

- For control of diseases in vegetable transplants (produced for sale only to residential consumers) grown in greenhouses, outdoor growing structures (including shade houses, lath houses and other outdoor growing structures) and nurseries

This supplemental label expires on 11/14/2019 and must not be used or distributed after this date.

Active Ingredient:	
Oxathiapiprolin (CAS No. 1003318-67-9).....	18.7%
Other Ingredients:	81.3%
Total:	100.0%

KEEP OUT OF REACH OF CHILDREN.

EPA Reg. No. 100-1533

All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed. Before using Segovis as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

The Syngenta logo, featuring the word "syngenta" in a lowercase, sans-serif font with a stylized leaf icon above the letter "a".

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

VEGETABLE TRANSPLANT USE DIRECTIONS

Vegetable transplants can be treated with Segovis prior to retail sale only to residential consumers.

Basil, fresh (Foliar Diseases)

Target Disease	Rate	Application Timing	Use Directions
Downy mildew (<i>Peronospora belbahrii</i>)	4 – 8 ml/5,000 sq ft (1.1 – 2.4 fl oz/A)	Begin foliar applications prior to disease development.	Use the higher application rate when disease is present, Apply in at least 15 gallons per acre, increasing the spray volume as the plants grow larger to ensure thorough coverage of the foliage.
Resistance Management:			
<ul style="list-style-type: none"> • Make no more than 2 applications per crop. • Refer to Section 3.2 for additional information. 			
USE RESTRICTIONS			
<ol style="list-style-type: none"> 1) Maximum Single Application Rate: 2.4 fl oz/A 2) Minimum Application Interval: 5 days 3) Maximum Annual Rate: <ol style="list-style-type: none"> a) Plants Grown Outdoors in Nurseries (including outdoor growing structures): <ol style="list-style-type: none"> i) Do not apply greater than 4.8 fl oz/A (0.0625 lb ai/A of oxathiapiprolin-containing products) to a single crop. ii) When multiple crops are produced in the same production area, do not apply greater than 0.25 lb ai/A/year of oxathiapiprolin-containing products. b) Plants Grown in Greenhouses: <ol style="list-style-type: none"> i) Do not apply greater than 4.8 fl oz/A (0.0625 lb ai/A of oxathiapiprolin-containing products) to a single crop. 			

Segovis® and the Syngenta logo are trademarks of a Syngenta Group Company
©2018 Syngenta