



For control of a wide-spectrum of annual, biennial, and perennial broadleaf weeds and brush in Pastures, Rangeland, and Grass (Hay, Silage); Grass Grown for Seed; Wheat; Sugarcane, Teff, Conservation Reserve Program Land; General Farmstead Areas; Post-Harvest, Fallow, Crop Stubble and Set Aside Acres

ACTIVE INGREDIENT(S):

| | |
|---------------------------------------|---------|
| 3,6-dichloromethoxybenzoic acid | 18.28% |
| 2,4-Dichlorophenoxyacetic acid | 24.62% |
| OTHER INGREDIENTS: | 57.10% |
| TOTAL | 100.00% |

Equivalent to: Dicamba Acid, 1.8 lbs./gal. and 2,4-D Acid, 2.4 lbs./gal.
 Isomer specific by AOAC Method 6.D01-5 (12th Ed.)
 Patent No. 5,877,112, other patents pending.

**KEEP OUT OF REACH OF CHILDREN
 DANGER – PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you do not understand the label, find someone to explain it you in detail.)

| FIRST AID | |
|--|--|
| IF IN EYES: | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| IF SWALLOWED: | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious or convulsing person. |
| IF ON SKIN OR CLOTHING: | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for further treatment advice. |
| HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information. | |
| NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. | |

EPA Reg. No. 5905-564
 EPA Est. No. 42750-MO-001
 AD 040417

NET CONTENTS:

- 1 Gallon (3.785 Liters)
- 2.5 Gallons (9.46 Liters)
- 55 Gallons (208.18 Liters)
- 250 Gallons (946.1 Liters)

MANUFACTURED FOR:
HELENA CHEMICAL COMPANY
 225 SCHILLING BOULEVARD, SUITE 300
 COLLIERVILLE, TN 38017

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER**

Corrosive. Causes irreversible eye damage. Harmful if swallowed. Harmful if absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All mixers, loaders, applicators, flaggers, and other handlers must wear:

1. Long-sleeved shirt and long pants
2. Shoes and socks
3. Chemical resistant gloves (made of Barrier laminate, butyl rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyethylene, polyvinyl chloride (PVC) \geq 14 mils, or viton \geq 14 mils)
4. Chemical resistant apron when applying with any handheld nozzle or equipment, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate
5. Protective eyewear (goggles or face shield)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

See "Engineering Controls" for additional requirements.

If this container contains over 1 gallon and less than 5 gallons, mixers and loaders who do not use a mechanical system (probe and pump) to transfer the contents of this container must wear coveralls or a chemical-resistant apron in addition to the other required PPE.

ENGINEERING CONTROL STATEMENTS

Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)].

When handlers use enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and change into clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate. Apply this product only as directed on label.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Groundwater Contamination: Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce

the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Endangered Species Concerns: The use of any pesticide in a manner that may kill or otherwise harm and endangered species or adversely modify their habitat is a violation of federal law.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Unless otherwise directed in supplemented labeling, all applicable directions, restrictions, precautions and Conditions of Sale and Warranty are to be followed. This labeling must be in the user's possession during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

1. Coveralls worn over short-sleeved shirt and short pants.
2. Chemical resistant footwear plus socks
3. Chemical resistant gloves made of any waterproof material.
4. Chemical resistant headgear for overhead exposure.
5. Protective eyewear

Notify workers of application by warning them orally and by posting warning signs at entrances to treated area.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

USE REQUIREMENTS FOR PASTURES, PERENNIAL GRASSLANDS, GRASS GROWN FOR SEED, RANGELAND, FALLOW LAND AND NONCROP AREAS: Do not enter treated areas until spray has dried. For early entry to treated areas, wear eye protection, chemical-resistant gloves made of any waterproof material, long-sleeved shirt, long pants, shoes and socks.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Do not store under conditions that might adversely affect the container or its ability to function properly.

PESTICIDE STORAGE: Do not store below temperature of 32°F or above 100°F. Store in original container in a well-ventilated area separately from fertilizer, feed, and foodstuffs. Keep container tightly closed when not in use. Reduce stacking height where local conditions can affect package strength.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Wastes resulting from this product must be disposed of on site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

(non-refillable ≤5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(non-refillable >5 gallons): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Pressure rinse as follows (all sizes): Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use for disposal. Insert pressure rinsing nozzle inside of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

[Optional text for refillable plastic bulk containers]

Refillable container (110 & 250 gallon & bulk): Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing process two more times.

In Case of Spill: In case of large-scale spillage regarding this product, call ChemTrec 1-800-424-9300.

Steps to be taken in case material is released or spilled: Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal. Remove contaminated clothing, and wash affected skin areas with soap and water. Wash clothing before re-use. Keep the spill out of all sewers and open bodies of water.

I. PRODUCT INFORMATION

LATIGO® is a postemergence herbicide for controlling a wide spectrum of annual, biennial, and perennial broadleaf weeds and brush in pastures, rangeland, and grass (hay, silage); grass grown for seed; sugarcane; teff; wheat; conservation reserve program land; postharvest, fallow, crop stubble, set-aside acres; general farmstead areas; certain noncrop areas; and for forest management.

Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order of Injunctive Relief in Washington Toxics Coalition et al vs. EPA CO1-132C (W.D.WA.). For information, please refer to www.epa.gov/espp/wtc/.

Mode of Action

LATIGO® contains two active ingredients uniquely formulated to be used alone or tank mixed with other listed products as well as liquid fertilizer solutions. LATIGO® is readily absorbed by plants through shoot and root uptake, translocates throughout the plant's system, and accumulates in areas of active growth. LATIGO® interferes with the plant's growth hormones (auxins) resulting in death of many broadleaf weeds.

For best results, thoroughly clean sprayer equipment (tank, lines and nozzles) immediately after use by flushing system with water and heavy duty detergent or other suitable tank cleaner.

II. APPLICATION INSTRUCTIONS

Apply LATIGO® at the rates and growth stages listed in Tables 1 and 2 as follows unless instructed differently by section on “Food/Feed Crop Specific Information” or “Non-Food/Feed Use (Land not Harvested, Grazed or Foraged)-Specific Information.” LATIGO® may be applied using water or sprayable fluid fertilizer as a carrier. Sprayable fluid fertilizer may be used as the carrier in preplant or pre-emergence use for all crops listed on this label. Postemergence uses with sprayable fluid fertilizer may be made on pasture, hayland, or wheat crops only. The most effective application rate and timing varies based on the target weed species (refer to Table I). In mixed populations of weeds the correct rate is determined by the weed species requiring the highest rate. Delaying application permits weeds to exceed the maximum size and will prevent adequate control. For certain specified applications liquid fertilizer or oil may replace part or all of the water as diluent. If dry flowable (DF), wettable powder (WP) or flowable (F) tank mix products are to be used, these should generally be added to the spray tank first. Refer to the mixing directions on the labels of the tank mix products.

Irrigation: In irrigated areas, it may be necessary to irrigate before treatment to ensure active weed growth.

CHEMIGATION PROHIBITION

Do not apply this product through any type of irrigation system.

Spray Coverage:

Weeds must be thoroughly covered with spray. Dense leaf canopies shelter smaller weeds and prevent adequate spray coverage.

Sensitive Crop Precautions:

LATIGO® may cause injury to desirable trees and plants, particularly beans, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tobacco, tomatoes and other broadleaf plants when contacting their roots, stems or foliage. At high temperatures (about 85 degrees or higher), vapors from this product may cause injury to the aforementioned susceptible crops. These plants are most sensitive to LATIGO® during their development or growing stage. Do not treat areas where either possible downward movement into the soil or surface washing may cause contact of LATIGO® with the roots of desirable trees and shrubs.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a Coarse or coarser spray, apply only as a Coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for non-target species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to cotton, okra, flowers,

grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All aerial equipment and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

For aerial equipment, the boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications.

When applications are made in a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

For ground boom application, do not apply with a nozzle height greater than 4 feet above the crop canopy.

Table 1. Application Rate and Timing – Annual Weeds

(Applicators must follow maximum application rates in the Use Specific Information sections of the label.)

(For use in non-food/feed crops only: the addition of liquid fertilizer (28-0-0,32-0-0) solutions at ½ the GPA spray solution has shown to give increased efficacy.)

| Weeds Controlled (including ALS- and triazine-resistant) | Rate Per Acre (according to weed growth stage) | | | | | |
|--|--|-----------------|-----------|-------------|------------------------|------------------|
| | 1/3 pint | 2/3 pint | 1 pint | 1-1/8 pints | 1-2/3 pints | 2 pints |
| Beebalm, Spotted | - | - | - | pre-bloom | post-bloom | - |
| Broomweed | 1-3" | 3" branching | - | branching | - | after branching |
| Buckwheat, Wild | - | 1-6" | - | - | - | - |
| Buffalobur | - | - | - | 1-6" | - | Flowering |
| Burdock | - | pre-flower | - | - | - | - |
| Buttercup | - | pre-flower | - | early bloom | late bloom | - |
| Chickweed, Common | - | Seedling | 1-3" | - | - | - |
| Cockle, Cow | - | <3" | - | - | - | - |
| Cocklebur, Common | - | 1-6" | 6-12" | 12-18" | - | - |
| Coreopsis, Plains | 1-4" | 1-6" | - | - | - | - |
| Croton, Woolly | - | 4-12" | 12-30" | - | - | - |
| Dogfennel | - | - | - | 10-15" | - | - |
| Evening Primrose | - | <2" | - | 2-6" | - | - |
| Flax | - | <2" | - | - | - | - |
| Fleabane, Annual | - | 1-4" | 4-8" | 8" | - | - |
| Flixweed | - | <3" | - | - | - | - |
| Henbit | - | - | preflower | - | flower | - |
| Horseweed/Marestail | - | - | pre-bolt | - | post-bolt: up to 4" | - |
| Knotweed spp. | - | <3" runners | - | >3" runners | - | actively growing |
| Kochia | - | 1-6" | 6-10" | 10-20" | - | actively growing |

| | | | | | | |
|---|------|-----------------|-----------------|------------------|-----------------|------------------|
| Lambsquarters, Common | - | 1-6" | 6-10" | 10-20" | - | actively growing |
| Mallow, Common | - | <3" | - | - | - | - |
| Morningglory, Ivyleaf | - | pre-flower | - | - | - | - |
| Morningglory, Tall | - | pre-flower | - | post-flower | - | - |
| Mustards, Annual | | rosette | | early bolt | - | - |
| Mustards, Tansy | - | <3" | | | - | - |
| Pennycress, Field | - | - | - | rosette | - | - |
| Pepperweed, Virginia | - | - | 1-3" | 3-6" | after branching | - |
| Pigweed, Prostrate | - | <3" | - | - | - | - |
| Pigweed, Redroot | - | <3" | 3-10" | - | - | - |
| Pigweed, Smooth | - | <3" | - | - | - | - |
| Pigweed, Tumble | - | <3" | - | mature | - | - |
| Poorjoe | - | prior to flower | - | - | - | actively growing |
| Purslane, Common | | <3" | 3-8" | | - | - |
| Ragweed, Common | | | | >10" | - | - |
| Ragweed, Western & Lanceleaf | 1-3" | 3-6" | 6-10" | actively growing | - | - |
| Sedge ¹ | - | - | - | - | - | - |
| Shepherdspurse | - | rosette | - | - | - | - |
| Smartweed, Pennsylvania | - | <4" | - | - | 4-12" | - |
| Sneezeweed, Bitter | - | 1-4" | prior to flower | flower | - | - |
| Sowthistle | - | rosette | - | bolting | - | - |
| Sunflower | - | 1-3" | 3-6" | 6-24" | - | - |
| Thistle, Russian | - | - | - | rosette | - | - |
| Velvetleaf | - | <6" | 6-20" | >20" | - | - |
| ¹ For use in non-food/feed crop only. Adding crop oil concentrate has shown to improve performance on actively growing annual sedge. | | | | | | |

Table 2. Application Rate and Timing – Biennial and Perennial Weeds

(Applicators must follow maximum application rates in the Use Specific Information sections of the label.)

(The addition of liquid fertilizer [28-0-0,32-0-0] at ½ the GPA of the spray solution has proven to give increased suppression or control on certain species of weeds.)

| Weeds Controlled | Rate Per Acre (according to weed growth stage) | | | | | |
|---------------------------------|--|----------|------------------|-------------|----------------|------------------|
| | 1/3 pint | 2/3 pint | 1 pint | 1-1/8 pints | 1-2/3 pints | 2 to 3-1/4 pints |
| Bindweed, Field | - | - | - | - | - | actively growing |
| Bittercress | - | 2-3" | - | - | - | - |
| Buckeye species ¹ | - | - | - | - | full leaf | - |
| Bullnettle ² | - | - | - | flower | - | - |
| Chicory | - | - | - | - | early bolting | - |
| Clove, Bur | - | - | pre-flower | - | - | - |
| Dandelion, Common | - | rosette | - | bolting | - | - |
| Dewberry, Southern ¹ | - | - | - | - | - | spring or fall |
| Dock, Curly | - | - | prior to bolting | - | after bolting | - |
| Elderberry ² | - | - | - | - | - | actively growing |
| Goldenrod, Missouri | - | - | - | 3-15" | flower | - |
| Groundsel, Texas | - | rosette | post-bolting | - | - | - |
| Honeysuckle, Hairy | - | - | - | - | spring or fall | - |

| | | | | | | |
|-------------------------------------|---|------|---------|---------------------|--------------|------------------|
| Horsenettle, Carolina ¹ | - | - | - | - | - | flower or berry |
| Ivy, Poison | - | - | - | after bloom | - | - |
| Knapweed, Black ² | - | - | - | - | - | actively growing |
| Knapweed, Russian ² | - | - | - | - | - | actively growing |
| Knapweed, Spotted | - | - | - | - | - | actively growing |
| Marshelder | - | - | - | <12" | 12"/prebloom | |
| Mesquite ³ | - | - | - | - | - | 45-90 days |
| | | | | | | after budbreak |
| Milkweed, Antelopehorn ² | - | - | - | pre-flower | - | flower |
| Nightshade, Silverleaf ¹ | - | - | - | full flower | - | - |
| Nightshade, Black ¹ | - | - | - | full flower | - | actively growing |
| Persimmon, Eastern ³ | - | - | - | - | - | actively growing |
| Prickly, Lettuce | - | - | - | rosette | - | actively growing |
| Rabbitbrush ² | - | - | - | - | - | - |
| Ragwort, Tansy | - | - | - | rosette | - | actively growing |
| Redvine ² | - | - | - | - | - | actively growing |
| Sagebrush, Fringed ² | - | - | - | - | - | actively growing |
| Smartweed | - | - | - | - | - | - |
| Sorrel, Red | - | - | rosette | bolting | flower | actively growing |
| Sowthistle ² | - | - | - | - | - | actively growing |
| Spurge, Leafy ² | - | - | - | - | - | full leaf |
| Tallow Tree, Chinese ⁴ | - | - | - | - | - | - |
| Thistle, Bull | - | - | rosette | bolting | - | actively growing |
| Thistle, Canada ² | - | - | - | - | - | - |
| Thistle, Musk | - | - | - | rosette/ bolting | - | - |
| Thistle, Plumeless | - | - | rosette | bolting | - | - |
| Vetch, Hairy | - | 1-4" | 4-8" | 8" full flower | - | - |
| Yankee weed | - | - | - | 10-18" | - | rosette |
| Yellow Starthistle ¹ | - | - | - | - | - | - |

¹ May require repeat applications

² Specified rate will provide top growth suppression only.

³ For improved root kill or woody species such as mesquite and eastern persimmon spray 2 pints of per acre LATIGO® each year for 3 consecutive years.

⁴ Under dense populations, a second application may be needed the following growing season. For increased control of weeds such as blackberry and dewberry, LATIGO® may be tank mixed with Ally® herbicide, if labeled for the use site.

Ground Application (Banding)

When applying LATIGO® herbicide by banding, determine the amount of herbicide and water volume needed using the following formula:

$$\begin{array}{l} \text{Bandwidth in inches} \\ \text{Row width in inches} \end{array} \times \begin{array}{l} \text{Broadcast rate} \\ \text{per acre} \end{array} = \begin{array}{l} \text{Banding herbicide} \\ \text{rate per acre} \end{array}$$

$$\begin{array}{l} \text{Bandwidth in inches} \\ \text{Row width in inches} \end{array} \times \begin{array}{l} \text{Broadcast rate} \\ \text{volume per acre} \end{array} = \begin{array}{l} \text{Banding water} \\ \text{volume per acre} \end{array}$$

Ground Application (Broadcast)

Water volume: Use 10-25 gallons of spray solution per broadcast acre for optimal performance. Use the higher spray volume when treating dense or tall vegetation.

Application Equipment: Select nozzle design to produce minimal amounts of fine spray particles. Spray nozzles as close to the weeds as is practical for good weed coverage.

Spot or Small Area Application

LATIGO® may be applied to individual clumps or small areas of undesirable vegetation using handgun or similar types of application equipment. Apply diluted sprays to allow complete wetting (up to runoff) of foliage and stems. For knapsack or other small capacity sprayers, prepare a solution of LATIGO® in water according to Table 3 (assuming that the spot treatment rate equates to 40 gallons per acre on the broadcast basis.) Adding a surfactant (0.5% by volume) can help improve control.

Do not make spot treatments in addition to broadcast or band treatments.

Application equipment: Select nozzles designed to produce minimal amounts of fine spray particles. Spray with nozzles as close to the weeds as is practical for good weed coverage.

Table 3. – Knapsack Sprayer Dilution Instructions

| Sprayer Capacity (gallons of water) | Amount of LATIGO® to add to the spray tank |
|--|---|
| 1 gallon | 2/3 fluid ounce* |
| 3 gallons | 2 fluid ounces |
| 5 gallons | 3 fluid ounces |

*1 fluid ounce = 2 tablespoons

III. ADDITIVES

To improve burndown of emerged weeds, surfactants and/or low use rates of liquid fertilizers (28-0-0; 32-0-0), or crop oil concentrate may be used with LATIGO® herbicide or LATIGO® tank mixes applied after the weeds have emerged. Crop oil concentrate is for non-food/feed crop uses only. Do not apply tank mixes that include Ammonium Sulfate or Crop Oil Concentrate to any food/feed crop use listed on this label. For food/feed crop use, do not use liquid fertilizers that contain Ammonium Sulfate (AMS) as a source of nitrogen as tolerances in commodities derived from the crop may contain residues that exceed established tolerances.

Oil Concentrate

A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all of the following criteria:

- be non-phytotoxic
- contain only EPA-exempt ingredients
- provide good mixing quality in the jar test, and
- be successful in local experience

The exact composition of suitable products will vary; however, vegetable oil and petroleum oil concentrates should contain emulsifiers to provide good mixing quality. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. For additional information, see “Compatibility Test for Mix Components”.

Adjuvants containing crop oil concentrates may be used for preplant, pre-emergence and between cropping applications. Do not use crop oil concentrate for postemergence applications in food/feed crops (i.e., grass (hay or silage), grass grown for seed, pastures, rangeland, and wheat).

Nitrogen Source

Sprayable liquid fertilizers: Use ½ GPA of sprayable liquid fertilizers (28-0-0; 32-0-0) per acre. Do not use brass or aluminum nozzles when spraying fertilizers.

Non-ionic Surfactant

The standard label recommendation is 2-4 pints of an appropriate 80% active non-ionic spray surfactant per 100 gallons of water. (Rate will vary with the size and condition of weeds to be controlled. Use lowest rate per 100 gallons when weeds are small and actively growing. As weeds increase in size and/or become hardened off, the rate of non-ionic surfactant will have to be increased to give optimum coverage and control.)

Table 4. Additive Rate Per Acre

| Additive ¹ | Rate Additive Per Acre |
|---|--|
| Non-ionic Surfactant | 2-4 pints per 100 gallons ² |
| Sprayable Liquid Fertilizers (28-0-0; 32-0-0) | ½ GPA of spray solution |
| Crop Oil Concentrate | 1 quart |

¹ See manufacturer's label for specific rate specifications.

² Use lowest rate per 100 gallons when weeds are small and actively growing. As weeds increase in size and/or become hardened off, the rate of non-ionic surfactant will have to be increased to give optimum coverage and control.

IV. TANK MIXING INFORMATION

Tank Mix Partners/Components

The following products may be tank mixed with LATIGO® according to the specific tank mixing instructions in this label and respective product labels.

| | |
|--|---|
| Aim™ EC (carfentrazone-ethyl) | Harmony® Extra SG Herbicide |
| Ally® XP Herbicide (metsulfuron-methyl) | (thifensulfuron-methyl + tribenuron-methyl) |
| Amber® (triasulfuron) | Karmex® DF (diuron) |
| Asulox® Herbicide (asulam) | Kerb™ 50 WP (pronamide) |
| Atrazine | MCPA |
| Basagran® (bentazon) | Paramount® Herbicide (quinclorac) |
| Cyclone® Star (paraquat dichloride) | Peak® (prosulfuron) |
| Dakota® (fenoxaprop-p-ethyl + MCPA) | Permit® (halosulfuron-methyl) |
| Evik® DF (ametryn) | Roundup® Ultra (glyphosate) |
| Express® Herbicide (tribenuron-methyl) | Sencor® 75% Turf Herbicide (metribuzin) |
| Finesse® Cereal and Fallow Herbicide (chlorsulfuron + metsulfuron-methyl) | Sinbar® WDG (terbacil) |
| Glean® XP Herbicide (chlorsulfuron) | Stinger™ (clopyralid) |
| Gly Star® 5 Extra (glyphosate) | Tordon™ K (picloram) |
| Gramoxone® Inteon (paraquat dichloride) | Touchdown® (glyphosate) |
| | 2,4-D ¹ |

¹ LATIGO® contains 0.30 pound a.e. of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed the combined total of pounds of a.e. per acre per crop cycle of 2,4-D for the use site being applied to.

See "VI. Food/Feed Crop Specific Information" section for more information for more details. Read and follow the applicable Restrictions and Limitations and Directions for Use on all products involved in tank mixing. The most restrictive labeling applies to tank mixes. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture. Physical incompatibility, reduced weed control, or crop injury may result from mixing LATIGO® with other pesticides (fungicides, herbicides, insecticides, or miticides), additives, or fertilizers.

Compatibility Test for Mix Components

Before mixing components, always perform a compatibility jar test.

For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water. For other spray volumes adjust accordingly. Only use water from the intended source at the source temperature.

Add components in the sequence indicated in the Mixing Order using 2 teaspoons for each pound or 1 teaspoon for each pint of specified label rate per acre.

Always cap the jar and invert 10 cycles between component additions.

When the components have all been added to the jar, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent. If the solution is still incompatible, do not mix the ingredients in the same tank.

Mixing Order

If an inductor is used, rinse it thoroughly after each component has been added. Maintain constant agitation during application.

1. Water. Begin by agitating a thoroughly clean sprayer tank half full of clean water.
2. Agitation. Maintain constant agitation throughout mixing and application.
3. Products in PVA bags. Place any product contained in water-soluble bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
4. Water-dispersible products (such as dry flowables, wettable powders, suspension concentrates, and suspo-emulsions)
5. Water-soluble products (such as LATIGO®).
6. Emulsifiable concentrates (such as oil concentrate, when applicable).
7. Water-soluble additives (such as liquid fertilizers [28-0-0; 32-0-0], when applicable).*
8. Remaining quantity of water.

*If sprayable fluid fertilizer is used as the carrier.

Always perform the Compatibility Test before mixing into the spray tank. Also, when using a sprayable fluid fertilizer as the carrier, any product contained in PVA bags must first be completely dissolved in water before the contents can be added to the fertilizer mix.

V. RESTRICTIONS AND LIMITATIONS

- Maximum seasonal use rate: Refer to Table 5.
- Preharvest Interval (PHI): Refer to “Food/Feed Crop Specific Information”
- Restricted-Entry Interval (REI): 48 Hours
- Crop Rotational Restrictions:

The interval between application and planting rotational crop is given below. Always exclude counting days when the ground is frozen. Planting at intervals less than specified below may result in crop injury. Moisture is essential for the degradation of this herbicide in soil.

| Crop | MINIMUM DAYS PLANTBACK INTERVAL (Areas >½" rainfall or irrigation after application)* | | | MINIMUM DAYS PLANTBACK INTERVAL (Areas <½" rainfall or irrigation after application) | | |
|--|--|-------------------|-----------------|---|-------------------|-------------------|
| | 2/3-1 pint/A | >1-3 ½ pints/A | >3-½ pints/A | 2/3-1 pints/A | >1-3 ½ pints/A | >3-1/2 pints/A |
| Corn (field and pop)** | 7 | 21 | 120 | 30 | 60 | 120 |
| Cotton | 21 | 45 | 120 | 30 | 90 | 120 |
| Barley, Oats, Wheat and other small grains | 14 | 21 | 120 | 21 | 60 | 120 |
| Sorghum | 14 | 21 | 120 | 30 | 60 | 120 |
| Soybean*** | 15 (one application) 30 (two applications) | 45 | 120 | 45 | 90 | 120 |
| All other crops | 120 | 120 | DO NOT ROTATE | 120 | 120 | DO NOT ROTATE |

*NOTE: A cumulative 1/2 inch of rainfall or irrigation must occur in 2 or less rainfalls and/or irrigations before calculating plantback interval.

Application Precautions:

- Arid (dry) conditions: it is extremely important that the addition of a suitable Nonionic Surfactant, Oil, or sprayable fertilizer be used when applying LATIGO®. Higher rates of LATIGO® may be needed to control susceptible weeds in this environment.
- Rainfast Period: Rainfall or irrigation occurring within 4 hours after postemergence applications may reduce effectiveness of LATIGO®.
- Stress: Do not apply to crops under stress such as stress due to lack of moisture, hail damage, flooding, herbicide injury, mechanical injury, or widely fluctuating temperatures, as unsatisfactory control may result.

Application Restrictions:

- **Make only one Corn preplant application per crop cycle.
- ***Make only one Soybean preplant application for the 15-day plantback interval.
- Do not apply to crops that show injury (leaf phytotoxicity or plant stunting) produced by any other prior herbicide applications, because this injury may be enhanced or prolonged.
- Do not apply this product through any type of irrigation equipment. Do not contaminate irrigation ditches or water used for domestic purposes.
- This product cannot be used to formulate or reformulate another pesticide product.

Table 5. Crop Specific Restrictions and Limitations

| Crop | Maximum Rate Per Acre Per Application | Maximum Rate Per Acre Per Season | Livestock Grazing or Feeding ¹ | Aircraft Application |
|---------------------------|---------------------------------------|----------------------------------|---|----------------------|
| Between Crop Applications | 3-2/3 pints | 7-1/3 pints | Yes | Yes |
| Pasture, Hay, Silage | 2-1/2 pints | 5 pints | Yes | Yes |
| Grass Grown for Seed | 2 pints | 4 pints | Yes | Yes |
| Wheat | 1-1/4 pints | 2 pints | Yes | Yes |
| Sugarcane | 4 pints | 8 pints | Yes | Yes |

¹ Refer to "Food/Feed Crop Specific Information" for grazing and feeding restrictions.

**VI. FOOD/FEED CROP SPECIFIC INFORMATION
PASTURES, RANGELAND AND GRASS (Hay, Silage)**

LATIGO® may be used for pasture (including pasture grown for hay), rangeland, grass grown for hay or silage, fallow systems, Conservation Reserve Programs, and general farmstead (non-cropland only).

Refer to Tables 1 and 2 for rate selection based on targeted weed or brush species. Some weed species will require tank mixes for adequate control.

Uses described in this section also pertain to small grains (including barley, corn, oats, rye, sudangrass, or wheat) grown for pasture, hay, and silage only. Newly seeded areas including small grains grown for pasture or hay may be injured if rates of LATIGO® greater than 1-1/4 pints per acre are applied.

In newly established hybrid Bermudagrass, Pangolagrass, and stargrasses (*Cynodon* spp.) use 1 to 2 pints of LATIGO® per acre to control or suppress weeds after planting vegetative propagules (stolons) of hybrid bermudagrasses. In addition to the weeds listed in Tables 1 and 2, this rate of LATIGO® will control or suppress annual sedges, broadleaf signalgrass, crabgrass, and goosegrass. Best results will be obtained if LATIGO® is applied at the germinating stage of weeds. Under favorable conditions, this is usually 7-10 days after planting these grasses. Reduced control can be expected if weeds are allowed to reach 1" in height before application or if germination of weeds occurs 10 days after application.

Do not use on bentgrass, susceptible grass pastures (such as carpetgrass, buffalograss, or St. Augustinegrass), lespedeza, wild winter peas, vetch, clover, and alfalfa pastures as injury will occur.

When perennial weeds are reaching maturity, mowing and allowing some regrowth will enhance control. Difficult-to-control weeds may require a repeat application.

For pasture renovations, wait 3 weeks per 1-1/4 pints of LATIGO® used per acre before interseeding or injury may occur.

Grazing and Feeding Non-Lactating Animals: There is no waiting period between treatment and grazing for non-lactating animals. Do not permit meat animals being finished for slaughter to graze treated fields within 30 days of slaughter.

Grazing and Feeding Lactating Animals: Do not graze lactating dairy animals within 7 days of treatment.

Pre-harvest Interval (PHI): Dry hay and Silage: Treated grasses may be harvested for dry hay or silage but do not harvest within 7 days of treatment.

Pasture and Rangeland Tank Mixes – LATIGO® may be applied in tank mixes with one or more of the following herbicides:
Ally®
Amber®

PASTURE & RANGELAND RESTRICTIONS:

- Do not apply more than 2-1/2 pints per acre per application.
- Rates above 2-1/2 pints of LATIGO® per acre are for spot treatments only.
- Do not make more than 2 applications per year.
- Minimum spray interval between applications is 30 days.
- If grass is to be cut for hay, Agricultural Use Requirements for the Worker Protection Standard are applicable.
- For spot treatment, do not exceed 4.4 pints per acre per application, or 8.8 pints per acre per year.

LATIGO® contains 0.225 pound a.e. of dicamba per pint. If applied with other products containing dicamba, either as a tank mix or separately during the same growing season do not exceed 2.0 lbs. of dicamba a.e. per crop cycle.

LATIGO® contains 0.30 pound a.e. of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 4.0 pounds of a.e. per acre per year.

GRASS GROWN FOR SEED

To control many species of emerged broadleaf weeds, apply 8 to 32 oz per acre. Use on newly established or established stands of cool season grasses grown for seed including bentgrasses, bluegrasses, fine fescues, tall fescue, orchardgrass, annual ryegrass and perennial ryegrass.

Apply LATIGO® in the spring or fall when broadleaf weeds are in the 2-4 leaf stage and rosettes are less than 2" across. Larger or more difficult to control broadleaf weeds may require high rates of LATIGO® and a tank mix partner for optimum control.

On newly established stands, the grass must have at least reached the 1 tiller stage prior to treatment.

On established stands, where at least one seed crop has been harvested, two applications separated by 30 days may be made with rates up to 32 oz per acre.

For ground applications apply in 10-20 gallons of spray solution per acre. For aerial applications apply in 5-10 gallons of spray solution per acre.

GRASS GROWN FOR SEED RESTRICTIONS:

- Do not make more than 2 applications per year.
- Do not apply to grasses in the boot stage of development.
- The maximum individual application rate is 32 oz per acre per application.
- The retreatment interval is 30 days.
- Do not use more than 64 oz of LATIGO® per acre per year.

SUGARCANE

Applications of LATIGO® can be made any time after the weeds have emerged and are actively growing but prior to the close-in stage of sugarcane. When possible, direct the spray beneath the sugarcane canopy in order to minimize the likelihood of crop injury. The use of directed sprays will also aid in maximizing spray coverage of weed foliage. Application rates and timing are given below. Use the higher level of listed rate ranges when treating dense vegetative growth.

- For control of listed ANNUAL broadleaf weeds, apply 1 quart of LATIGO® per treated acre.
- For suppression of listed PERENNIALS, apply 1-2 quarts of LATIGO® per treated acre.

SUGARCANE Tank Mixes: LATIGO® may be tank mixed with one or more of the following herbicides: Asulox, Atrazine, Evik, Sencor, Sinbar

SUGARCANE RESTRICTIONS:

- Pre-emergent Application: Do not make more than one pre-emergence application per crop cycle.
- Pre-emergent Application: Do not apply more than 4 pints per application.
- Post-emergent Application: Do not make more than one post-emergence application per crop cycle.
- Post-emergent Application: Do not apply more than 4 pints per application.
- Do not harvest cane prior to crop maturity.
- Preharvest interval (PHI) is 87 days.

LATIGO® contains 0.225 pound a.e. of dicamba per pint. If applied with other products containing dicamba, either as a tank mix or separately during the same growing season do not exceed 2.0 lbs. of dicamba a.e. per crop cycle.

LATIGO® contains 0.30 pound a.e. of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 4.0 pounds of a.e. per acre per year.

TEFF

LATIGO® may be applied by ground or air.

For ground applications, apply in 10 to 20 gallons of total broadcast spray mix per acre. For aerial application, apply in 3 to 10 gallons of water per acre.

GRASS SEED CROPS

LATIGO® application rates and spray volumes will vary with the growth stage and population of broadleaf weeds to be controlled. In general the smaller the weed, the lower use of the specified rate range will provide satisfactory control.

The larger the weed, the population and environmental conditions will require the higher end of the rate range to achieve the satisfactory control especially for many of the perennial weeds.

To control emerged broadleaf weeds, apply 0.75-1.25 pints of LATIGO® per acre. Apply after the two-leaf stage and prior to the boot stage. Refer to "Plant Response" and "Livestock Feeding Restrictions" under the GRASS PASTURES section of the label.

PASTURES, RANGELAND AND GRASS (Hay, Silage)

In newly established Teff, use 0.75 to 1.25 pints of LATIGO® per acre to control or suppress weeds after planting. Newly seeded areas including small grains grown for pasture or hay, may be injured if rates of LATIGO® are greater than 1.25 pints per acre are applied. Apply after the two-leaf stage and prior to the boot stage.

SMALL GRAIN CROPS (Grains used for Gluten-Free Flour)

Apply up to 1.25 pints of LATIGO® per acre to Teff. Apply after the two-leaf stage and prior to the boot stage. If small grains are grown for pasture or hay only, refer to Pastures, Rangeland and Grass (Hay, Silage).

TEFF RESTRICTIONS:

Grazing and Feeding Non-Lactating Animals: There is no waiting period between treatments and grazing for non-lactating animals. Do not permit meat animals being finished for slaughter to graze treated fields within 30 days of slaughter.

Grazing and Feeding Lactating Animals: Do not graze lactating dairy animals within 7 days of treatment.

Pre-harvest Interval (PHI): Dry Hay and Silage: Treated grasses may be harvested for dry hay or silage but do not harvest within 37 days of treatment.

The Restricted Entry Interval (REI) is 48 hours.

The Pre-harvest Interval (PHI) (Not for Dry Hay and Silage) is 14 days.

Limited to one postemergence application per crop cycle.

Limited to one preharvest application per crop cycle.

WHEAT
(Fall- and Spring-seeded)

If small grains are grown for pasture or hay only, refer to Pastures, Rangeland and Grass (Hay, Silage).

Do not graze or harvest for livestock feed prior to crop maturity.

Do not use LATIGO® in wheat underseeded with legumes.

EARLY-SEASON POST-EMERGENCE APPLICATION (Fall- and Spring-Seeded):

Apply 1-1/4 pints of LATIGO® per acre to wheat unless using one of the wheat specific programs below.

Early-season applications to spring-seeded wheat must be made after tillering and before wheat reaches the 6-leaf stage.

Early-season applications to fall-seeded wheat must be made after tillering and prior to the jointing stage. Care should be taken in staging early developing wheat varieties such as TAM 107, Madison, or Wakefield to be certain that the application occurs prior to the jointing stage.

SPECIFIC USE PROGRAMS FOR FALL-SEEDED WHEAT ONLY:

Up to 3/4 pint of LATIGO® per acre may be applied post-emergent on fall-seeded wheat after the wheat begins to tiller for suppression of perennial weeds, such as field bindweed. Applications may be made in the fall following a frost but before a killing freeze. Periods of extended stress such as cold and wet weather may enhance the possibility of crop injury. For fall applications only, do not use if the potential for crop injury is not acceptable.

PREHARVEST APPLICATIONS:

LATIGO® can be used to control weeds that may interfere with harvest of wheat. Apply up to 1-1/4 pints of LATIGO® per acre as a broadcast or spot treatment to annual broadleaf weeds when wheat is in the hard dough stage and the green color is gone from the nodes (joints) of the stem. Best results will be obtained if application can be made when weeds are actively growing but before weeds canopy.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture.

Do not use preharvest-treated wheat for seed unless a germination test is performed on the seed with an acceptable result of 95% germination or better. For control of additional broadleaf weeds or grasses, LATIGO® may be tank mixed with other herbicides such as Ally® or Gly Star® Plus that are registered for preharvest use in wheat.

Preharvest use of LATIGO® is not registered for use in California.

Table 6 – Wheat Tank Mixes

| LATIGO® can be tank mixed with one or more of the following herbicides | | | |
|--|-------------------|-----------------------|-------------------|
| Bromoxynil | Diuron | Metsulfuron-methyl | Tribenuron-methyl |
| Carfentrazone-ethyl | Fenoxypop-p-ethyl | Prosulfuron | |
| Chlorsulfuron | MCPA | Triasulfuron | |
| Clopyralid | Metribuzin | Thifensulfuron-methyl | |

¹ Do not use low rates of sulfonyleurea herbicide on more mature weeds or on dense vegetative growth.

² Do not use as a tank mix treatment with Fenoxypop-p-ethyl or MCPA on Durum wheat.

³ Tank mixes with Diuron and Metribuzin are for use in fall-seeded wheat only.

Fallow Systems, Conservation Reserve Programs, and General Farmstead

These uses are considered Food/Feed Crops when harvested, grazed or foraged. Consult section on "General Tank Mixing Information" for adjuvant restrictions and section on "Additives" for specific use directions.

WHEAT RESTRICTIONS:

- Postemergence:
 - Make no more than one application per crop cycle.
 - Do not apply more than 1-1/4 pints per acre per application.
- Preharvest:

- Make no more than one application per crop cycle.
- Do not apply more than 1-1/4 pints per acre per application.
- Pre-Harvest interval is 14 days.

LATIGO® contains 0.225 pound a.e. of dicamba per pint. If applied with other products containing dicamba, either as a tank mix or separately during the same growing season do not exceed 2.0 lbs. of dicamba a.e. per crop cycle.

LATIGO® contains 0.30 pound a.e. of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not:

- Exceed a combined total of 1.25 pounds of a.e. per acre per crop cycle of 2,4-D for post emergent use.
- Exceed 0.5 pound of a.e. per acre per crop cycle of 2,4-D for pre-harvest application.
- Exceed a total of 1.75 pounds of a.e. per acre per crop cycle for all uses.

**VII. NON-FOOD/FEED USE (LAND NOT HARVESTED, GRAZED OR FORAGED) –
SPECIFIC INFORMATION BETWEEN CROP APPLICATIONS**

PREPLANT DIRECTIONS (POSTHARVEST, FALLOW, CROP STUBBLE, SET-ASIDE) FOR BROADLEAF WEED CONTROL

LATIGO® can be applied postharvest in the fall, spring, or summer during the fallow period or to crop stubble/set-aside acres. Apply to weeds after crop harvest (postharvest) and before a killing frost or in the fallow cropland or crop stubble the following spring or summer. To aid in suppressing certain perennial or biennial broadleaf weeds (including cotton regrowth), this product may be applied either alone or in combination with other registered herbicides. For cotton regrowth, a minimum rate of 1-1/5 pints/acre is recommended.

See “V. Restrictions and Limitations” for the recommended interval between application and planting to prevent crop injury.

Rates and Timings:

Apply 1 to 3-2/3 pints of LATIGO® per acre. Refer to Table 1 to determine use rates for specific targeted weed species. For best performance, apply LATIGO® when annual weeds are less than 6" tall, when biennial weeds are in the rosette stage and to perennial weed regrowth in late summer or fall following a mowing or tillage treatment. The most effective control of upright perennial broadleaf weeds such as Canada thistle and Jerusalem artichoke occurs if LATIGO® is applied when the majority of weeds have at least 4-6" of regrowth or for weeds such as field bindweed and hedge bindweed that are in or beyond the full bloom stage. The addition of liquid fertilizers (28-0-0, 32-0-0) at ½ GPA has shown to increase efficacy.

Avoid disturbing treated areas following application. Treatments may not kill weeds that develop from seed or underground plant parts such as rhizomes or bulblets, after the effective period for LATIGO®. For seedling control, a follow-up program or other cultural practices could be instituted.

Between Crop Tank Mixes:

It is the pesticide user’s responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture.

In tank mixes with one or more of the following herbicides, apply 1.0-1.25 pints of LATIGO® per acre for control of annual weeds, or 1.25-4.25 pints of LATIGO® per acre for control of biennial and perennial weeds:

| | | | |
|---------------------|--------------------|---------------------|--------------|
| Atrazine | Glyphosate | Paraquat dichloride | Quinclorac |
| Carfentrazone-ethyl | Metribuzin | Picloram | Triasulfuron |
| Chlorsulfuron | Metsulfuron-methyl | Pronamide | |

APPLICATIONS TO FALLOW GROUND PRIOR TO PLANTING COTTON

Rates and Timings

Apply LATIGO® as a broadcast or spot treatment to emerged and actively growing weeds at the rate of 1 to 3-2/3 pints per acre. The most effective control of weeds occurs if application is made when weeds are in the 2- to 4-leaf stage and rosettes are less than 2" across.

Cropping Restrictions

Refer to the Crop Rotational Restrictions Table in Section V. RESTRICTIONS AND LIMITATIONS for appropriate pre-plant application intervals for cotton.

Tank Mix Treatments

For control of grasses or additional broadleaf weeds, LATIGO® may be tank mixed with CAPROL® and GRAMAXONE® Extra, and glyphosate herbicides.

BETWEEN CROP APPLICATION RESTRICTIONS:

- Do not apply more than 3-2/3 pints per acre per application.
- Do not make more than 2 applications per year.
- Minimum spray interval between applications is 30 days.
- Only labeled crops can be planted within 30 days of application.

LATIGO® contains 0.225 pound a.e. of dicamba per pint. If applied with other products containing dicamba, either as a tank mix or separately during the same growing season, do not exceed 2.0 lbs of dicamba a.e. per crop cycle.

LATIGO® contains 0.30 pound a.e. of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 4.0 pounds of a.e. per acre per year.

CONSERVATION RESERVE PROGRAMS

LATIGO® may be used for Conservation Reserve Programs, general farmstead (non-cropland only), weed and brush control, or use in State Recognized Noxious Weed areas (non-cropland areas).

Refer to Tables 1 and 2 for rate selection based on targeted weed or brush species. Some weed species will require tank mixes for adequate control.

CONSERVATION RESERVE PROGRAMS RESTRICTIONS:

- Do not apply more than 4 pints per acre per application.
- Do not make more than 2 applications per year.
- Minimum spray interval between applications is 30 days.
- If grass is to be cut for hay, Agricultural Use requirements for the Worker Protection Standard are applicable.

LATIGO® contains 0.225 pound a.e. of dicamba per pint. If applied with other products containing dicamba, either as a tank mix or separately during the same growing season do not exceed 2.0 lbs. of dicamba a.e. per crop cycle.

For program lands, including Conservation Reserve Program, consult program rules to determine whether grass or hay may be used. The more restrictive requirements of the program rules or this label must be followed.

LATIGO® contains 0.30 pound a.e. of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 4.0 pounds of a.e. per acre per year.

GENERAL FARMSTEAD

Farmstead and Fence-row Treatment Application Instructions

LATIGO® may be applied using water or oil and water emulsions in spot application to control undesirable vegetation using handgun or similar types of application equipment. In addition to weed species listed in Tables 1 and 2, these treatments may be used to control or suppress woody plant species listed in Table 7.

To prepare soil and water emulsions, mix in the order and proportions indicated below.

The solution should remain milky colored without an oily layer on top when under agitation. If an oily layer forms, increase the amount of emulsifier or change to a more effective emulsifier.

Do not exceed 40 gallons of spray solution per treated acre per application. 4 pints of LATIGO® in forty gallons of spray solution contains 0.9 pound acid equivalent of dicamba and 1.2 pounds acid equivalent of 2,4-D. Spray plants to wet. Do not allow this spray mix to contact desirable vegetation.

To control brush, briars, and weeds along fence-rows surrounding pasture and ranch lands, and fallow fields, use a tank mix of 1.5% LATIGO®, 88.5% water, 10% diesel oil, and sufficient emulsifier (to mix the diesel and emulsifier). The diesel oil in this

tank mix will damage or kill desirable grasses and should not be used in pastures or where damage to desirable species cannot be tolerated.

1. Water: Begin by agitating a thoroughly clean sprayer tank with the desired quantity of clean water. Maintain constant agitation during complete mixing procedure.
 2. Emulsifier: Add 0.5% volume to volume of water.
 3. LATIGO®: Add 1.5 gallons per 100 gallons of total intended solution.
 4. Diesel Oil: Add 10 gallons per 100 gallons of total intended solution.
- Maintain constant agitation during application. Under good agitation, the spray solution should be milky white with no oil layer on top. If oil layer forms, increase the amount of emulsifier or change to a more effective emulsifier.

FOR SPRAYING FOLIAR APPLICATIONS:

1. Spray when leaves have reached full size but have not hardened due to drought or maturity.
2. Spray individual plants to wet with handgun.
3. For larger stems (up to 3" in diameter) and hard to control species, direct spray stream to base of stems to wet the stem at soil surface in addition to wetting the foliage.
4. Do not apply under drip line of desirable trees or adjacent to desirable vegetation.

FOR DORMANT BASAL APPLICATIONS:

1. Increase diesel oil content to 15% or 15 gallons of diesel oil per 100 gallons of total solution.
2. Spray in late winter and early spring before plants break dormancy.
3. Spray the bottom 24" of the target stem to wet on all sides.
4. For larger stems (up to 3" in diameter) and hard-to-kill species direct the spray solution to the base of target stems to wet the soil at the stem/soil junction in addition to wetting the stem.
5. Do not apply under drip line of desirable trees or adjacent to desirable vegetation.

FARMSTEAD AND FENCEROW RESTRICTIONS:

- Postemergence (annual & perennial weeds):
 - Do not make more than 2 applications per year.
 - Do not apply more than 4 pints per acre per application.
 - Minimum spray interval between applications is 30 days.
- Postemergence (woody plants):
 - Do not make more than 1 application per year.
 - Do not apply more than 4 pints per acre per application.

Applications to non-cropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

LATIGO® contains 0.225 pound a.e. of dicamba per pint. If applied with other products containing dicamba, either as a tank mix or separately during the same growing season do not exceed 2.0 lbs. of dicamba a.e. per crop cycle.

LATIGO® contains 0.30 pound a.e. of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 4.0 pounds of a.e. per acre per year.

FOR CUT SURFACE TREATMENTS:

Apply LATIGO® in an undiluted state as a cut surface treatment to control unwanted trees and prevent sprouts of cut trees.

- Frill or Girdle Treatments: Make a continuous cut or a series of overlapping cuts using an axe to girdle tree trunk. Spray or paint the cut surface with LATIGO®.
- Stump Treatments: Spray or paint freshly cut surface with LATIGO®. The cambium layer (the area adjacent to the bark) should be thoroughly wet. Treat stumps within 6 hours after cutting.

Table 7. The following list of trees and vines can be controlled on farmsteads and fencerows as foliar, basal, or cut surface treatments:

| | | |
|-----------------------|------------------------|-------------------|
| Alder | Hemlock | Poplar |
| Ash | Hickory | Rabbitbrush |
| Aspen | Honeylocust | Redcedar, Eastern |
| Basswood | Honeysuckle | Rose, McCartney |
| Beech | Hornbeam | Rose, Multiflora |
| Blackberry | Huckleberry | Sagebrush, Fringe |
| Blackgum | Huisache | Sassafras |
| Cedar | Ivy, Poison | Spruce |
| Cherry | Kudzu | Sumac |
| Chinquapin | Locust, Black | Sweetgum |
| Cottonwood | Maple | Sycamore |
| Creosotebush | Mesquite | Tarbrush |
| Dewberry | Oak | Willow |
| Dogwood | Oak, Poison | Witchhazel |
| Elm | Olive, Russian | Yaupon |
| Grape | Persimmon, Eastern | Yucca |
| Greenbriar | Pine | |
| Hawthorn (Thornapple) | Plum, Sand (Wild Plum) | |

CUT SURFACE RESTRICTIONS:

- Do not make more than one cut surface application per year.
- Do not use more than 8.8 pints per 100 gallons of spray solution.

Weeds listed in this label:

| Common Name | Scientific Name |
|--------------------------|-----------------------------------|
| ANNUALS | |
| Beebalm, Spotted | <i>Monarda punctata</i> |
| Broomweed, Common | <i>Gutierrezia dracunculoides</i> |
| Buckwheat, Wild | <i>Polygonum convulvulus</i> |
| Buffalobur | <i>Solanum rostratum</i> |
| Burdock | <i>Arctium spp.</i> |
| Buttercup, Corn | <i>Ranunculus arvensis</i> |
| Chickweed, Common | <i>Stellaria media</i> |
| Cockle, Corn | <i>Agrostemma githago</i> |
| Cocklebur, Common | <i>Xanthium strumarium</i> |
| Coreopsis, Plains | <i>Coreopsis tinctoria</i> |
| Croton, Woolly | <i>Croton capitatus</i> |
| Devilsclaw | <i>Proboscidea luisianica</i> |
| Dogfennel (Cypressweed) | <i>Eupatorium capillifolium</i> |
| Eveningprimrose, Cutleaf | <i>Oenothera lacinata</i> |
| Flax | <i>Linum catharticum</i> |
| Fleabane, Annual | <i>Erigeron annuus</i> |
| Flixweed | <i>Descurainia sophia</i> |
| Henbit | <i>Lamium amplexicaule</i> |
| Horseweed/Marestail | <i>Conyza canadensis</i> |
| Knotweed, Prostrate | <i>Polygonum aviculare</i> |
| Kochia | <i>Kochia scoparia</i> |
| Lambsquarters, Common | <i>Chenopodium album</i> |
| Lettuce, Prickly | <i>Lactuca serriola</i> |
| Mallow, Common | <i>Malva neglecta</i> |
| Morningglory, Ivyleaf | <i>Ipomea hederacea</i> |
| Morningglory, Tall | <i>Ipomea purpurea</i> |
| Mustard, Annual | <i>Brassica spp.</i> |

| | |
|--------------------------|---------------------------------|
| Mustard, Tansy | <i>Descurainia pinnata</i> |
| Pennycress, Field | <i>Thlaspi arvense</i> |
| Pepperweed, Virginia | <i>Lepidium virginicum</i> |
| Pigweed, Prostrate | <i>Amaranthus blitoides</i> |
| Pigweed, Redroot | <i>Amaranthus retroflexus</i> |
| Pigweed, Smooth | <i>Amaranthus hybridus</i> |
| Pigweed, Tumble | <i>Amaranthus albus</i> |
| Poorjoe | <i>Diodia teres</i> |
| Purslane, Common | <i>Portulaca oleracea</i> |
| Ragweed, Common | <i>Ambrosia artemisiifolia</i> |
| Ragweed, Lance-leaf | <i>Ambrosia bidentata</i> |
| Ragweed, Western | <i>Ambrosia psilostachya</i> |
| Sedge | <i>Cyperus compressus</i> |
| Shepherdspurse | <i>Capsella bursa-pastoris</i> |
| Smartweed, Pennsylvania | <i>Polygonum pennsylvanicum</i> |
| Sneezeweed, Bitter | <i>Helenium amurum</i> |
| Sunflower, Common (wild) | <i>Helianthus annuus</i> |
| Thistle, Russian | <i>Salsola iberica</i> |

| Common Name | Scientific Name |
|---------------------------------|---------------------------------|
| BIENNIALS AND PERENNIALS | |
| Bindweed, field | <i>Convolvulus arvensis</i> |
| Bittercress | <i>Cardamine spp.</i> |
| Buckeye | <i>Aesculus spp.</i> |
| Bullnettle | <i>Cnidoscopus stimulosus</i> |
| Chicory | <i>Cichorium intybus</i> |
| Clover, Hop | <i>Trifolium aureum</i> |
| Dandelion | <i>Taraxacum officinale</i> |
| Dock, Curly | <i>Rumex crispus</i> |
| Elderberry | <i>Sambucus canadensis</i> |
| Goldenrod, Missouri | <i>Solidago missouriensis</i> |
| Goldenweed, Common | <i>Isocoma coronopifolia</i> |
| Groundsel | <i>Senecio vulgaris</i> |
| Honeysuckle, Hairy | <i>Lonicera</i> |
| Horsenettle | <i>Solanum carolinense</i> |
| Ivy, Poison | <i>Rhus radicans</i> |
| Knapweed, Black | <i>Centaurea nigra</i> |
| Knapweed, Russian | <i>Centaurea repens</i> |
| Knapweed, Spotted | <i>Centaurea maculosus</i> |
| Marshelder | <i>Ina annua</i> |
| Mesquite | <i>Prosopis juliflora</i> |
| Milkweed, Antelopehorn | <i>Asclepius</i> |
| Nightshade, Silverleaf | <i>Solanum elaeagnifolium</i> |
| Nightshade, Black | <i>Solanum nigrum</i> |
| Persimmon, Eastern | <i>Diospyros virginiana</i> |
| Rabbitbrush | <i>Chrysanthemum pulchellus</i> |
| Ragwort, Tansy | <i>Senecio jacobia</i> |
| Redvine | <i>Brunnichia ovata</i> |
| Sagebrush, Fringed | <i>Artemisia frigida</i> |
| Smartweed, Swamp | <i>Polygonum coccineum</i> |
| Sorrel, Red (Sheep Sorrel) | <i>Rumex acetosella</i> |
| Sowthistle, Perennial | <i>Sonchus arvensis</i> |
| Spurge, Leafy | <i>Euphorbia esula</i> |
| Starthistle, Yellow | <i>Centaurea solstitialis</i> |

| | |
|----------------------|-----------------------------------|
| Tallow Tree, Chinese | <i>Sapium sebiferum</i> |
| Thistle, Bull | <i>Cirsium vulgare</i> |
| Thistle, Canada | <i>Cirsium arvense</i> |
| Thistle, Musk | <i>Carduus nutans</i> |
| Thistle, Plumeless | <i>Carduus acanthoides</i> |
| Vetch | <i>Vicia spp.</i> |
| Yankeeeweed | <i>Eupatorium compositifolium</i> |

Food/Feed Crop Uses

This product can be used on the following:

- Conservation Reserve Program Land
- Fallow Systems (Between Crop Application)
- General Farmstead
- Grass (Hay or Silage)
- Pastures
- Grass Grown for Seed
- Rangeland
- Sugarcane
- Teff
- Wheat

Look inside for complete Restrictions and Limitations and Application Instructions.

These crops are considered Food/Feed crops only when harvested, grazed, or foraged. Otherwise, they are considered non-Food/Feed uses.

CONDITIONS OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions or presence of other materials. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

HELENA CHEMICAL COMPANY warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, HELENA CHEMICAL COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS FOR ANY PARTICULAR PURPOSE OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. When Buyer suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), Buyer must promptly notify Seller in writing of any claims to be eligible to receive either remedy stated above. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL HELENA CHEMICAL COMPANY OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. HELENA CHEMICAL COMPANY and the Seller offer this product, and the Buyer accepts it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of HELENA CHEMICAL COMPANY. No employee or agent of HELENA CHEMICAL COMPANY or the Seller is authorized to vary or exceed the terms of this Warranty in any other manner.

LATIGO® is a registered trademark of Helena Holding Company.

Ally®, Canvas®, Express®, Finesse®, Glean®, Harmony®, Karmex® and Sinbar® are registered trademarks of E.I. du Pont de Nemours & Company.

Banvel®, Clarity®, Distinct®, Fallowmaster® and Paramount® are registered trademarks of BASF Corporation.

Basagran® and Laddok® S-12 are registered trademarks of BASF AG.

Asulox®, Bronate®, Butril®, Dakota® and Tiller® are registered trademarks of Bayer CropScience.

Amber®, Evik®, Caparol®, Cyclone®, Sencor®, Gramoxone®, Peak®, Rave™ and Touchdown® are trademarks and/or registered trademarks of Syngenta Crop Protection.

Landmaster®, Permit® and Roundup® Ultra are registered trademarks of Monsanto Company.

Curtail™, Stinger™, Grazon™, Kerb™ and Tordon™ are trademarks of Dow AgroSciences LLC.
Aim™ is a trademark of FMC Corporation.
Fallow Star® and Gly Star® Plus are registered trademarks of Albaugh, LLC.