

GROUP 4A INSECTICIDE

GROUP 3 4 7 FUNGICIDES

Do not use, sell or distribute this product within or into California.

WARDEN® CEREALS 360

A seed treatment product for protection against damage from certain insects and diseases of small grain cereals (barley, oat, rye, triticale, wheat) and enhanced germination, root and top growth

Active Ingredient:

Thiamethoxam	5.79%
Difenoconazole	3.47%
Mefenoxam	0.87%
Sedaxane	0.87%
Cytokinin, as Kinetin.....	0.054%
Gibberellic Acid	0.018%
Indole Butyric Acid.....	0.027%
Other Ingredients:	88.901%
Total:	100.000%

WARDEN® CEREALS 360 is a flowable concentrate containing 0.51 lb thiamethoxam; 0.31 lb difenoconazole; 0.08 lb mefenoxam; 0.08 lb sedaxane; 0.0048 lb cytokinin; 0.0016 lb gibberellic acid and 0.0024 lb indole butyric acid per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use in booklet [on label].

EPA Reg. No. 100-1564-1381

EPA Est. No. 100-NE-001
Net Contents

Distributed by:
Winfield Solutions, LLC
P.O. Box 64589
St. Paul, MN 55164-0589

1/0212/6

WINFIELD™

FIRST AID

Have the product container with you when calling a poison control center or doctor, or going for treatment.

HOT LINE NUMBER

For 24 Hour Medical Emergency Assistance (Human or Animal) Call **1-877-424-7452** or
Chemical Emergency Assistance (Spill, Leak, Fire, or Accident),
Call
1-800-424-9300

PRECAUTIONARY STATEMENTS**Hazards to Humans and Domestic Animals****CAUTION**

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)**Mixers, loaders, applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves such as polyvinyl chloride [PVC] ≥14 mils, nitrile rubber ≥14 mils, butyl rubber ≥14 mils, barrier laminate, neoprene rubber ≥14 mils, or Viton® ≥14 mils

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticides get inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product is toxic to wildlife, freshwater and estuarine/marine fish and highly toxic to aquatic invertebrates. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate. Exposed treated seed may be hazardous to wildlife.

Pollinator Precautions

Thiamethoxam is highly toxic to bees, and effects are possible as a result of exposure to translocated residues in blooming crops.

If treated seed is spilled outdoors or in areas accessible to birds, promptly clean up or bury to prevent ingestion.

Groundwater Advisory

Mefenoxam is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may lead to groundwater contamination. Thiamethoxam has properties and characteristics associated with chemicals detected in ground water. Mefenoxam and Thiamethoxam may leach into the ground water if used in areas where soils are permeable, particularly where the water table is shallow.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of WINFIELD SOLUTIONS, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold WINFIELD SOLUTIONS, LLC and Seller harmless for any claims relating to such factors.

WINFIELD SOLUTIONS, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or WINFIELD SOLUTIONS, LLC, and (2) Buyer and User assume the risk of any such use. To the extent permitted by applicable law, WINFIELD SOLUTIONS, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall WINFIELD SOLUTIONS, LLC be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF WINFIELD SOLUTIONS, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF WINFIELD SOLUTIONS, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

WINFIELD SOLUTIONS, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of WINFIELD SOLUTIONS, LLC.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use in commercial seed treatment facilities. Use is also permitted as an end-use seed treatment on agricultural establishments at planting, or immediately before planting, as specified in the specific Crop Use Directions. This product is to be used in liquid or slurry treaters only.

Do not make any soil or foliar application of products containing a neonicotinoid to crops grown from seed treated with WARDEN® CEREALS 360.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigor and poor quality, except for the purpose of curative control of existing disease pests, may result in reduced germination and/or reduction of seed and seedling vigor. Conduct germination tests on a small portion of seed before committing the total seed lot to a selected seed treatment. Treat using equipment similar to that planned for treating the total seed lot. Due to seed quality and seed storage conditions beyond the control of WINFIELD SOLUTIONS, LLC, no claims are made to guarantee the germination of carry-over treated seed.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notifications to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride [PVC] ≥14 mils, nitrile rubber ≥14 mils, butyl rubber ≥14 mils, barrier laminate, neoprene rubber ≥14 mils, or Viton® ≥14 mils
- Shoes plus socks

PRODUCT INFORMATION

WARDEN® CEREALS 360 is a seed treatment product containing one insecticide active ingredient (thiamethoxam), three fungicide active ingredients (difenoconazole, mefenoxam, and sedaxane) and three plant-growth regulators (cytokinin, indole butyric acid and gibberellic acid). WARDEN® CEREALS 360 seed treatment protects against damage from certain early-season insects and certain seed- and soil-borne diseases of cereal crops.

Difenoconazole provides protection against several seed, seedling and certain foliar disease including bunts and smuts. Mefenoxam protects against damping-off caused by *Pythium* spp. Sedaxane protects against *Rhizoctonia* spp. and loose smuts (*Ustilago* spp.).

Thiamethoxam is a systemic seed treatment insecticide belonging to the neoticotinoid class of chemistry. Thiamethoxam in WARDEN® CEREALS 360 protects against aphids, wireworm and European chafer.

WARDEN® CEREALS 360 contains the plant-growth regulators cytokinin, gibberellic acid and indolebutyric acid to aid in enhanced germination and early season root and top growth.

GROUP 4A INSECTICIDE

Resistance Management

WARDEN® CEREALS 360 contains thiamethoxam, a Group 4A insecticide. Some insect pests are known to develop resistance to products after repeated use. Because resistance development cannot be predicted, the use of this product should conform to sound resistance management strategies established for the crop and use area. WINFIELD SOLUTIONS, LLC encourages responsible product stewardship to ensure effective long-term control of the insects on this label.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot

be attributed to improper application or weather conditions, a resistant strain of insect or pathogen may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

Insect biotypes with acquired or inherent resistance to Group 4A insecticides may eventually dominate the insect population if Group 4A insecticides are used repeatedly as the predominant method of control for targeted species. This may result in partial or total loss of control of those species by WARDEN® CEREALS 360 or other Group 4A insecticides.

In order to maintain susceptibility to this class of chemistry:

- Avoid using Group 4A insecticides exclusively for season long control of insect species with more than one generation per crop season.
- For insect species with successive or overlapping generations, apply WARDEN® CEREALS 360 and other Group 4A insecticides using a “treatment window” approach. A treatment window is a period of time as defined by the stage of crop development and/or the biology of the pests of concern. Within the treatment window, depending on the length of residual activity, there may either be single or consecutive applications (seed treatment, soil, foliar, unless otherwise stated) of the Group 4A insecticides. Do not exceed the maximum WARDEN® CEREALS 360 allowed per calendar year.
- Following a treatment window of Group 4A insecticides, rotate to a treatment window of effective products with a different mode of action before making additional applications of Group 4A insecticides.
- A treatment window rotation, along with other IPM practices for the crop and use area, is considered an effective strategy for preventing or delaying a pest’s ability to develop resistance to this class of chemistry.
- If resistance is suspected, do not reapply WARDEN® CEREALS 360 or any other Group 4A insecticides.

Other Insect Resistance Management (IRM) practices include:

- Incorporating IPM techniques into your insect control program.
- Monitoring treated insect populations for loss of field efficacy.
- Using tank-mixtures or premixes with insecticides from a different target site of action group as long as the involved products are all registered for the same crop and effective rates are applied.

For additional information on Insect Resistance Management:

- Contact your local extension specialist, certified crop advisor and/or product manufacturer for additional insect resistance management recommendations.
- Visit the Insecticide Resistance Action Committee (IRAC) on the web at:
<http://www.irc-online.org>

GROUP	3	4	7	FUNGICIDES
--------------	----------	----------	----------	-------------------

WARDEN® CEREALS 360 contains difenoconazole, a Group 3 fungicide; mefenoxam, a Group 4 fungicide; and sedaxane, a Group 7 fungicide. Difenoconazole belongs to the triazole class of chemistry and is a demethylation inhibitor of sterol biosynthesis (DMI). DMI's disrupt membrane synthesis by blocking demethylation. Mefenoxam is a systemic fungicide belonging to the phenylamide class of chemistry which adversely affects fungal RNA synthesis. Sedaxane belongs to the carboxamide class of chemistry and is a succinate dehydrogenase inhibitor (SDHI). SDHI's inhibit fungal metabolism by binding to the succinate dehydrogenase enzyme thereby disrupting cellular respiration and energy generation.

Fungal populations may contain individuals naturally resistant to Group 3, 4, or 7 fungicides. If Group 3, 4, or 7 fungicides are used repeatedly in the same fields, then resistant members may eventually dominate the population. Because resistance development cannot be predicted, the use of this product should conform to sound resistance management strategies established for the crop and use area.

Seed treatment use should be based on an IPM program that includes field sanitation, scouting, historical information related to pesticide use, and crop rotation. The IPM program should also consider cultural, biological, and other chemical control practices.

Where possible, rotate or mix the use of Group 3, 4, or 7 fungicides with different fungicide groups to which resistance has not developed.

WINFIELD SOLUTIONS, LLC encourages responsible product stewardship to ensure effective long term control of the fungal diseases on this label.

For additional information on fungicide resistance management:

- Contact WINFIELD SOLUTIONS, LLC representatives at 1-800-334-9481
- Contact your local extension specialist or certified crop advisor
- Visit the Fungicide Resistance Action Committee (FRAC) on the web at: <http://www.frac.info>

ROTATIONAL CROP RESTRICTIONS

In the event of crop failure or harvest of a crop grown from seed treated with WARDEN® CEREALS 360, crops may be replanted according to the following schedule:

Plantback Interval Table

Immediate Plantback	Minimum 30-Day Plantback Interval
Cereal Grains: Barley, Oats, Rye, Triticale, and Wheat	Alfalfa
	Cereal Grains Crop Group 15
	Cotton
	Cucurbit Vegetables Crop Group 9
	Fruiting Vegetables Crop Group 8
	Head and Stem Brassica Crop Subgroup 5A
	Leafy Brassica Greens Crop Subgroup 5B
	Leafy Vegetable (Except <i>Brassica</i> Vegetables) Crop Group 4
	Legume Vegetables (Succulent or Dried) Crop Group 6
	Mint: Peppermint and Spearmint
	Oilseeds: Borage, Canola (Rapeseed), Crambe, Flax Seed, Mustard Seed, and Safflower
	Onion, Bulb
	Peanut
	Root Vegetables (including sugarbeet) Crop Subgroup 1A
	Tuberous and Corm Vegetables (including potato) Crop Subgroup 1C
Strawberry	
Sunflower	
Tobacco	

For any other crops the minimum plantback interval is 120 days from the date the seeds treated with WARDEN® CEREALS 360 were planted. A cover crop other than the crops listed above that is planted for erosion control or soil improvement may be planted sooner than the 120 day interval; however, the crop may not be grazed or harvested for food or feed.

MIXING INSTRUCTIONS

Important: Always re-circulate WARDEN® CEREALS 360 thoroughly before using.

Apply WARDEN® CEREALS 360 as a water-based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of insect and disease control.

WARDEN® CEREALS 360 may be applied in tank mixes or sequentially with other EPA-registered seed treatment pesticide products. If using WARDEN® CEREALS 360 in a tank mixture with other seed treatment products, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank-mix partner label(s).

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products of the mixture (for example, first aid from one product, spray drift management from another).

Consult the manufacturer of the application equipment you plan to employ for suitability for this application and for instructions on operation and calibration of the equipment. Follow the manufacturer application instructions for the seed treatment equipment being used.

Continuous agitation or mixing of the slurry mixture is necessary to prevent settling out of the solution.

Allow seed to dry before bagging.

Follow planter manufacturer specifications for use of talc or other hopper box additives at planting. Seed must be completely dry before adding to planter.

WARDEN® CEREALS 360 contains an EPA-approved dye/colorant that imparts an unnatural color to the seed as required by the Federal Seed Act.

SEED BAG LABEL REQUIREMENTS

The Federal Seed Act requires that bags containing treated seeds shall be labeled with the following statements:

- This seed has been treated with difenoconazole, mefenoxam and sedaxane fungicides, thiamethoxam insecticide, and cytokinin, indole butyric acid and gibberellic acid plant growth regulators.
- Do not use for feed, food, or oil purposes.

In addition, the U.S. Environmental Protection Agency requires the following statements on bags containing seeds treated with WARDEN® CEREALS 360:

- Pollinator Precaution: Thiamethoxam is highly toxic to bees, and effects are possible as a result of exposure to translocated residues in blooming crops.
- Groundwater Advisory:
Mefenoxam is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may lead to groundwater contamination. Thiamethoxam has properties and characteristics associated with chemicals detected in ground water. These chemicals may leach into the ground water if used in areas where soils are permeable, particularly where the water table is shallow.
- Store away from feeds and foodstuffs.
- Wear long-sleeved shirt, shoes, long pants and chemical resistant gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- Do not allow children, pets, or livestock to have access to treated seed.
- Dispose of all excess treated seed. Leftover treated seed may be doublesown around the headland or buried away from water sources in accordance with local requirements.
- Do not contaminate water bodies when disposing of planting equipment wash waters.
- Dispose of seed packaging in accordance with local requirements.
- Treated seed must be planted into the soil at a depth greater than 0.5 inch.
- In the event of crop failure or harvest of a crop grown from seed treated with WARDEN® CEREALS 360, crops may be replanted according to the following schedule:

Plantback Interval Table

Immediate Plantback	Minimum 30-Day Plantback Interval
Cereal Grains: Barley, Oats, Rye, Triticale, and Wheat	Alfalfa
	Cereal Grains Crop Group 15
	Cotton
	Cucurbit Vegetables Crop Group 9
	Fruiting Vegetables Crop Group 8
	Head and Stem Brassica Crop Subgroup 5A
	Leafy Brassica Greens Crop Subgroup 5B
	Leafy Vegetable (Except Brassica Vegetables) Crop Group 4
	Legume Vegetables (Succulent or Dried) Crop Group 6
	Mint: Peppermint and Spearmint
	Oilseeds: Borage, Canola (Rapeseed), Crambe, Flax Seed, Mustard Seed, and Safflower
	Onion, Bulb
	Peanut
	Root Vegetables (including sugarbeet) Crop Subgroup 1A
	Tuberous and Corm Vegetables (including potato) Crop Subgroup 1C
Strawberry	
Sunflower	
Tobacco	

- For any other crop, the minimum plant back interval is 120 days from the date the WARDEN® CEREALS 360 treated seed was planted. A cover crop other than the crops listed above that is planted for erosion control or soil improvement may be planted sooner than the 120-day interval; however, the crop may not be grazed or harvested for food or feed.

- Do not make any soil or foliar application of products containing a neonicotinoid to crops grown from seed treated with WARDEN® CEREALS 360.
- Do not apply any product containing cytokinin, gibberellic acid, or indole butyric acid as an in-furrow, band, side dress or mark out application.
- Do not use at a rate that will result in more than 0.05 lb thiamethoxam per acre (22.7 grams ai/A) per calendar year as a seed treatment application.
- This seed has been treated with 0.008 mg ai thiamethoxam per seed assuming an average seed size of 12,000 seeds per pound.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed, and (2) no measurable residues of pesticide remain in the ethanol byproducts that are used in agronomic practice.

CROP USE DIRECTIONS

Crop	Effect	Use rate for all crops (5 fl oz per 100 lb seed)
Barley Oats Rye Triticale Wheat	Growth Regulation: Enhanced germination and early-season root and top growth	
Barley Oats Rye Spring Wheat Winter Wheat	Insect Protection against: Aphids ¹ European Chafer Wireworms ¹	
Barley	Disease Protection against: General Seed Rots ² Seedling Blight, Root Rot, and Damping-Off caused by seed- and soil-borne <i>Fusarium</i> or <i>Rhizoctonia</i> spp. Seedling Blight, Root Rot, and Damping-Off caused by soil-borne <i>Pythium</i> spp. Seed-borne Septoria Covered Smut False Loose Smut True Loose Smut Diseases Suppressed³: Common Root Rot (<i>Cochliobolus</i> spp.) Fusarium Crown and Foot Rot Take-All	
Oats	Disease Protection against: General Seed Rots ² Seedling Blight, Root Rot, and Damping-Off caused by seed- and soil-borne <i>Fusarium</i> or <i>Rhizoctonia</i> spp. Seedling Blight, Root Rot, and Damping-Off caused by soil-borne <i>Pythium</i> spp. Seed-borne Septoria Covered Smut Loose Smut	

	<p>Diseases Suppressed³: Common Root Rot (<i>Cochliobolus</i> spp.)</p>	
Rye	<p>Disease Protection against: General Seed Rots² Seedling Blight, Root Rot, and Damping-Off caused by seed- and soil-borne <i>Fusarium</i> or <i>Rhizoctonia</i> spp. Seedling Blight, Root Rot, and Damping-Off caused by soil-borne <i>Pythium</i> spp. Seed-borne Septoria Common Bunt⁴ Dwarf Bunt⁴</p> <p>Diseases Suppressed³: Common Root Rot (<i>Cochliobolus</i> spp.) Fusarium Crown and Foot Rot Take-All</p>	
Spring Wheat	<p>Disease Protection against: General Seed Rots² Seedling Blight, Root Rot, and Damping-Off caused by seed- and soil-borne <i>Fusarium</i> or <i>Rhizoctonia</i> spp. Seedling Blight, Root Rot, and Damping-Off caused by soil-borne <i>Pythium</i> spp. Seed-borne Septoria Fusarium Seed Scab Common Bunt⁴ Karnal Bunt Loose Smut Pythium Damping-Off</p> <p>Diseases Suppressed³: Common Root Rot (<i>Cochliobolus</i> spp.) Fusarium Crown and Foot and Root Rot Take-All</p>	<p>Use rate for all crops (5 fl oz per 100 lb seed)</p>
Triticale Winter Wheat	<p>Disease Protection against: General Seed Rots² Seedling Blight, Root Rot, and Damping-Off caused by seed- and soil-borne <i>Fusarium</i> or <i>Rhizoctonia</i> spp. Seedling Blight, Root Rot, and Damping-Off caused by soil-borne <i>Pythium</i> spp. Seed-borne Septoria Septoria Leaf Blotch⁵ Common Bunt⁴ Flag Smut Fusarium Seed Scab Dwarf Bunt⁴ Karnal Bunt Loose Smut Pythium Damping-Off</p> <p>Diseases Suppressed³: Common Root Rot (<i>Cochliobolus</i> spp.) Fusarium Crown and Foot and Root Rot Take-All</p>	

¹The 5 fluid oz/100 lb rate of WARDEN® CEREALS 360 provides suppression of wireworm activity, aphids and aphid vectored barley yellow dwarf virus. If pressure is moderate or high or control is required, apply up to 0.8 fl oz additional Cruiser 5FS.

²This includes rots caused by *Fusarium*, *Pythium*, *Rhizoctonia*, *Penicillium* and *Aspergillus* spp.

³Suppression means consistent protection at a level which is not optimal but is still of commercial benefit.

⁴Protects against both seed- and soil-borne bunts.

⁵Provides early-season foliar disease protection for first 4 weeks after planting. For full-season protection, apply a foliar fungicide according to label directions.

Restrictions: If seed has been treated with Warden Cereals 360, do not apply any product containing cytokinin, gibberellic acid, or indole butyric acid as an in-furrow, band, side dress or mark out application. Do not apply more than 0.05 lbs of thiamethoxam per acre per calendar year.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling [less than or equal to 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Container Handling [greater than 5 gallons – mini-bulk]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Container Handling [greater than 5 gallons – bulk]

Refillable container. Refill this container with pesticide only. Do not use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

Warden® is a Trademark of Winfield Solutions, LLC

Viton® trademark of E. I. du Pont de Nemours and Company