



Bayer CropScience LP  
 P.O. Box 12014  
 2 T.W. Alexander Drive  
 Research Triangle Park, North Carolina 27709  
 1-866-99BAYER (1-866-992-2937)

## Luna Sensation

For control of Anthracnose in Leafy  
 Vegetables (4A and 4B)  
 EPA Reg. No. 264-1090

### FIFRA 2(ee) Recommendation

#### FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF CALIFORNIA

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

#### Leafy greens subgroup 4A (except Watercress)

Amaranth; Arugula; Chervil; Chrysanthemum, Edible-Leaved; Chrysanthemum, Garland; Corn Salad; Cress, Garden; Cress, Upland; Dandelion; Dock; Endive; Lettuce (head and leaf); Orach; Parsley; Purslane, Garden; Purslane, Winter; Radicchio (Red Chicory); Spinach; Spinach, New Zealand; Spinach, Vine.

Disease Control	Application Rate	Product Instructions
Anthracnose ( <i>Microdochium panattonianum</i> <i>Colletotrichum spp.</i> )	7.6 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval. When disease pressure is severe, use the higher rates.  Can be applied in a band, lettuce (head and leaf) only.

#### Restrictions:

- Do not apply more than 15.3 fl oz of LUNA SENSATION per acre per year.
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, do not apply more than 0.446 lbs fluopyram or 0.375 lbs trifloxystrobin per acre per year, including soil and foliar uses.
- Can be applied the day of harvest for broadcast foliar uses, 20 days for banded applications for lettuce (head and leaf) only.
- To limit the potential for development of disease resistance to these fungicide classes, do not make more than 2 sequential applications of LUNA SENSATION or any Group 7 or Group 11 containing fungicide before rotating with a fungicide from a different Group.

**Leaf petioles subgroup 4B**

Cardoon; Celery; Celery, Chinese; Celtuce; Fennel, Florence; Rhubarb; Swiss Chard.

<b>Disease Control</b>	<b>Application Rate</b>	<b>Product Instructions</b>
Anthracnose ( <i>Colletotrichum spp.</i> )	5.8 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval. When disease pressure is severe, use the higher rates.

**Restrictions:**

- Do not apply more than 15.3 fl oz of LUNA SENSATION per acre per year.
- Can be applied using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, do not apply more than 0.446 lbs fluopyram or 0.375 lbs trifloxystrobin per acre per year, including soil and foliar uses.
- Do not apply LUNA SENSATION within 7 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, do not make more than 2 sequential applications of LUNA SENSATION or any Group 7 or Group 11 containing fungicide before rotating with a fungicide from a different Group.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.  
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any crop-protection product, always read and follow label directions.  
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*