



FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MINNESOTA

EVEREST® 2.0 HERBICIDE

EPA Reg. No. 66330-391

EPA SLN No. MN-110002

**For Application on Kentucky Bluegrass Grown For Seed (Only During the Year of Establishment) in the State of Minnesota
FIFRA §24(c) Special Local Need Registration**

| Active Ingredient | By weight |
|---|------------------|
| Flucarbazono-sodium, 4,5-Dihydro-3-methoxy-4-methyl-5-oxo-N- [[2-(trifluoromethoxy)phenyl]sulfonyl]-1H- 1,2,4-triazole-1-carboxamide, sodium salt..... | 35.0% |
| Other Ingredients | 65.0% |
| Total..... | 100.0% |

This formulation contains 3.5 lbs of Flucarbazono-Sodium active ingredient per gallon (419 g ai/l)

KEEP OUT OF REACH OF CHILDREN

CAUTION

**In Case of Medical Emergency Involving This Product,
Call Day or Night: 866-303-6952**

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This labeling and the main label must be in the possession of the user at the time of application.

Read the entire label for EVEREST 2.0 HERBICIDE before proceeding with the use directions contained in this supplemental labeling. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label and this supplemental label.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor

drift from areas treated. Do not contaminate water when disposing of equipment washwaters or rinsate.

Do not allow sprays to drift onto adjacent desirable plants.

USE RATES

Apply EVEREST 2.0 Herbicide to Kentucky bluegrass grown for seed (only during the year of establishment) for control of wild oats, windgrass, and other grass and broadleaf weeds at a rate of 1.0 fl oz/A, with a non-ionic surfactant at a rate of 0.25% v/v.

| Recommended Rates for Kentucky Bluegrass Grown For Seed | | |
|---|----------------|--|
| Grass Weeds | Rate (fl oz/A) | Notes |
| Wild Oats (<i>Avena fatua</i>) | 1.0 oz/A | Control |
| Windgrass (<i>Apera spica-venti</i>) | 1.0 oz/A | Control |
| Cheat (<i>Bromus secalinus</i>) | 1.0 oz/A | Control with fall applications. Suppression with spring applications. |
| Ventenata (<i>Ventenata dubia</i>) | 1.0 oz/A | Suppression |
| Broadleaf Weeds | Rate | Notes |
| Wild Mustard | 1.0 | Control |
| Black Mustard | 1.0 | Control |
| Blue Mustard | 1.0 | Control |
| Tansy Mustard | 1.0 | Control |
| Tumble Mustard | 1.0 | Control |

GROUND APPLICATION

Apply in water using a spray volume of 5 to 10 gallons/acre (or 50 to 100 liters/hectare) at 30 to 50 psi to ensure proper weed coverage. Flat fan nozzles of 80 or 110 degrees are recommended for optimum coverage. Do not use floodjet or controlled droplet application (CDA) equipment. Nozzles may be oriented 45 degrees forward to enhance crop penetration and to give better weed coverage.

AERIAL APPLICATION

Apply in water using a minimum spray volume of 3 gal/A (or 30 L/ha). For best results, particularly in dry conditions or where heavy weed infestations exist, use a minimum of 5 gal/A (or 50 L/ha). Use nozzles that provide 200 to 350 micron size droplets for best results and to insure uniform spray coverage. Aerial applications with EVEREST 2.0 Herbicide should be made with low drift nozzles at a maximum height of 10 feet above the crop and at a maximum pressure of 40 psi. Do not apply aurally when wind speed is greater than 10 mph. Do not allow spray to drift onto adjacent crops, as injury or loss may occur.

Refer to the **SPRAY DRIFT MANAGEMENT** section of the federal label for additional information on how to reduce drift during aerial application.

ENDANGERED SPECIES PROTECTION

To avoid adverse effects on endangered dicot plant species, the following measures will be required where endangered plant species occur in the counties listed in the table below:

| State | County |
|-----------|---|
| Minnesota | Clay Kittson Norman Polk Red Lake |

For ground applications, the applicator must:

- Apply when there is sustained wind away from native plant communities, OR
- Use low-pressure nozzles according to manufacturer's specifications that produce only coarse or very coarse droplets, OR
- Leave a 50 foot untreated buffer between the treatment and native plant communities.

For aerial applications, the applicator must:

- Apply only when there is sustained wind away from native plant communities, OR
- Leave a 350 foot untreated buffer between the treatment and native plant communities.

RESTRICTIONS/PRECAUTIONS

Do not apply more than 1.0 fl oz/A of EVEREST 2.0 Herbicide per year.

Do not make more than one postemergence application of EVEREST 2.0 Herbicide per year.

Do not use EVEREST 2.0 Herbicide in annual production systems of Kentucky bluegrass grown for seed.

For use under this SLN label do not apply EVEREST 2.0 Herbicide through any type of irrigation system.

Under this SLN label, use of EVEREST is deemed a non-food use by the Minnesota Department of Agriculture and may be applied to Kentucky bluegrass seed fields only during the year of establishment.

Do not apply EVEREST 2.0 Herbicide within 1 year (365 days) of harvesting Kentucky bluegrass seed.

Do not graze livestock or harvest forage from treated Kentucky bluegrass seed fields within 1 year (365 days) of application.

CONTAINER DISPOSAL GUIDANCE

Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal.

ARYSTA LIFESCIENCE NORTH AMERICA, LLC'S WARRANTY DISCLAIMER FOR USE OF EVEREST 2.0 HERBICIDE ON KENTUCKY BLUEGRASS GROWN FOR SEED

EVEREST 2.0 Herbicide when used on Kentucky bluegrass grown for seed may lead to crop injury, loss, or damage. Arysta LifeScience North America, LLC recommends that the user and/or grower test EVEREST 2.0 Herbicide in order to determine its suitability for such intended use. Arysta LifeScience North America, LLC makes EVEREST 2.0 Herbicide available to the user and/or grower solely to the extent the benefit and utility, in the sole opinion of the user and/or grower, outweigh the extent of potential injury associated with the use of this product. The decision to use or not to use EVEREST 2.0 Herbicide must be made by each individual EVEREST 2.0 Herbicide user and/or grower on the basis of possible crop injury from EVEREST 2.0 Herbicide, the severity of weed infestation, the cost of alternative weed controls, and other factors. Arysta LifeScience North America, LLC intends that because of the risk of crop damage that all such use is at the user's and/or grower's risk. Arysta LifeScience North America, LLC makes no warranty, express or implied, with respect to the suitability of EVEREST 2.0 Herbicide for use on Kentucky bluegrass grown for seed.

**ARYSTA LIFESCIENCE NORTH AMERICA, LLC
15401 Weston Parkway, Suite 150
Cary, North Carolina 27513**

EVEREST is a registered trademark of Arysta LifeScience North America, LLC.
Label Revision Date: April 1, 2011