



Bayer CropScience LP
 P.O. Box 12014
 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina 27709
 1-866-99BAYER (1-866-992-2937)

LUNA[®] SENSATION
For control of Citrus diseases
at reduced application rates
 EPA Reg. No. 264-1090

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF TEXAS

This recommendation expires December 31, 2019

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

Fruit, citrus, group 10-10

Australian Desert Lime; Australian Finger Lime; Australian Round Lime; Brown River Finger Lime; Calamondin; Citron; Citrus Hybrids; Grapefruit; Japanese Summer Grapefruit; Kumquat; Lemon; Lime; Mediterranean Mandarin; Mount White Lime; New Guinea Wild Lime; Orange, Sour; Orange, Sweet; Pummelo; Russell River Lime; Satsuma Mandarin; Sweet Lime; Tachibana Orange; Tahiti Lime; Tangelo; Tangerine (Mandarin); Tangor; Trifoliate Orange; Uniq Fruit; cultivars, varieties and/or hybrids of these.

Disease Control	Application Rate	Product Instructions
Greasy spot (<i>Mycosphaerella citri</i>)	4.0 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 21-day interval. When disease pressure is severe, use the shorter intervals.
Melanose (<i>Diaporthe citri</i>)	5.0 fl oz/acre	

Restrictions:

- Do not apply more than 27.3 fl oz of LUNA SENSATION per acre per year.
- Apply using ground equipment.
- Regardless of formulation or method of application, do not apply more than 0.446 lbs fluopyram or 0.5 lbs trifloxystrobin per acre per year, including soil and foliar uses.
- Do not apply LUNA SENSATION within 7 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, do not make more than 2 sequential applications of LUNA SENSATION or any Group 7 or Group 11 containing fungicide before rotating with a fungicide from a different Group.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.
 For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any crop-protection product, always read and follow label directions.
 For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*