

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.



EPA Reg. No. 279-3588

2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF CT, GA, NJ, MA, MI, MD, NC, OH, PA, SC, VA, WV

THIS FIFRA SECTION 2(EE) EXPIRES JANUARY 30, 2021

To permit the use of Rhyme[®] fungicide in peach as a foliar application for control of rusty spot (*Podosphaera leucotricha*).

Directions For Use

- IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.
- ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.
- This recommendation must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for Rhyme[®] Fungicide before applying.

Crop

- Peach

Rates and Timing

Diseases	Product Rate (fl. oz./A)	Application Directions	Application Interval (Days)
Rusty spot (<i>Podosphaera leucotricha</i>)	7 (0.113 lb ai/A)	Start applications at petal fall and continue through 3rd Cover (3C) or until terminal shoot growth ceases and shoots harden off or conditions are no longer favorable for disease development	7

Restrictions:

- No single application may exceed 7 fl. oz. of product/A.
- Do not apply more than 4 applications per year.
- Do not apply more than 28 fl. oz. of product/A per year.
- Do not apply more than 0.455 lb. ai. of flutriafol/A per year.
- Restricted Entry Interval (REI) is 12 hours.
- Preharvest Interval (PHI): Do not apply within 7 days of harvest.



FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104

FMC and Rhyme are trademarks of FMC Corporation
Copyright © 2018