



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

ADMIRE® PRO Systemic Protectant
EPA Reg. No. 264-827
plus
Maxim® 4FS
EPA Reg. No. 100-758
Tank Mixture for use on Potatoes

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF CONNECTICUT, DELAWARE, NEW JERSEY, MARYLAND, RHODE ISLAND, VERMONT, VIRGINIA and WEST VIRGINIA,

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. It is a violation of federal law to use this product in a manner inconsistent with its label. This product bulletin must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

ADMIRE® PRO + Maxim® 4SC can be applied to potato seed for the control of many sucking and chewing potato insects and for protection against several seed-borne and soil-borne fungi.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling.

Product	Rate Fluid Ounces/100 lbs. of seed	Crop	Pests Controlled
ADMIRE® PRO Systemic Protectant + Maxim® 4FS	0.25 to 0.35 + 0.08 to 0.16	Potato	Aphids (including green peach aphid), Colorado potato beetles, flea beetles, leafhoppers, potato psyllids and wireworms (seed-piece protection). <i>Fusarium</i> dry rot seed decay, seed-borne <i>Rhizoctonia</i> (stem canker and tuber black scurf) and seed-borne <i>Helminthosporium solani</i> (silver scurf disease on potato tubers).

Notes and Restrictions:

Do not use treated seed-pieces for food, feed or fodder. Do not apply any subsequent application of a neonicotinoid insecticide following an ADMIRE PRO seed-piece treatment.

Applications:

To assure maximum efficacy, proper coverage of potato pieces is necessary. Use a minimum of 1 fl oz of total treatment for each 100 pounds of potato seed. When combining ADMIRE PRO and Maxim 4FS, water must be added as the middle component. Add ADMIRE PRO to the mix tank first followed by the calculated amount of water. Stir until ADMIRE PRO is uniformly distributed in solution. Finally, add Maxim 4FS and mix into solution. Maintain adequate agitation to keep materials suspended.

Example: To treat 100 lbs of potato seed pieces

- Add 0.35 fl oz of ADMIRE PRO
- Followed by 0.57 fl oz of water, stir
- Add 0.08 fl oz Maxim 4FS
- Maintain agitation

Fungicidal or inert absorbent dusts may be applied after ADMIRE PRO + Maxim 4FS application.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937) or

Visit our worldwide web site at www.bayercropscienceus.com

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).