

Product Bulletin



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Blackhawk®

EPA Reg. No. 62719-523

Special 2(ee) Recommendation†
(For Distribution and Use in the State of Texas)

For Suppression of Potato Tuberworm in Potato

This label expires and must not be distributed or used in accordance with this product bulletin after December 31, 2020.

Refer to product label for Blackhawk for general use precautions, mixing directions, application directions, and restrictions.

ATTENTION

†This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Blackhawk Naturalyte insecticide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Blackhawk Naturalyte according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Blackhawk Naturalyte.

Directions for Use

Pest	Blackhawk (oz/acre)
Potato tuberworm	3.5 (0.08 lb ai/acre)

Refer to the Blackhawk product label for General Use Precautions, Mixing Directions and Application Directions. Blackhawk should be used as part of an integrated program to manage potato tuberworm which may require multiple applications.

Application Timing and Coverage: Treat in accordance with local economic thresholds. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area. Application coverage is important for potato tuberworm management, regardless of method used. Ensure coverage is optimized.

Blackhawk is a Group 5 insecticide (spinosad and spinetoram active ingredients). Do not make more than four applications of a Group 5 insecticide per crop. It is also recommended in the case of potato tuberworm to not make more than two consecutive applications of a Group 5 insecticide. Follow additional resistance management recommendations on the product label.

Restrictions:

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 7 days apart.
- Do not apply more than a total of 14.4 oz of Blackhawk (0.33 lb ai of spinosad) per acre per year.
- **Maximum Number of Applications:** Do not make more than four applications per acre per year.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow
R373-014

Approved: 09/12/17

Replaces R373-012