

# MycoApply® Endo



MycoApply® Endo is a granular mycorrhizal inoculum that consists of a blend four endomycorrhizal fungi species. Endo contains 60,000 propagules per pound.

## CONTAINS NON-PLANT FOOD INGREDIENTS:

### Soil Amending Guarantee Analysis:

ACTIVE INGREDIENTS: Endomycorrhizae: *Glomus intraradices*,  
*Glomus aggregatum*, *Glomus mosseae*, *Glomus etunicatum*  
(33 propagules/gram each) ..... 75%

Inert Ingredients (Clay) ..... 25%

### INDEX:

- 1.0 First Aid
- 2.0 Recommendations for Use
- 3.0 Storage and Handling
- 4.0 Warranty and Disclaimer

KEEP OUT OF REACH OF CHILDREN

### CAUTION

#### 1.0

First Aid	
If on skin	Wash with soap and water. Exposure to skin may cause mild irritation and dry skin.
If in eyes	Rinse opened eye for several minutes under

	running water. If symptoms persist, consult a doctor.
If swallowed	If used material is ingested, seek medical attention.
If inhaled	Supply fresh air. If irritation or other symptoms occurs, seek medical attention.

## 2.0 RECOMMENDATIONS FOR USE

Endo can be applied to existing plants, seed beds or roots at transplant time. The goal is to make physical contact between the inoculum and the roots/seed.

Endo can be banded under seed, worked into seed beds, placed under cuttings, blended into potting soil, or sprinkled on roots at transplant time. The goal is to create physical contact between the roots and the inoculant.

Agriculture: Endo can be banded in rows or side dressed before or during planting at 2 pounds per acre.

Restoration: Endo can be hydromulched or banded in rows with seed during planting at 20 pounds per acre. Use 40 to 60 pounds per acre broadcast and rototill before planting.

Nurseries: Endo can be mixed into the soil before/during filling cavities, pots and trays. Endo may be mixed mechanically using equipment that allows passage of a #8 screen (2.5 mm) or larger particles. For propagation, use 1.0 to 2.0 lbs. per cubic yard. Use higher rates if filling smaller cavities.

Turf: Apply 0.25 to 0.5 lb. of Endo per 1,000 square feet during installation or aerification. Water thoroughly following application, using overhead irrigation.

For Planting: Apply Endo in-furrow at a rate of 1 teaspoon per row foot. For use on cuttings, apply 0.5 teaspoon under the cutting. For potted plants or transplants, apply 1 to 2 tablespoons of Endo per gallon-size pot. For ball and burlap, apply 0.5 to 1.5 ounce of Endo per inch of stem caliper.

### 3.0 STORAGE AND HANDLING

**Storage:** Store product in a dry area. Avoid prolonged storage above 120 ° Fahrenheit. Keep container tightly sealed.

**Handling:** Avoid breathing dust. Ensure good ventilation/exhaustion at the workplace. For operations where the exposure limit may be exceeded, a NIOSH/MSHA approved high efficiency particulate respirator is recommended. Use of safety glasses or goggles is recommended.

**Spills:** Stop spill at source. Reclaim material if possible for reuse. Containerize contaminated material for disposal. After all visible traces have been removed, flush area thoroughly with water.

**Product & Waste Disposal:** Dispose of all waste in accordance with local, state and federal regulations.

### 4.0 WARRANTY AND DISCLAIMER

To the extent consistent with applicable law, seller makes no warranty, express or implied of merchantability, fitness or otherwise concerning the use of this product other than as indicated on the label. To the extent consistent with applicable law, user assumes all risks of use, storage or handling not in strict accordance with accompanying directions.

“This Product is not a Plant Food” Information regarding the contents and levels of metals in this product is available on the internet at:

<http://www.aapfco.org/metals.htm>

MycoApply® is a registered trademark of Mycorrhizal Applications

Manufactured by:  
Mycorrhizal Applications  
P.O. Box 1029  
Grants Pass, OR 97528

Distributed by:  
Valent U.S.A. Corporation  
P.O. Box 8025  
Walnut Creek, CA 94596-8025