


SMARTLINE US61384098B (170628) LABMC MOVENTO 32 FOZ - CMYK 9/18/17



2.75"

COPY AREA

FIRST AID

If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first five minutes, then continue rinsing.• Call a poison control center or doctor immediately for treatment advice.
Have a product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-334-7577 for emergency medical treatment information.	
Note to Physician: No specific antidote is available. Treat the patient symptomatically.	

1.75"

2.25"

3.25"

2.75"

COPY AREA

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

- Harmful if swallowed or absorbed through skin.
- Causes moderate eye irritation.
- Avoid contact with skin, eyes, or clothing.
- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.
- Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.


Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical resistant gloves made of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, natural rubber \geq 14 mils, polyethylene, polyvinyl chloride \geq 14 mils, or viton \geq 14 mils.
- Shoes plus socks.

1.75"

2.25"

3.25"



2.75"
COPY AREA

ENVIRONMENTAL HAZARDS

For Terrestrial Use: This pesticide is toxic to aquatic invertebrates and oysters. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. This product may contaminate water through drift of spray in wind. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters or rinsate.


This chemical has properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.


This product is potentially toxic to honey bee larvae through residues in pollen and nectar, but not to adult honeybees. Exposure of adult bees to direct treatment or residues on blooming crops can lead to effects on honeybee larvae. See the "Directions for Use" section of this label for specific crop application instructions that minimize risk to honey bee larvae.

1.75"

2.25"

3.25"





2.75"

COPY AREA

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

DIRECTIONS FOR USE


**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Read the entire label before using this product.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

1.75"

2.25"

3.25"



2.75"

COPY AREA


AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It also contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours following application.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water), is:



- Coveralls.
- Chemical resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride ≥ 14 mils, or viton ≥ 14 mils.
- Shoes plus socks.





PRODUCT INFORMATION

MOVENTO®:

- Is a suspension concentrate formulation and is active primarily by ingestion against immature target pest life stages. In addition, fertility of adult female target pests, such as aphids and whiteflies, may be reduced.
- Can be applied by air, ground equipment or through chemigation as a preventative treatment or timed to coincide with an early threshold level in developing insect populations.
- Must be tank-mixed with a spray adjuvant / additive having spreading and penetrating properties to maximize leaf uptake and systemicity of the active ingredient within treated plants; please contact your local Bayer CropScience representative or PCA for specific recommendations by crop.
- It is widely known that tank mixtures and/or sequential treatments of horticultural spray oil with Captan and/or sulfur may cause adverse plant compatibility in tree and vine crops; including MOVENTO in this tankmix and/or sequential treatment scenario is not recommended.
- Following application to plant foliage, MOVENTO is fully systemic, moving through phloem and xylem to new shoot, leaf and root tissues; systemicity and efficacy may be hindered during periods of cold temperatures, under drought conditions, or when plants are not actively growing.







2.75"

COPY AREA

INSECT RESISTANCE MANAGEMENT RECOMMENDATIONS

MOVENTO contains an active ingredient with a mode of action classified as a Group 23 Insecticide, i.e., a lipid biosynthesis inhibitor (LBI). To delay insecticide resistance:


- Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area.
- Bayer CropScience strongly encourages that MOVENTO, applied alone or in tankmix combination with another Group 23 product, be applied in a block rotation or windowed approach with products from other chemical classes having a different mode of action before using additional applications of Group 23 insecticides against the same target pest. Using a block rotation or windowed approach, along with other IPM practices, is considered an effective use strategy for preventing or delaying an insect pest's ability to develop resistance to a given class of chemistry.

Contact your local extension specialist, certified crop advisor, and/or Bayer CropScience representative for additional resistance management or IPM recommendations. Also, for more information on Insect Resistance Management (IRM), visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://irac-online.org>.

1.75"

2.25"

3.25"



2.75"

COPY AREA

Chemigation monitoring

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Required system safety devices

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor/engine stops or in cases where there is no water pump, when water pressure decreases to the point where pesticide distribution is adversely affected. The irrigation line
(continued)

1.75"

2.25"


3.25"




or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Using water from public water systems

Do not connect an irrigation system (including greenhouse system) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet




(continued)




end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional automatic quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must contain a functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Injection for chemigation

Inject the specified dosage of MOVENTO into the irrigation main water stream: (1) through a constant flow, metering device; (2) into the center of the main line flow via a pitot tube or equivalent; (3) at a point ahead of at least one, right-angle turn in the main stream flow such that thorough mixing with the irrigation water is ensured.







Flushing and Cleaning the chemical injection system


At the end of the application period, allow time for all lines to flush the pesticide through all nozzles or emitters before turning off irrigation water. To ensure the lines are flushed and free of pesticides, a dye indicator may be injected into the lines to mark the end of the application period.

In order to apply pesticides accurately, the chemical injection system must be kept clean, free of chemical or fertilizer residues and sediments. Refer to your owner's manual or ask your equipment supplier for the cleaning procedure for your injection system.

SPRAY DRIFT MANAGEMENT

Do not apply when wind speed favors drift beyond the area intended for treatment. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all of these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator.





2.75"

COPY AREA

1.75"

Temperature Inversions

Do not make aerial or ground applications during temperature inversions. Drift potential is high during temperature inversions. Temperature inversions restrict vertical air mixing, which causes small-suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog. However, if fog is not present, the movement of smoke from a ground source can also identify inversions. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Airblast (Air Assist) Applications for Tree Crops and Vineyards


Airblast sprayers carry droplets into the canopy of trees/vines via a radial, or lateral directed air stream. The following drift management practices should be followed:

- Adjust deflectors and aiming devices so that spray is only directed into the canopy;
- Block off upward pointed nozzles when there is no overhanging canopy;
- Use enough air volume to penetrate the canopy and provide good coverage;

(continued)

2.25"

3.25"



2.75"

COPY AREA

1.75"

- For further information, contact your local Bayer representative.
- If your supplier and Bayer representative have no experience with the combination you are considering, you should conduct a test to determine physical compatibility.
- To determine physical compatibility, add the recommended proportions of each chemical with the same proportion of water, as will be present in the chemical supply tank, into a suitable container, mix thoroughly and allow to stand for five minutes. If the combination remains mixed, or can be readily re-mixed, the mixture is considered physically compatible.


Compatibility

MOVENTO is physically and biologically compatible with many registered pesticides and fertilizers or micronutrients. However, it is known that many components, including crop protection products, fertilizers, micronutrients, and spray adjuvants, may be present in a tank mix combination. There is potential for adverse chemical reactions. It is impossible to determine physical, biological, and plant compatibility for all scenarios that may be encountered; therefore, it is recommended that users determine the chemical, physical, biological and plant compatibility of such mixes prior to making applications on a broad commercial scale. Observe the most restrictive of the labeling instructions and precautions of all products used in mixtures.

22

2.25"

3.25"




Order of Mixing

The proper mixing procedure for MOVENTO alone or in tank mix combinations with other pesticides is:

1. Fill the spray tank 1/4 to 1/3 full with clean water;
2. While recirculating and with the agitator running, add any products in Polyvinyl acetate (PVA) bags **(See Note)**. Allow time for thorough mixing;
3. Continue to fill spray tank with water until 1/2 full;
4. Add any other wettable powder (WP) or wettable granules (WG) products;
5. Add the required amount of MOVENTO, and any other "flowable" (FL or SC) type products; add required amount of MOVENTO, and;
6. Allow enough time for thorough mixing of each product added to tank;
7. If applicable, add any remaining tank mix components: emulsifiable concentrates (EC), fertilizers and micronutrients;
8. Fill spray tank to desired level and maintain constant agitation to ensure uniformity of spray mixture.

NOTE: Do not use PVA packets in a tank mix with products that contain boron or release free chlorine. The resultant reaction of PVA and boron or free chlorine is a plastic that is not soluble in water or solvents. For tank mixing with MOVENTO, WSP packaged product user must carefully follow the label directions provided on those product labels.



2.75"
COPY AREA

SPECIFIC CROP DIRECTIONS
CROP USE DIRECTIONS

Apply specified dosage of MOVENTO early in the infestation as the population begins to develop or at early threshold for the target insect pest. Apply higher dosages specified within the crop specific sections when applied as a preventive application, for moderate to heavy insect pressure, or where longer residual control is desired. Degree of efficacy against labeled pests will be determined, in part, by the stage of pest development at application and infestation level of those pests.


Apply in adequate water for uniform coverage. For tree and vine crops, apply in a minimum of 50 GPA for conventional ground airblast sprayer, 30 GPA for high air velocity, low volume or air curtain sprayers, 10 GPA for aerial application; rates for tree and vine crops are based on full-size mature trees and vines. For vegetable and potato crops, apply in a minimum of 15 GPA by ground and 5 GPA by aerial application. MOVENTO may also be applied through overhead irrigation systems as designated in the CHEMIGATION section of this label under *Chemigation Application Instructions*.

25

1.75"

2.25"

3.25"



MOVENTO must be tank-mixed with a spray adjuvant / additive having spreading and penetrating properties to maximize leaf uptake and systemicity of the active ingredient within treated plants; please contact your local Bayer CropScience representative or PCA for specific recommendations by crop. However, the use of Induce[®] adjuvant in combination with MOVENTO on grape, pome fruits, and stone fruits is prohibited when fruit is present due to adverse plant compatibility on harvested commodities. The tank-mixture of MOVENTO with an adjuvant / additive having sticking properties or crop protection product formulations containing built-in stickers have been shown to interfere with leaf uptake and should be avoided. Sufficient leaf tissue must be present for uptake and translocation of this product; due to this requirement, do not apply prior to petal-fall on pome fruits, stone fruits, and tree nut crops.

It is widely known that tank mixtures and/or sequential treatments of horticultural spray oil with Captan and/or sulfur may cause adverse plant compatibility in tree and vine crops; including MOVENTO in this tank mix and/or sequential treatment scenario is not recommended.

← 2.75" →
COPY AREA

↑ 1.75" ↓

BANANA and PLANTAIN (HI, PR, FL only)		
Pests Controlled	Product Rate	
	(fl oz/A)	(lb ai/A)
Aphids	10.0 - 16.0	0.16 - 0.25

Foliar Application Restrictions:
Pre-Harvest Interval (PHI): **1 day(s)**
Minimum interval between applications: **14 days**
Maximum MOVENTO allowed per crop season: **80 fl oz/A**
Maximum spirotetramat per crop season: **1.25 lb ai/A**

↑ 2.25" ↓

← 3.25" →

2.75"
COPY AREA

1.75"


Pests Controlled		Product Rate	
Aphids	Whiteflies	(fl oz/A)	(lb ai/A)
Swede midge		4.0 - 5.0	0.06 - 0.08
Pests Suppressed			
Diamondback moth			
Nematodes			
Onion thrips (larvae)			

2.25"

28

(continued)

3.25"



← 2.75" →
COPY AREA

↑ 1.75" ↓


BULB VEGETABLES
Crops of Crop SubGroup 3-07A Including: Daylily (bulb), Fritillaria (bulb), Garlic (bulbs of common, great-headed, Serpent), Lily (bulb), Onion (bulbs of common, Chinese, Pearl, potato onion), Shallot (bulb), plus cultivars, varieties, and/or hybrids of these.
Crops of Crop Subgroup 3-07B Including: Chinese Chive (fresh leaves), Chive (fresh leaves), Elegans hosta, Fritillaria (leaves), Kurrat, Leek (Allium porrum, Lady's, Wild), Onion (Beltsville bunching, fresh, green, macrostem, tree [tops], Welsh [tops]), Shallot (fresh leaves), plus cultivars, varieties, and/or hybrids of these.

Pests Controlled	Product Rate	
	(fl oz/A)	(lb ai/A)
Onion thrips (larvae)	5.0	0.08

(continued)

↑ 2.25" ↓

← 3.25" →



2.75"
COPY AREA

1.75"

BUSHBERRY SUBGROUP		Product Rate	
LOW GROWING BERRY SUBGROUP		(fl oz/A)	(lb ai/A)
Crops of Crop Subgroups 13-07B and 13-07H including: Aronia berry, Bearberry, Bilberry, Blueberry (highbush and lowbush), Chilean guava, Cloudberry, Cranberry, Currant (black, buffalo, native, and red), Elderberry, European barberry, Gooseberry, Edible honeysuckle, Jostaberry, Juneberry, Muntries, Lingonberry, Partridgeberry, Salal, Sea buckthorn, and cultivars, varieties, and/or hybrids of these.			
Pests Controlled		Product Rate	
Aphids	Cranberry Tipworm	8.0 - 10.0	0.13 - 0.16
Blueberry Gall Midge	Thrips (larvae)		
Pests Suppressed		Product Rate	
Blueberry Maggot		10.0	0.16
Leathoppers			

2.25"

3.25"

2.75"
COPY AREA

Foliar Application Restrictions:
Pre-Harvest Interval (PHI): **7 days**
Minimum interval between applications: **7 days**
Maximum MOVENTO allowed per calendar year: **30 fl oz/A**
Maximum spirotetramat per crop season: **0.47 lb ai/A**
Do not apply until after petal fall

Pests Controlled	Product Rate	
	(fl oz/A)	(lb ai/A)
Aphids		
Whiteflies	5.0	0.08

Foliar Application Restrictions:
Pre-Harvest Interval (PHI): **1 day**
Minimum interval between applications: **7 days**
Maximum MOVENTO allowed per crop season: **10 fl oz/A**
Maximum spirotetramat per crop season: **0.16 lb ai/A**

33



1.75"

2.25"

3.25"



CHRISTMAS TREE PLANTATIONS		
Pests Controlled	Product Rate	
	(fl oz/A)	(lb ai/A)
Aphids (including root aphids and Adelgids)	5.0 - 10.0	0.08 - 0.16
Scales		
Foliar Application Restrictions: Minimum interval between applications: 14 days Maximum MOVENTO allowed per calendar year: 20 fl oz/A Maximum spirotetramat per calendar year: 0.31 lb ai/A		



2.75"
COPY AREA

1.75"

Pests Controlled		Product Rate	
		(fl oz/A)	(lb ai/A)
Aphids	Citrus snow scale	8.0 - 10.0	0.13 - 0.16
Asian citrus psyllid	Citrus thrips		
California red scale	Florida red scale		
Citrus leafminer	Mealybugs		
Citrus bud mite	Pink citrus rust mite		
Citrus red mite	Purple scale		
Citrus rust mite	Texas citrus mite		
(silver mite)	Whitflies		

35

(continued)

2.25"

3.25"

2.75"
COPY AREA

1.75"

Pests Suppressed		Product Rate	
Black scale	Cottony cushion scale	(fl oz/A)	(lb ai/A)
Brown scale	Nematodes	8.0 - 10.0	0.13 - 0.16
Citricola scale			

Foliar Application Restrictions:
 Pre-Harvest Interval (PHI): **1 day(s)**
 Minimum interval between applications: **21 days**
 Maximum MOVENTO allowed per calendar year: **20 fl oz/A**
 Maximum spirotetramat per calendar year: **0.31 lb ai/A**
 Do not apply this product within 10 days prior to bloom, during bloom, or until petal fall is complete.
 For production areas in Alabama, Florida, Georgia, Louisiana, Mississippi, South Carolina, and Texas, optimum control of target pests is obtained at application volumes up to 250 gpa; application volumes in excess of 350 gpa must be avoided.
 Do not apply non-ionic surfactants in tank mix combination with MOVENTO on white grapefruit.

36

2.25"

3.25"

← 2.75" →
COPY AREA

↑ 1.75" ↓

COFFEE		
Pests Controlled	Product Rate	
	(fl oz/A)	(lb ai/A)
Green Scale	8.0 - 10.0	0.13 - 0.16
Foliar Application Restrictions: Pre-Harvest Interval (PHI): 14 day(s) Minimum interval between applications: 21 days Maximum MOVENTO allowed per calendar year: 30 fl oz/A Maximum spirotetramat per calendar year: 0.47 lb ai/A		

↑ 2.25" ↓

← 3.25" →

2.75"

COPY AREA

1.75"

2.25"

3.25"

Pests Controlled		Product Rate	
		(fl oz/A)	(lb ai/A)
Aphids	Tomato russet mite	4.0 - 5.0	0.06 - 0.08
Broad mite	Whiteflies		
Psyllids			
Pests Suppressed			
Leafminers	Western flower thrips		
Nematodes	(larvae)		
Twospotted spider mite			

FRUITING VEGETABLES
Crops of Crop Group 8-10 Including: African eggplant, Bush tomato, Cocona, Currant tomato, Eggplant, Garden huckleberry, Goji berry, Groundcherry, Martynia, Naranjilla, Okra, Pea eggplant, Pepino, Pepper (bell and nonbell*), Roselle, Scarlet eggplant, Sunberry, Tomatillo, Tomato, Tree tomato, including cultivars, varieties and/or hybrids of these commodities.
 *including cayenne, chili pepper, habanero, jalapeno, poblano, pimiento, and serrano.

← 2.75" →
COPY AREA

↑ 1.75" ↓

Foliar Application Restrictions:
 Pre-Harvest Interval (PHI): **1 day(s)**
 Minimum interval between applications: **7 days**
 Maximum MOVENTO allowed per crop season: **10 fl oz/A**
 Maximum spirotetramat per crop season: **0.16 lb ai/A**

GLOBE ARTICHOKE		
Pests Controlled	Product Rate	
	(fl oz/A)	(lb ai/A)
Aphids	5.0 - 8.0	0.08 - 0.13

Foliar Application Restrictions:
 Pre-Harvest Interval (PHI): **3 day(s)**
 Minimum interval between applications: **7 days**
 Maximum MOVENTO allowed per crop season: **32 fl oz/A**
 Maximum spirotetramat per crop season: **0.5 lb ai/A**

39

↑ 2.25" ↓

← 3.25" →


← 2.75" →
COPY AREA

↑ 1.75" ↓

HOPS		
Pests Controlled	Product Rate	
	(fl oz/A)	(lb ai/A)
Hop aphid	5.0 - 6.0	0.08 - 0.09
Twospotted spider mite		
Foliar Application Restrictions: Pre-Harvest Interval (PHI): 7 day(s) Minimum interval between applications: 14 days Maximum MOVENTO allowed per calendar year: 12.5 fl oz/A Maximum spirotetramat per calendar year: 0.2 lb ai/A		

↑ 2.25" ↓

← 3.25" →



2.75"
COPY AREA


1.75"

Pests Controlled		Product Rate	
Aphids	Whiteflies	(fl oz/A)	(lb ai/A)
Pests Suppressed		4.0 - 5.0	0.06 - 0.08
Diamondback moth	Nematodes		
Leafminers	Western flower thrips (larvae)		
Foliar Application Restrictions: Pre-Harvest Interval (PHI): 3 day(s) Minimum interval between applications: 7 days Maximum MOVENTO allowed per crop season: 10 fl oz/A Maximum spirotetramat per crop season: 0.16 lb ai/A			

2.25"

42



3.25"





Pests Controlled		Product Rate	
Aphids	Whiteflies	(fl oz/A)	(lb ai/A)
Pests Suppressed		4.0 - 5.0	0.06 - 0.08
Leafminers	Twospotted spider mite		
Melon thrips (larvae)	Western flower thrips (larvae)		
Nematodes			

Foliar Application Restrictions:
 Pre-Harvest Interval (PHI): **1 day(s)** (edible podded and succulent beans and peas); **7 day** (dry shelled beans and peas).
 Minimum interval between applications: **7 days**
 Maximum MOVENTO allowed per crop season: **10 fl oz/A**
 Maximum spirotetramat per crop season: **0.16 lb ai/A**



← 2.75" →
COPY AREA

↑ 1.75" ↓

PINEAPPLE		
Pests Controlled	Product Rate	
	(fl oz/A)	(lb ai/A)
Mealybugs	10.0	0.16
	Foliar Application Restrictions: Pre-Harvest Interval (PHI): 1 day(s) Minimum interval between applications: 14 days Maximum MOVENTO allowed per crop season: 20 fl oz/A Maximum spirotetramat per crop season: 0.31 lb ai/A	

↑ 2.25" ↓

← 3.25" →

2.75"
COPY AREA

1.75"


2.25"

3.25"

Pests Controlled		Product Rate	
		(fl oz/A)	(lb ai/A)
Aphids (including Woolly Apple Aphid)	Pear psylla Pear rust mite San Jose scale	6.0 - 9.0	0.09 - 0.14
Apple rust mite Mealybugs	Whiteflies		
Pests Suppressed			
Apple gall midge Codling moth European red mite Micro-lepidoptera leafminers	Pear leaf midge Twospotted spider mite White apple leafhopper		

46

(continued)



2.75"
COPY AREA

1.75"

POMEGRANATE		
Pests Controlled	Product Rate	
	(fl oz/A)	(lb ai/A)
Aphids	8.0 - 10.0	0.13 - 0.16
Whiteflies		

Foliar Application Restrictions:
Pre-Harvest Interval (PHI): **1 day(s)**
Minimum interval between applications: **14 days**
Maximum MOVENTO allowed per calendar year: **20 fl oz/A**
Maximum spirotetramat per calendar year: **0.31 lb ai/A**

2.25"

48

3.25"

2.75"

COPY AREA

1.75"

2.25"

3.25"

POTATO and OTHER TUBEROUS AND CORM VEGETABLES			
Crops of Crop Subgroup 1C Including: Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Canna (edible), Cassava (bitter and sweet), Chayote (root), Chufa, Dasheen (taro), Ginger, Leren, Potato, Sweet potato, Tanier, Turmeric, Yam bean, Yam (true)			
Pests Controlled		Product Rate	
Aphids	Whiteflies	(fl oz/A)	(lb ai/A)
Psyllids		4.0 - 5.0	0.06 - 0.08
Pests Suppressed			
Nematodes	Western flower thrips (larvae)		
Twospotted spider mite	Wireworms		

Foliar Application Restrictions:
 Pre-Harvest Interval (PHI): **7 day(s)**
 Minimum interval between applications: **7 days**
 Maximum MOVENTO allowed per crop season: **10 fl oz/A**
 Maximum spirotetramat per crop season: **0.16 lb ai/A**

2.75"
COPY AREA

1.75"


SMALL FRUIT VINE CLIMBING SUBGROUP (Except Fuzzy Kiwifruit)		
Crops of Crop Subgroup 13-07F including: Amur river grape, Gooseberry, Grape, Kiwifruit (hardy), Maypop, Schisandra berry		
Pests Controlled	Product Rate	
	(fl oz/A)	(lb ai/A)
Grape tumid gallmaker	6.0 - 8.0	0.10 - 0.13
Mealybugs		
Phylloxera		
Willamette mite		
Whiteflies		
Pests Suppressed		
European fruit mite		
lecanium scale		
Pacific mite		
Twospotted spider mite		
Nematodes		

2.25"

50

(continued)

3.25"



2.75"
COPY AREA

1.75"

STONE FRUITS		
Crops of Crop Group 12-12 Including: Apricot; apricot, Japanese; capulin; cherry, black; cherry, Nanking; cherry, sweet; cherry, tart; Jujube, Chinese; nectarine; peach; plum; plum, American; plum, beach; plum, Canada; plum, cherry; plum, Chickasaw; plum, Damson; plum, Japanese; plum, Klamath; plum, prune; plumcot; sloe; cultivars, varieties, and/or hybrids of these		
Pests Controlled	Product Rate	
	(fl oz/A)	(lb ai/A)
Aphids	White peach scale	6.0 - 9.0
Mealybugs	Whiteflies	
San Jose scale		

2.25"

(continued)

3.25"

2.75"
COPY AREA

1.75"

Pests Suppressed		Product Rate	
		(fl oz/A)	(lb ai/A)
Black scale	Nematodes	6.0 - 9.0	0.09 - 0.14
Cherry fruit fly	Spotted Wing Drosophila		
European fruit lecanium scale	Twospotted spider mite		
European red mite			
Foliar Application Restrictions:			
Pre-Harvest Interval (PHI): 7 day(s)			
Minimum interval between applications: 14 days			
Maximum MOVENTO allowed per calendar year: 15.3 fl oz/A			
Maximum spirotetramat per calendar year: 0.24 lb ai/A			
Do not apply until after petal fall.			
For control of San Jose scale, apply immediately after petal fall; under heavy infestation pressure or where difficult control conditions exist, a second application may be necessary.			

2.25"

3.25"

2.75"
COPY AREA

1.75"

SUGAR BEET		
Pests Controlled	Product Rate	
	(fl oz/A)	(lb ai/A)
Bean aphid Root Aphid Whiteflies	5.0 - 9.0	0.08 - 0.14
Pests Suppressed		
Root Maggot Sugarbeet cyst nematode		

Foliar Application Restrictions:
Pre-Harvest Interval (PHI): **28 day(s)**
Minimum interval between applications: **14 days**
Maximum MOVENTO allowed per crop season: **18 fl oz/A**
Maximum spirotetramat per crop season: **0.28 lb ai/A**

54

2.25"

3.25"

2.75"
COPY AREA

1.75"

TREE NUTS			
Crops of Crop Group 14-12 Including: African nut-tree; almond; beechnut; Brazil nut; Brazilian pine; bunya; bur oak; butternut; Cajou nut; candlenut; cashew; chestnut; chinquapin; coconut; coquito nut; dika nut; ginkgo; Guiana chestnut; hazelnut (filbert); heartnut; hickory nut; Japanese horse-chestnut; macadamia nut; mongongo nut; monkey-pot; monkey puzzle nut; Okari nut; Pachira nut; peach palm nut; pecan; pequi; Pili nut; pine nut; pistachio; Sapucaia nut; tropical almond; walnut, black; walnut, English; yellowhorn; cultivars, varieties, and/or hybrids of these			
Pests Controlled			
Product Rate			
Aphids	San Jose scale	(fl oz/A)	(lb ai/A)
Mealybugs	Walnut scale	6.0 - 9.0	0.09 - 0.14
Phylloxera	Whiteflies		

2.25"

(continued)

3.25"

2.75"
COPY AREA

1.75"

2.25"

Pests Suppressed		Product Rate	
		(fl oz/A)	(lb ai/A)
European fruit lecanium scale	Pacific mite Twospotted spider mite	6.0 - 9.0	0.09 - 0.14
Nematodes Olive scale			
Foliar Application Restrictions: Pre-Harvest Interval (PHI): 7 day(s) Minimum interval between applications: 14 days Maximum MOVENTO allowed per calendar year: 21.5 fl oz/A Maximum spirotetramat per calendar year: 0.34 lb ai/A Do not apply until after petal fall. For control of San Jose scale, apply immediately after petal fall; under heavy infestation pressure or where difficult control conditions exist, a second application may be necessary.			

3.25"

2.75"
COPY AREA


1.75"

TROPICAL FRUIT			
Acerola, Avocado, Black sapote, Canistel, Feijoa, Jaboticaba, Guava, Longan, Lychee, Mamey sapote, Mango, Papaya, Passionfruit, Persimmon, Pulasan, Rambutan, Sapodilla, Spanish lime, Star apple, Starfruit, Wax jambu, White sapote (<i>Casimiroa</i> spp.)			
Pests Controlled			
Product Rate			
Aphids	Scales	(fl oz/A)	(lb ai/A)
Avocado thrips	Whiteflies	8.0 - 10.0	0.13 - 0.16
Mealybugs			



2.25"

(continued)

3.25"



Pests Suppressed		Product Rate	
		(fl oz/A)	(lb ai/A)
Avocado brown mite	Persea mite	8.0 - 10.0	0.13 - 0.16
Papaya leaf edge-roller mite	Twospotted spider mite		
Foliar Application Restrictions: Pre-Harvest Interval (PHI): 1 day(s) Minimum interval between applications: 14 days Maximum MOVENTO allowed per calendar year: 25 fl oz/A Maximum spirotetramat per calendar year: 0.39 lb ai/A			



2.75"

COPY AREA


1.75"

WATERCRESS		
Pests Controlled	Product Rate	
	(fl oz/A)	(lb ai/A)
Aphids	4.0 - 12.5	0.06 - 0.2
	Foliar Application Restrictions: Pre-Harvest Interval (PHI): 3 day(s) Minimum interval between applications: 7 days Maximum MOVENTO allowed per crop season: 25 fl oz/A Maximum spirotetramat per crop season: 0.39 lb ai/A Do not plant or replant any crop not listed on this label within 260 days after the last application.	

2.25"

59

3.25"



2.75"
COPY AREA

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide storage


MOVENTO is packaged in poly-ethylene containers. Do not allow product or containers to freeze. Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking, invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer CropScience Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer CropScience Emergency Response Telephone No. is (800) 334-7577, or contact Chemtrec at (800) 424-9300.

1.75"



2.25"

3.25"



Pesticide disposal
Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container handling
Rigid, Non-refillable containers (equal to or less than 5 gallons)
Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.



(continued)






Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.
Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill.

Induce is a registered trademark of Helena Holding Company

62

Bayer





BASE LABEL

.8125"

2.875"

.5625"

COPY AREA

1.75"

NET CONTENTS: 1 QT. (32 FL. OZ.)

2.25"

4.25"

MOVENTO®

GROUP 23 INSECTICIDE

For Agricultural Use Only: For control of listed insects on certain tree, tropical fruits, vine, and vegetable crops.

ACTIVE INGREDIENT: Spirotetramat: cis-3-(2,5-dimethylphenyl)-8-methoxy-2-oxo-1-azaspiro [4.5]dec-3-en-4-yl-ethyl carbonate 22.40%

OTHER INGREDIENTS: 77.60%

MOVENTO contains 2.00 pounds TOTAL: 100.00% spirotetramat per U.S. gallon (240 grams AI/liter)

EPA Reg. No. 264-1050

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

For **MEDICAL** And **TRANSPORTATION** Emergencies
ONLY Call 24 Hours A Day 1-800-334-7577
For **PRODUCT USE** Information Call 1-866-99BAYER
(1-866-922-2937)

Please refer to booklet for additional precautionary statements and directions for use.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

- Harmful if swallowed or absorbed through skin.
- Causes moderate eye irritation.
- Avoid contact with skin, eyes, or clothing.
- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.
- Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Read the entire label before using this product.

Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
US61384098B 170628B 09/17