

**FIFRA Section 2(ee) Recommendation**

**DISARM™480 SC  
FUNGICIDE  
EPA Reg. No. 66330-64**

**This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U. S. Environmental Protection Agency**

**FOR DISTRIBUTION ONLY IN STATES WHERE DISARM™ 480 SC IS CURRENTLY REGISTERED**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read and follow all applicable directions for use, precautions and limitations on the product label attached to the container for **DISARM™ 480 SC Fungicide**. Also observe all directions, precautions and limitations on the labeling of all other products mentioned below. This recommendation must be in the possession of the user at the time of application.

**ANTHRACNOSE AND GRAY LEAF SPOT**

**DISARM™ 480 SC** will provide control of Anthracnose (*Colletotricum graminicola*) and Gray leaf spot (*Pyricularia grisea*). Apply 8 – 16 fl. oz./A (0.18 – 0.36 fl. oz./1,000 sq. ft.) on a 21 day schedule when conditions are favorable for disease. Under heavy disease pressure, apply 16 fl. oz./A (0.36 fl. oz./1,000 sq. ft.) or see directions below for alternating fungicides to control turf diseases.

**ALTERNATE SPRAY PROGRAM**

For control of Anthracnose (*Colletotricum graminicola*), Gray leaf spot (*Pyricularia grisea*), Brown Patch (*Rhizoctonia solani*) and Dollar spot (*Sclerotinia homoeocarpa*), apply **DISARM™ 480 SC** at a rate of 8 - 16 fl. oz./A (0.18 - 0.36 fl. oz./1,000 sq. ft.) on a 28 day schedule alternated with a DMI fungicide (Fungicide Group 3) such as **triadimefon** or **propiconazole** on a 28 day schedule between each **DISARM™ 480 SC** application. When dollar spot is **NOT** a problem, **DISARM™ 480 SC** can be alternated with other products labeled for the target diseases except those in the strobilurin class of chemistry (Fungicide Group 11).

For control of Pythium blight (*Pythium aphanidermatum*) apply **DISARM™ 480 SC** at 12 - 16 fl. oz./A (0.28 - 0.36 fl. oz./1,000 sq. ft.) every 21 days in alternation with another fungicide labeled for the control of Pythium blight, between each **DISARM™ 480 SC** application.

**ARYSTA LIFESCIENCE NORTH AMERICA CORPORATION  
15401 Weston Parkway, Suite 150  
Cary, NC 27513  
[www.arystalifescience.com](http://www.arystalifescience.com)**

DISARM is a trademark of Arysta LifeScience North America Corporation

Issue date: April 27, 2006