
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1571B-S2 1116

GROUP	U15	FUNGICIDE
--------------	------------	------------------

Orondis® Gold 200

Fungicide

This supplemental label expires on 11/14/19 and must not be used or distributed after this date.

Active Ingredient:	
Oxathiapiprolin (CAS No. 1003318-67-9).....	18.7%
Other Ingredients:	81.3%
Total:	100.0%

KEEP OUT OF REACH OF CHILDREN.

EPA Reg. No. 100-1571

All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed. Before using Orondis Gold 200 as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

The Syngenta logo, featuring the word "syngenta" in a lowercase, sans-serif font with a registered trademark symbol. A stylized leaf icon is positioned above the letter "g".

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ROTATIONAL CROP RESTRICTIONS

The following crops may be planted at the specified interval following application of Orondis.

Crop, Crop Group, or Subgroup	Plant-back Interval (in Days)
Tuberous and Corm Vegetables (Subgroup 1C)	0
Bulb Vegetables (Group 3-07)	
Leafy Greens (Crop Subgroup 4-16A)	
Brassica, Head and Stem (Crop Group 5-16)	
Peas, Edible-Podded	
Peas, Succulent Shelled	
Fruiting Vegetables (Group 8-10)	
Cucurbit Vegetables (Group 9)	
Caneberry (Crop Subgroup 13-07A)	
Strawberries	
Oilseed (Group 20)	
Stalk and Stem Vegetables (Crop Subgroup 22A)	
Ginseng	
Tobacco	
Cereals (Group 15,16)	30
Grass animal feeds (Group 17)	
Herbs and Spices (Group 19)	180
Legume Vegetables, except succulent shelled and edible-podded peas	
Non-grass Animal feed (Group 18)	
Peanuts	
All other crops not listed	

CROP USE DIRECTIONS

Caneberry, Crop Subgroup 13-07A

Crops (including all cultivars, varieties, and/or hybrids of these)			
Blackberry Loganberry		Raspberry, red and black Wild raspberry	
Target Disease	Rate	Application Timing	Use Directions
Phytophthora root rot (<i>Phytophthora</i> spp.)	4.8 - 9.6 fl oz/A	<p>Make first application before plants start to grow in the spring.</p> <p>Make a second application during a period favorable for root growth and at least 7 days after the first application.</p>	<p>Apply as a banded, soil-directed spray in a minimum of 20 gal/A of water. Direct the spray along each side of the crop row and direct the application to the soil, near and under the lower leaves.</p> <p>Use a higher rate in the listed rate range for moderate to severe infections.</p> <p>The application is to be made at the per acre rate concentrated in a band.</p> <p>Apply ¼ - ½ inch of water after application, either by overhead sprinkler irrigation or garden hose drench on the row.</p>
<p>Resistance Management:</p> <ul style="list-style-type: none"> • Make no more than 2 sequential applications before rotating to a fungicide with a different mode of action. • When 3 or more applications are made, use Orondis (or other oxathiapiprolin-containing product) in no more than 33% of the applications, or a maximum of 4 applications, whichever is fewer. • Do not make more than two applications per year. 			
USE RESTRICTIONS			
<ol style="list-style-type: none"> 1) Refer to Use Restrictions section of the booklet for additional product use restrictions. 2) Maximum Single Application Rate: 9.6 fl oz/A/application (0.12 lb ai/A of oxathiapiprolin-containing products) 3) Minimum Application Interval: 7 days 4) Maximum Annual Rate: 19.2 fl oz/A/year (0.25 lb ai/A/year of oxathiapiprolin-containing products) 5) Maximum Number of Applications: Do not make more than two applications per year. 6) Pre-harvest Interval (PHI): 1 day 			

Orondis® and the Syngenta logo are trademarks of a Syngenta Group Company

©2017 Syngenta