

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

FIFRA 24(c) Special Local Need Label



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Lorsban[®] Advanced

EPA Reg. No. 62719-591

**EPA 24(c) Special Local Need Registration SLN CA-080011
For Distribution and Use Only in the State of California**

Control of Argentine Ants in Citrus

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Lorsban[®] Advanced insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Lorsban Advanced according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Lorsban Advanced.

Directions for Use

Argentine Ants Only (Spray Method):

Mix 3 to 8 fl oz of Lorsban Advanced per gallon of water. Apply 2.5 gallons of a 1% to 3% chlorpyrifos solution per acre (approximately 100 trees) as a ground broadcast spray directed to the orchard floor around the base of the tree. Apply as a coarse stream with a hand-held sprayer and spray uniformly to the soil surface, allowing some spray mixture to soak into the soil. Use a 1% ai chlorpyrifos solution (3 fl oz per gallon) if less than 50 ants per minute are going up a tree. If a spreader/sticker is needed, use it at a rate of 1 fl oz per 2.5 gallons of water.

Specific Use Restrictions for Argentine Ants Only (Spray Method):

- **Preharvest Interval:** Do not apply last treatment within 28 days before harvest.
- Do not make a second application of Lorsban Advanced or other product containing chlorpyrifos within 10 days of the first application.
- Do not allow meat or dairy animals to graze in treated areas.
- Maximum single application rate is 1 lb ai chlorpyrifos per acre.
- Do not apply more than 3 lb ai chlorpyrifos per acre per year.
- Do not make more than 3 applications of Lorsban Advanced per year.
- Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 5 days unless PPE required for early entry is worn.
- Growers may not treat the same citrus crop under the requirements of any SLN for Lorsban-4E.

Argentine Ants (Fiberglass Band Method)

Mix 3 to 8 fl oz of Lorsban Advanced per gallon of water. Apply 10 gallons of a 1% to 3% chlorpyrifos solution per acre (approximately 100 trees). Apply as a coarse stream with a hand-held sprayer and spray thoroughly and uniformly to wet an absorbent insulation grade fiberglass band securely wrapped around the trunk of skirt-pruned trees. Fiberglass material should contain a "backing" material that faces toward the tree trunk. If a spreader/sticker is needed, use it at a rate of 1 fl oz per 2.5 gallons of water.

Specific Use Restrictions Argentine Ants (Fiberglass Band Method):

- Do not allow meat or dairy animals to graze in treated areas.
- Maximum single application rate is 2.5 lb ai chlorpyrifos per acre
- Do not make more than 3 applications of Lorsban Advanced per year
- Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 5 days unless PPE required for early entry is worn.
- Growers may not treat the same citrus crop under the requirements of any SLN for Lorsban-4E.

Expiration Date: December 31, 2019

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R368-104

Approved: 10/16/14

Replaces R368-048