




# Dimension<sup>®</sup> 270G Specialty Herbicide

Now with BIODAC



A consistent sized granular carrier that is virtually dust free and environmentally friendly.

The BIODAC<sup>®</sup> trademark and logo are owned by Kadant Gran Tek, Inc.

## Turf and Landscape Ornamental Granular Pre-emergent Herbicide

- Provides control of listed annual grasses and broadleaf weeds in Ornamental turf (including but not limited to golf course fairways, roughs, tee boxes)
- Commercial sod farms
- Field-grown nursery ornamentals
- Container grown nursery ornamentals
- Ornamentals landscape areas
- Established lawns
- Non-crop land and industrial sites
- For use on plants intended for aesthetic purposes or climatic modification and being grown in interior landscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

In New York State this product may only be used by professional applicators at no more than 185 pounds per acre (4.25 lbs. per 1,000 sq. ft.), (0.5 pounds active ingredient per acre) per year and is prohibited from use in Nassau and Suffolk Counties.

### ACTIVE INGREDIENT:

Dithiopyr (CAS 97886-45-8) ..... \*0.27%

OTHER INGREDIENTS: ..... 99.73%

TOTAL : ..... 100.00%

\*This product contains 0.135 pounds of the active ingredient, dithiopyr, per 50 pound bag.

EPA Reg. No. 7001-375

EPA Est. No. 41124-WI-1

This Product is intended for professional use.

Notice: Read The Entire Label before using this product. Use only according to label instructions. Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions. If Conditions of Sale and Warranty are not acceptable, return at once unopened.

A 50 Lb. Bag Treats Up To 11,764 Sq. Ft.

At the 185 Lb./Acre Rate for Crabgrass Control

KEEP OUT OF REACH OF CHILDREN

## CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

### FIRST AID

- IF IN EYES:**
- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
  - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
  - Call a poison control center or doctor for treatment advice.

### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.

bestfertilizer.com

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes or clothing.

## PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category B on an EPA chemical-resistance category selection chart.

Applicators and other handlers who handle this product for any use covered by the Worker Protection Standard (40 CFR Part 170) must wear:

- Long-sleeved shirt and long pants
- Socks and Shoes

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

This product is toxic to fish. Drift and runoff may adversely affect aquatic organisms. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Read all directions carefully before applying this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protection equipment (PPE), restricted-entry interval and notification to workers. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours.

For early entry into treated areas that is permitted under the Worker protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear:

- Long-sleeved shirt and long pants
- Socks and shoes

## NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, nurseries or greenhouses. Keep unprotected persons out of treated areas until dusts have settled.

Spread The Word



## SPREADER SETTINGS

Pounds / 1,000 Sq. Ft.	Half Rate Application			Full Rate Application		
	1.15 lbs.	1.7 lbs.	2.1 lbs.	2.3 lbs.	3.4 lbs.	4.2 lbs.
BEST PRO						
SPYKER / BEST 34, 44, 64						
SCOTTS R8A						
LELY TYPE W						
EARTHWAY						

## USE DIRECTIONS FOR TURF

Dimension 270G Specialty Herbicide provides season-long control of crabgrass and control or suppression of other listed annual grasses and broadleaf weeds in established lawns and ornamental turfs, (including golf course fairways, roughs, and tee boxes). When used according to label directions, this product will not harm nearby established ornamentals identified as "Tolerant Ornamentals" elsewhere in this label.

### Precautions for Turf

- **Do not** use this product on golf greens.
- **Do not** apply this product to Colonial bentgrass.
- For best results, cultural practices that disturb the soil, such as core, spike, hydro-aerification, or verticutting, and topdressing should be done before application of this product. This product should not be applied until turfgrasses have recovered from stresses related to these or other mechanical operations. Irrigation or rain fall within 7 days after application will improve the performance of this product.
- **Reseeding or Sprigging:** Reseeding, overseeding, or sprigging of turfgrasses into areas that have been previously treated with this product should be delayed until 12 weeks from the time of application. If overseeding with perennial ryegrass, delay overseeding at least 8 weeks after application. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed-to-soil contact.
- This product may be used on seeded, sodded, or sprigged turfgrass that is well established. Allow the turf to develop a good root system and uniform stand before application. Use of this product on turfgrass not well established, weakened by lack of moisture, chemical injury, insects, disease, or temperature extremes that cause stress will increase the potential for injury. Allow turfgrass weakened by these stresses to recover before application of this product.
- This product may be applied as a single application or as a split application in the spring, summer or fall.
- This product will prevent germination of annual bluegrass.
- This product will not affect established annual bluegrass. If maintenance of annual bluegrass is desired, do not use this product during time of annual bluegrass germination. For AZ, CA, ID, MT, NM, NV, OR, UT, and WA. This product may contribute to thinning or stand reduction in established stands of annual bluegrass.
- **Do not** use clippings from treated turf for mulching around vegetables or fruit trees.
- **Do not** allow domestic animals to feed or forage on the treated turf or clippings from treated turf.

When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

### Cool-Season Grasses

bentgrass, creeping*	<i>Agrostis stolonifer</i>
bluegrass, Kentucky	<i>Poa pratensis</i>
fescue, fine**	<i>Festuca rubra</i>
fescue, tall	<i>Festuca arundinacea</i>
ryegrass, perennial	<i>Lolium perenne</i>

### Warm-Season Grasses

bahiagrass	<i>Paspalum notatum</i>
bermudagrass	<i>Cynodon dactylon</i>
buffalograss	<i>Buchloe dactyloides</i>
carpetgrass	<i>Axonopus affinis</i>
centipedegrass	<i>Eremochloa ophiuroides</i>
kikuyugrass	<i>Pennisetum clandestinum</i>
seashore paspalum	<i>Paspalum vaginatum</i>
St. Augustinegrass	<i>Stenotaphrum secundatum</i>
zoysiagrass	<i>Zoysia japonica</i>

\*DO NOT apply this product to Colonial bentgrass. Use of this product on certain varieties of creeping bentgrass, such as "Carmen", "Cohansey", or "Seaside" will result in undesirable turf injury. Not all varieties of creeping bentgrass have been tested.

\*\*The following fine fescue varieties have been found to be sensitive to this product: "Atlanta", "Banner", "Beauty", "Bligart", "CF-2", "Enjoy", "HF-93", "Highlight", "Ivalo", "Jamestown", "Koket", "Majenta", "Mary", "Pennlawn", "Tamara", "Tatjana", "Waldorf", and "Waldina".

### Sod Production

It is recommended that sod be established for at least six months before a Dimension 270G application is made.

Do not apply this product within 3 months of harvest.

### Other Use Precautions

Early postemergence applications of this product will control crabgrass only if applied prior to the fifth leaf (first tiller) state of growth of crabgrass.

For best results, cultural practices that disturb the soil, such as core-spike, or hydro-aerification, and verticutting, should be done before application of this product.

Do not apply this product until the grass has recovered from these cultural practices. Do not use clippings from treated turf for mulching around vegetables or fruit trees.

Do not graze livestock or feed foliage cut from areas treated with this product.

### Application Equipment and Instructions

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Calibrate application equipment prior to usage, and apply this product uniformly over the turfgrass area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction of application. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

### Preemergence Control of Annual Grasses and/or Annual Broadleaf Weeds

Used as directed, this product will control or suppress the following annual grass and broadleaf weeds when applied prior to their emergence. This product will not control established weeds, except for crabgrass in early stages of development, but only if application is made to crabgrass prior to the 5-leaf stage of growth or first tiller. Applications to crabgrass after initiation of tillering will not provide satisfactory control. All other applications of this product should be made preemergence (prior to germination of target weeds. Use the higher application rates where weed infestations are dense and/or applications are made early and longer residual control is desirable.

## WEEDS CONTROLLED AND APPLICATION RATES

Weeds Controlled	Suggested Use Rate Per Application	
	Pounds/Acre	Pounds/1000 sq ft
barley brome crabgrass, large crabgrass, smooth crabgrass, southern	dallisgrass (seedling) foxtail, giant foxtail, green foxtail, yellow	100 to 150 ** 2.3 to 3.4 *
barnyardgrass chickweed clover, hop crabgrass, tropical crowfootgrass dichondra, Carolina lespedeza, common kikuyugrass Mary's grass marestail medic, black mulberry weed mustard	oats, wild oxalis, buttercup oxalis (creeping woodsorrel) oxalis (yellow woodsorrel) pineappleweed ryegrass, annual ryegrass, perennial sandbur smutgrass speedwell, corn spurge, garden spurge, prostrate spurge, spotted	150 to 185 ** 3.4 to 4.25 *
bluegrass, annual bittercress carpetweed chickweed geranium, Carolina goosegrass henbit	knotweed, prostrate parsley-piert pigweed, redroot purslane, common rocket, London shepherd's-purse	185 ** 4.25 **

\* Use the higher application rates where weed infestations are heavy and/or dense.

\*\* Use of this product at this rate will provide only partial control and/or suppression of listed weeds.

## Poa Annu (annual bluegrass) Control

Apply Dimension® 270G for preemergence control of Poa annua (annual bluegrass) at a rate of 0.38 to 0.5 lbs active ingredient per acre. This is equivalent to 150 to 185 pounds per acre. Use the higher application rates where weed infestations are dense and/or applications are made early and longer residual control is desirable.

- Apply 6 to 8 weeks before overseeding bermudagrass with perennial ryegrass. Specific to perennial ryegrass; not for use on Poa trivialis or bentgrass.
- Minimum seeding rate of perennial ryegrass is 400 lbs per acre.
- Use limited to fairways and roughs
- Perennial varieties of Poa annua (var. repens) may not be controlled as well as the true annual.
- Do not apply earlier than 16 weeks after over-seeding unless injury to the rye grass can be tolerated.
- A follow-up treatment 16 weeks after overseeding offers an early season crabgrass treatment and helps suppress some winter annual broadleaf weeds.

## Control of Crabgrass (Preemergence and early Postemergence Control)

This product provides preemergence control of crabgrass (including the large, smooth, and southern species when applied prior to the emergence of crabgrass in established lawns and ornamental turfs. It can also provide "early postemergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged. However, it is often difficult to see the very small, early stages of crabgrass in well-established lawns and ornamental turfs. Early post-emergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the lawn or turf. The benefit of Dimension® 270G's additional, early post-emergence activity is that (compared to strictly preemergence crabgrass products) it can give the user a 2-8 week longer period of time (depending on climatic conditions and crabgrass growth rates) to make applications and still control crabgrass.

### Restrictions:

- Do not apply more than 185 pounds of this product per acre per application and no more than 3 applications per year (555 pounds) of this product per acre per year.
- In New York State, this product may only be used by Commercial Applicators at no more than 185 pounds per acre (4.25 lb/1000 sq ft), (0.5 lb active ingredient per acre) per year; and is prohibited from use in Nassau and Suffolk Counties.

## NONCROPLAND AND INDUSTRIAL SITES

Dimension 270G may be applied for preemergence control of listed annual grasses and broadleaf weeds in terrestrial noncrop areas such as farm yards, fence rows, highway or roadside utility and railroad rights-of way, airports, recreation areas, campgrounds, and industrial sites such as lumber yards, tank farms, and storage areas. Applied preemergence, Dimension 270G controls weeds as they germinate. This product will not control established weeds. Make applications prior to germination of target weeds or to bare ground. The best weed control is obtained when applications are made to soil that is free of clods, weeds and debris such as leaves. Prior to making an application, control existing vegetation by cultivation, hand weeding, or use of a postemergence herbicide.

To be effective, Dimension 270G must be activated by ½ inch of rainfall or irrigation prior to germination of target weeds. Once the treatment is activated, avoid mixing of the soils surface that will expose treated soil.

### Application rates:

Pounds Product/Acre	Ounces Product/1000 Square Feet	Ounces (Grams) Product/100 Square Feet
185	68	6.8 (193 grams)

Sequential applications may be made at 3 to 4 month intervals for extended preemergence weed control. Do not exceed maximum use rates per year.

### Maximum Use Rates

Do not apply more than 185 lb. of this product per acre per application and no more than 3 applications (555 lbs.) of this product per acre per year.

In New York State, this product may only be used by Commercial Applicators at no more than 185 lbs. per acre (4.25 ounces/1000 sq.ft.) (0.5 lb. active ingredient per acre) per year, and is prohibited from use in Nassau and Suffolk Counties.

### Use Precautions for Noncrop Areas

- Do not apply when weather conditions favor drift to non-target areas. This product may injure foliage of non-target plants.
- Do not graze livestock or feed forage cut from areas treated with this product.
- For ornamentals within non-crop areas, apply only after transplanting when soil around roots has been thoroughly settled by rainfall or irrigation or injury may result.

## ORNAMENTALS (Landscape, Field Grown, and Container Grown)

DIMENSION® 270G provides preemergence control of listed annual grass and broadleaf weeds in ornamental plantings. Ornamental plants on this label have shown tolerance to this product when applied as directed. It is impossible, however, to evaluate tolerance for all ornamental plant species or varieties or under all possible growing conditions. Under your growing conditions, a limited area involving only a few plants should be treated for tolerance evaluation prior to large-scale applications.

### Use Rates

Apply DIMENSION® 270G with a properly calibrated spreader which will assure accurate, even particle distribution. This product may be applied as a single application or as a split application. Refer to the following table for product application rates when treating a small area.

Pounds Product/Acre	Ounces Product/1000 Square Feet	Ounces (Grams) Product/100 Square Feet
185	68	6.8 (193 grams)

### Other Use Precautions

Early postemergence applications of this product will control crabgrass only if applied prior to the fifth leaf (first tiller) state of growth of crabgrass.

For best results, cultural practices that disturb the soil, such as core-spike, or hydro-aerification, and verticutting, should be done before application of this product.

- Do not apply this product until the grass has recovered from these cultural practices.
- Do not use clippings from treated turf for mulching around vegetables or fruit trees.
- Do not graze livestock or feed foliage cut from areas treated with this product.

### Application Timing and Use

This product is a preemergence herbicide that will not control established weeds. Make applications prior to weed seed germination or to bare ground. The best weed control is obtained when applied to soil that is free of clods, weeds and debris such as leaves. Prior to making an application, control existing vegetation by hand weeding, cultivation or with a postemergence herbicide. To be effective, Dimension® 270G must be activated by 1/2 inch or more of rainfall or irrigation prior to germination of target weeds. Once the treatment is activated, avoid excessive soil disruption that may break down the herbicide barrier. Minimal surface disruption such as raking should not break down the herbicide barrier.

### Weed Controlled

Used as directed, DIMENSION® 270G will control or suppress the following annual grass and broadleaf weeds when applied prior to their emergence. This product will not control established weeds, except for crabgrass in early stages of development, but only if application is made to crabgrass prior to the 5-leaf stage of growth or first tiller. Therefore, the area to be treated should be free of weeds prior to application. Repeat applications may be made during the growing season at 3 to 4 month intervals to maintain weed control.

### Weed controlled or suppressed by Dimension® 270G when applied at 185 lbs. per acre:

#### Annual Grasses

barley <i>Hordeum spp.</i>	foxtail, giant <i>Setaria faberi</i>
barnyard grass <i>Echinochloa crus-galli</i>	foxtail, green <i>Pennisetum glaucum</i>
bluegrass, annual <i>Poa annua</i>	foxail, yellow <i>Pennisetum glaucum</i>
brome <i>Bromus spp.</i>	goosegrass <i>Eleusine indica</i>
crabgrass, large <i>Digitaria sanguinalis</i>	oats, wild <i>Avena fatua</i>
crabgrass, Southern <i>Digitaria ciliaris</i>	ryegrass (annual & perennial) <i>Lolium spp.</i>
crowfootgrass <i>Dactyloctenium aegyptium</i>	sandbur <i>Cenchrus spp.</i>
dallisgrass (seedling) <i>Paspalum dilatatum</i>	smutgrass <i>Sporobolus indicus</i>

#### Broadleaf Weeds

bittercress <i>Cardamine spp.</i>	parsley-piert <i>Alchemilla arvensis</i>
--------------------------------------	---

carpetweed <i>Mullugo verticillata</i>	pigweed, redroot <i>Amaranthus retroflexus</i>
chickweed <i>Stellaria spp.</i>	pineappleweed <i>Matricaria matricarioides</i>
geranium, Carolina Geranium carolinianum	purslane, common Protulaca oleracea
henbit <i>Lamium spp.</i>	rocket, London <i>Sisymbrium irio</i>
knotweed, prostrate <i>Polygonum saviculare</i>	shepherdspurge <i>Capsella bursa-pastoris</i>
lespedeza, common <i>Lespedeza striata</i>	speedwell, corn <i>Veronica arvensis</i>
marestail <i>Hippuris vulgaris</i>	spurge, garden <i>Euphorbia hirta</i>
medic, black <i>Medicago lupulina</i>	spurge, prostrate <i>Euphorbia humistrata</i>
mulberry weed <i>Fatoua villosa</i>	spurge, spotted <i>Euphorbia maculate</i>
mustard <i>Brassica spp.</i>	woodsorrel, creeping <i>Oxalis corniculata</i>
oxalis, buttercup <i>Oxalis pes-caprae</i>	woodsorrel, yellow <i>Oxalis stricta</i>

### Application Equipment and Instructions

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that will apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate listed above. Apply this product uniformly over the area to be treated. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules. Shortly after application apply overhead irrigation to activate the herbicide and wash Dimension® 270G from plant surface onto soil surface.

### Use Precautions for Ornamental Plantings:

- Only apply this product to established ornamentals as directed in the Tolerant Ornamental table below.
- Do not incorporate this product into soil. Dilution of active ingredient and possible injury to plant roots may occur.
- Do not apply around ornamental plants that have been weakened or are under stress (due to drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides or injury due to insects, nematodes or disease).
- Do not make applications when weather conditions favor drift to non-target areas. This product may injure foliage of non-target plants.
- Do not apply this product on grasses grown for seed.
- Do not graze livestock or feed forage cut from areas treated with this product.
- Do not apply this product in enclosed structures and greenhouses.
- Do not apply another herbicide within 4 weeks of application of this product.
- Do not apply this product to bare roots of ornamental plants as injury may result. Following transplanting, care must be taken that soil or planting mixes have settled firmly through irrigation, rainfall or packing and that there are no cracks that would allow direct contact of this product to the plant roots or plant injury may occur.

### Treatment of Ornamental species not listed on this label:

It is impossible to evaluate tolerance to this product on all ornamental species or varieties or under all growing conditions. Users who wish to use Dimension® 270G on species not listed on this label may determine the suitability for use by treating a small number of plants at an application rate under your growing conditions. Treated plants should be observed for any symptoms of herbicidal injury such as foliar damage, reduced vigor or stand reduction during 30 to 60 days of normal growing conditions to determine if the treatment is non-injurious.

### Restrictions:

Do not apply more than 185 lbs. of this product per acre per application and no more than 3 applications (555 lbs.) of this product per acre per year. In New York State, this product may only be used by Commercial Applicators at no more than 185 pounds per acre (4.25 ounces/1000 sq. ft.), (0.5 lbs. active ingredient per acre) per year, and is prohibited from use Nassau and Suffolk Counties.

### TOLERANT ORNAMENTALS

When applied as directed under the conditions described on this label, ornamentals listed below have shown tolerance in field trials.

		Application to Field / Landscape or Container Grown Ornamentals Noted by a (x)	
Name	Tolerant Cultivars	Container Grown	Field / Landscape Grown
abelia, dwarf <i>(Abelia x grandiflora)</i>	nana		x
abyssinian red banana <i>(Ensete ventricosum)</i>	maureli	x	x
acacia redolens <i>(Acacia redolens)</i>		x	x
Ajuga carpet bugle <i>Ajuga reptans</i> <i>(Ajuga genevensis)</i>	bronze bronze beauty		x x
almond, flowering <i>(Prunus gladiolosa)</i>			x
apple* <i>(Malus pumila)</i>			x
arborvitae <i>(Thuja occidentalis)</i>	nigra pyramidalis smaragh techny woodwardii		x x x x x
arborvitae, dwarf golden <i>(Thuja orientalis)</i>	aurea nana	x	x
ash, green <i>(Fraxinus pennsylvanica)</i>			x
ash, autumn purple <i>(Fraxinus americana)</i>	autumn purple		x
aster, Chinese <i>(Callistephus chinensis)</i>	dwarf queen		x
azalea <i>(Rhododendron spp.)</i>	brilliant buccaneer carror chimes (Belgian) Elsie Lee exbury fashion hardijzer beauty hershey red higasa hinocrimson Holland (hybrid) Marion Lee northern lights orange cup orchid lights snow southern charm		x x x x x x x x x x x x x x x x x x x x
azalea, flame <i>(Rhododendron calendulaceum)</i>			x
azalea, kirishima <i>(Rhododendron obtusum)</i>			x
bamboo, heavenly <i>(Nandina domestica)</i>	compacta nana		x x
barberry, Japanese <i>(Berberis thunbergii)</i>	aurea dwarf pygmy green kobold pygmy red rose glow		x x x x x x
barberry, purple <i>(Berberis thunbergii var atropurpurea)</i>	atropurpurea		x
basket flower <i>(Gaillardia grandiflora)</i>			x
beach grass <i>(Ammophila breviligulata)</i>			x
bearberry (common) <i>(Arctostaphylos uva-ursi)</i>	Massachusetts		x



Name	Tolerant Cultivars	Application to Field / Landscape or Container Grown Ornamentals Noted by a (x)	
		Container Grown	Field / Landscape Grown
gardenia ( <i>Gardenia jasminoides</i> )	mystery radicans white gem	x	x x x
geranium ( <i>Pelargonium x hortorum</i> )			x
gum ( <i>Eucalyptus citriodora</i> )			x
hawthorn ( <i>Crataegus spp.</i> )	cockspur white crimson cloud enchantress Jack Evans Washington white		x x x x x
hawthorn, Indian ( <i>Rhamphiolepis indica</i> )	ballerina	x	x
heather, twisted ( <i>Erica cinerea</i> )	Mediterranean pink		x
hemlock, Canada ( <i>Tsuga canadensis</i> )			x
hibiscus ( <i>Hibiscus spp.</i> )	blue bird brilliant hula girl		x x x
holly ( <i>Ilex x 'Nellie R. Stevens'</i> ) ( <i>Ilex x attenuata</i> )	Nellie R. Stevens fosteri Savannah	x	x x x
holly, blue ( <i>Ilex x meserveae</i> )	blue boy blue girl China girl		x x x
holly, cassine ( <i>Ilex cassine</i> )		x	x
holly, Chinese ( <i>Ilex cornuta</i> )	burfordii needlepoint		x x
holly, Japanese ( <i>Ilex crenata</i> )	compacta hellerie Japanese northern beauty steeds		x x x x
holly, yaupon ( <i>Ilex vomitoria</i> )			x
honeysuckle ( <i>Lonicera xylosteum</i> ) ( <i>Lonicera japonica</i> ) ( <i>Lonicera tatarica</i> )	Claveyí dwarf halliana Canadian white Zabelli		x x x x
hosta ( <i>Hosta spp.</i> ) ( <i>Hosta Lancifolia</i> )	Albo marginata		x x
ice plant ( <i>Carpobrotus edulis</i> )		x	x
ice plant, rosea ( <i>Drosanthemum floribundum</i> )		x	x
ice plant, white trailing ( <i>Delosperma alba</i> )		x	x
impatiens ( <i>Impatiens spp.</i> )			x
iris ( <i>Iris spp.</i> )	dwarf blue wedgewood		x x
ivy, English ( <i>Hedera helix</i> )	Bulgaria thorndale		x x
jasmine, Asian ( <i>Trachelospermum asiaticum</i> )			x
jasmine, star ( <i>Trachelospermum jasminoides</i> )			x

Name	Tolerant Cultivars	Application to Field / Landscape or Container Grown Ornamentals Noted by a (x)	
		Container Grown	Field / Landscape Grown
juniper ( <i>Juniperus spp.</i> )	Arcadia Armstrong bar harbor blue chip blue Pacific blue rug blue star broadmoor buffalo calgary carpet emerald sea emerald spreader endora compacta fruitlandi green gold tip hetzi hughes Manhattan blue parsoni pfitzeriana plumosa Prince of Wales procumbens dwarf San Jose sargent blue sargent green scandia scopulorum moonglow spartan tamariscifolia weberi Youngstown Yukon belle		x x
juniper ( <i>Juniperus chinensis</i> )	gold coast torulosa-Hollywood twisted	x x	x x
juniper ( <i>Juniperus procumbnes nana</i> )	prostrate Japanese garden	x	x
juniper ( <i>Juniperus sabina</i> 'tamariscifolia')	tamarix	x	x
juniper ( <i>Juniperus scopulorum</i> )	scopulorum skyrocket	x	x
king palm ( <i>Archontophoenix cunninghamiana</i> )		x	x
laurel, Carolina cherry ( <i>Prunus caroliniana</i> )	bright n' tight	x	x
laurel, mountain ( <i>Kalmia latifolia</i> )			x
leucothoe ( <i>Leucothoe fontanesiana</i> )			x
ligustrum, Japanese ( <i>Ligustrum japonicum</i> )			x
lily ( <i>agapanthus spp.</i> )	streamline		x
Lily of the Nile ( <i>agapanthus africanus</i> )	Albus Peter Pan		x x
lilyturf ( <i>Liriope muscari</i> )	evergreen giant lilac beauty majestic monroe white silvery sunproof variegata	x x	x x x x x
lilyturf, creeping ( <i>Liriope spicata</i> )			x

Name	Tolerant Cultivars	Application to Field / Landscape or Container Grown Ornamentals Noted by a (x)	
		Container Grown	Field / Landscape Grown
maple, amur ( <i>Acer ginnala</i> )	emerald elf	x	x
maple, Japanese ( <i>Acer palmatum</i> )			x
maple, Norway ( <i>Acer platanoides</i> )			x
maple, red* ( <i>Acer rubrum</i> )	red sunset	x	x
maple, silver ( <i>Acer saccharinum</i> )			x
maple sugar* ( <i>Acer saccharum</i> )			x
marigold ( <i>Tagetes patula</i> )	honeycomb variegata wheeler's dwarf		x x x
mock orange* ( <i>Philadelphus spp</i> )	golden snowflake double white		x x
mondo grass ( <i>Ophiopogon japonicus</i> )			x
moss rose ( <i>Portulaca grandiflora</i> )	sunnyside		x
mountainash ( <i>Sorbus aucuparia</i> )			x
myrtle, crape ( <i>Lagerstroemia indica</i> )	faurei langer muskogee standard pink		x x x x
myrtle, wax ( <i>Myrica californica</i> )			x
narcissus ( <i>Narcissus spp.</i> )			x
oak, laurel ( <i>Quercus laurifolia</i> )			x
oak, pin ( <i>Quercus palustris</i> )			x
oak, red ( <i>Quercus rubra</i> )			x
oak, southern ( <i>Quercus virginiana</i> )			x
oak, willow ( <i>Quercus phellos</i> )			x
oleander ( <i>Nerium oleander</i> )	hardy red petite pink sister agnes		x x x
osteospermum ( <i>Osteospermum fruticosum</i> )	whirligig		x
pachysandra ( <i>Pachysandra terminalis</i> )			x
palm, king ( <i>Archontophoenix cunninghamiana</i> )			x
palm, queen ( <i>Syagrus romanzoffianum</i> )		x	x
pampas grass ( <i>Cortaderia selloana</i> )			x
pansy ( <i>Viola x wittrockiana</i> )			x
paper flower ( <i>Bougainvillea glabra</i> )	Barbara Karst	x	x
peach* ( <i>Prunus persica</i> )			x
periwinkle, dwarf ( <i>Vinca minor</i> )			x

Name	Tolerant Cultivars	Application to Field / Landscape or Container Grown Ornamentals Noted by a (x)	
		Container Grown	Field / Landscape Grown
petunia ( <i>Petunia x hybrida</i> )	picoti		x
photinia, red tip ( <i>Photinia x fraseri</i> )			x
pieris ( <i>Pieris taiwanensis</i> )			x
pieris, Japanese ( <i>Pieris japonica</i> )	mountain fire		x
pine, Afghan ( <i>Pinus eldarica</i> )		x	x
pine, Austrian black ( <i>Pinus nigra</i> )		x	x
pine, Japanese black ( <i>Pinus thunbergii</i> )			x
pine, loblolly ( <i>Pinus taeda</i> )			x
pine, longleaf ( <i>Pinus palustris</i> )			x
pine, mugo or Swiss Mt. ( <i>Pinus mugo</i> )			x
pine, Scotch ( <i>Pinus sylvestris</i> )			x
pine, slash ( <i>Pinus elliottii</i> )			x
pine, Virginia ( <i>Pinus virginiana</i> )			x
pine, white ( <i>Pinus strobus</i> )		x	x
pineapple, guava* ( <i>Feijoa sellowiana</i> )			x
pittosporum, tobira ( <i>Pittosporum tobira</i> )	Wheeler's dwarf	x	x
plumbago, cape ( <i>Plumbago auriculata</i> )	royal cape	x	x
potentilla ( <i>Potentilla fruticosa</i> ) ( <i>Potentilla nepalensis</i> )	abbotswood		x x
privet ( <i>Ligustrum x vicaryii</i> ) ( <i>Ligustrum japonicum</i> )	golden vicary regal texanum yellow tipped	x	x x x x
privet, glossy ( <i>Ligustrum lucidum</i> )			x
Pyracantha or firethorn ( <i>Pyracantha x 'Gnome'</i> ) ( <i>Pyracantha coccinea</i> ) ( <i>Pyracantha koidzumii</i> )	gnome lalandei victory	x	x x x
queen palm ( <i>Arecastrum rammanzoffianum</i> )			x
quince, Japanese* ( <i>Chaenomeles japonica</i> )			x
Rhododendron ( <i>Rhododendron spp</i> )	album Cunningham white PJM purple gem silvery pink		x x x x x
rhododendron, Carolina ( <i>Rhododendron carolinianum</i> )			x
rhododendron, catawba ( <i>Rhododendron catawbiense</i> )			x

Name	Tolerant Cultivars	Application to Field / Landscape or Container Grown Ornamentals Noted by a (x)	
		Container Grown	Field / Landscape Grown
rhododendron, rhodie max-rosebay ( <i>Rhododendron maximum</i> )			x
ribbon grass ( <i>Phalaris arundinacea</i> )			x
rockcress ( <i>Arabis caucasica</i> )	snowcap		x
Rose* ( <i>Rosa banksiae</i> )	luta		x
rose, knockout shrub ( <i>Rosa spp. hybrid</i> )	knockout	x	x
Rosemary* ( <i>Rosmarinus officinalis</i> )			x
rosemary, bog ( <i>Andromeda polifolia</i> )	nana		x
salvia ( <i>Salvia farinacea</i> )	rhea		x
sedum ( <i>Sedum spurium</i> )	dragon blood red red carpet yellow		x x x
smoketree, royal purple ( <i>Cotinus coggygria</i> )	royal purple		x
snapdragon ( <i>Antirrhinum spp.</i> )			x
snowball, common ( <i>Viburnum opulus</i> )	sterile	x	x
sourwood ( <i>Oxydendrum arboreum</i> )			x
spiraea ( <i>Astilbe X arendsii</i> )	fanall		x
spiraea ( <i>Spiraea spp.</i> )	Anthony Waterer red dolchica froebeli pink goldenflame red snowmound white Van Houtte white		x x x x x x
spiraea, garland ( <i>Spiraea x arguta</i> )			x
spruce, Black Hills ( <i>Picea glauca var densata</i> )			x
spruce, Colorado blue ( <i>Picea pungens</i> )	glauca	x	x
spruce, Norway ( <i>Picea abies</i> )			x
spruce, white ( <i>Picea glauca</i> )	conica		x
sweetflag grass ( <i>Acorus calimus</i> )			x
sweetgum ( <i>Liquidambar styraciflua</i> )			x
sycamore, American ( <i>Platanus occidentalis</i> )			x
tea tree, New Zealand ( <i>Leptospermum scoparium</i> )	ruby glow	x	x
tree fern (tiki fern) ( <i>Asparagus virgatus</i> )			x
trumpet flower or Carolina Jessamine ( <i>Gelsemium sempervirens</i> )			x
tufted hairgrass ( <i>Deschampsia caespitosa</i> )			x

Name	Tolerant Cultivars	Application to Field / Landscape or Container Grown Ornamentals Noted by a (x)	
		Container Grown	Field / Landscape Grown
tulip ( <i>Tulip spp</i> )	apeldorn		x
verbena, shrub ( <i>Lantana sellowiana</i> )			x
Verbena, St. Paul's ( <i>Verbena peruviana</i> )	St. Paul		x
viburnum ( <i>Viburnum spp.</i> )	American cranberry bush arrowwood European cranberry bush linden Mohican wright		x x x x x
vinca (periwinkle) ( <i>Vinca minor</i> )			x
windmill palm ( <i>Trachycarpus fortunei</i> )			x
xylosma ( <i>Xylosma congestum</i> )			x
yarrow ( <i>Achillea spp.</i> )			x
yaupon ( <i>Ilex vomitoria</i> )	dwarf		x
yew ( <i>Taxus cuspidata</i> ) ( <i>Taxus x media</i> )	capitata denisiformis	x	x x

\*Ornamental species only. Do not apply this product directly to plants that are grown for food.

## STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

**Pesticide Storage:** Store this product only in its original container. In case of spill, sweep up material and dispose of material according to label directions.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

**EMPTY CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty bag into application equipment. Dispose of empty bag(s) in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## CONDITIONS OF SALE AND WARRANTY

J.R. Simplot Company warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. To the extent consistent with applicable law J.R. SIMPLOT COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE. Handling, storage and use of the product by Buyer or User are beyond the control of J.R. Simplot Company and Seller. To the extent consistent with applicable law risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pest, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. To the extent consistent with applicable law, in no case will J.R. Simplot Company or Seller be held liable for consequential, special or indirect damages resulting from the handling, storage or use of this product.

DIMENSION® is a registered trademark of Dow AgroSciences LLC.

EPA Approved 12/23/2010



Tech sheets, MSDS and other information on BEST products available at [www.bestfertilizer.com](http://www.bestfertilizer.com)

**NET WEIGHT: 50 LBS. (22.68 Kgs.)**

**Simplot**

Manufactured in U.S.A. for:  
J.R. SIMPLOT COMPANY  
P.O. Box 198, Lathrop, CA 95330