



Water-Soluble Foliar Fertilizer

Folo Spray 20-20-20 WITH MICRONUTRIENTS

GUARANTEED ANALYSIS:

TOTAL NITROGEN (N)	20.00%
2.40% Ammoniacal Nitrogen	
4.60% Nitrate Nitrogen	
13.00% Urea Nitrogen	
AVAILABLE PHOSPHATE (P ₂ O ₅)	20.00%
SOLUBLE POTASH (K ₂ O)	20.00%
Boron (B)	0.02%
Copper (Cu)	0.05%
0.05% Chelated Copper	
Iron (Fe)	0.10%
0.10% Chelated Iron	
Manganese (Mn)	0.05%
0.05% Chelated Manganese	
Molybdenum (Mo)	0.0005%
Zinc (Zn)	0.05%
0.05% Chelated Zinc	

Derived from Monoammonium Phosphate, Monopotassium Phosphate, Potassium Nitrate, Urea, Sodium Borate, Copper EDTA, Iron EDTA, Manganese EDTA, Zinc EDTA and Sodium Molybdate.

KEEP OUT OF REACH OF CHILDREN
WARNING
HARMFUL IF INHALED. CAUSES EYE IRRITATION.

PRECAUTIONARY STATEMENTS: Avoid breathing dust. Use only outdoors or in a well-ventilated area. Wash hands thoroughly after handling.

FIRST AID: IF INHALED Remove person to fresh air and keep comfortable for breathing. Call a poison control center (1-877-424-7452) or doctor for treatment advice if you feel unwell.

IF IN EYES Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists get medical advice / attention.

GENERAL USE DIRECTIONS

DILUTION RATE: Unless otherwise specified, use the rate per acre and dilute as follows:

Air: 5-30 gallons of water per acre.

Dilute: 100 plus gallons of water per acre.

Concentrate: 10-100 gallons of water per acre.

Do not exceed 2 lbs. of product per gallon of water.

Add approximately ½ water to tank before adding Folo Spray 20-20-20, agitate thoroughly while adding remaining water, add pesticide last.

When temperatures are cold allow extra time for this product to completely dissolve. Do not use when temperatures are expected to exceed 90°F or when plants are under stress conditions such as heat or water stress.

THIS PRODUCT WILL LOWER THE pH OF THE SPRAY SOLUTION.

CAUTION: This product contains Boron and Molybdenum. Use of this product at rates other than those recommended, may cause injury to susceptible plants.

DO NOT EXCEED RECOMMENDED AMOUNTS PER ACRE.

MIXING INSTRUCTIONS: Add to tank when half full, agitation is advised. Water temperature plays an important role in determining fertilizer solubility. The higher the water temperature the greater the solubility and quicker the dissolving time. As fertilizers are added to the solution, there is an immediate decrease in water temperature. It is important when preparing concentrated solutions to allow for this temperature drop to ensure that all the fertilizer is properly dissolved.

COMPATIBILITY: Folo Spray 20-20-20 is compatible with most commonly used pesticides and fungicides. However, a compatibility test is recommended before full scale mixing, if the desired combination has not been used in the past. Do not combine with copper fungicide or other strongly alkaline materials.

APPLES, APRICOTS, CHERRIES, CITRUS (ALL TYPES), NECTARINES, PEACHES, PEARS, PLUMS, PRUNES, ALMONDS, PECANS, PISTACHIOS, WALNUTS: As a foliar spray apply at the rate of 5 to 10 lbs. per acre. For dilute sprays use 2 lbs. per 100 gallons of water. Repeat as necessary. For best results, make 3 to 4 applications per season. **CAUTION: DO NOT USE Folo Spray 20-20-20 in late season sprays where fruit color and maturity is delayed by a nitrogen application.** Do not use on Golden Delicious during slow drying conditions.

STRAWBERRIES: Apply at the rate of 5 to 10 lbs. per acre when the first fruit buds show in the crown. For best results, apply 3 or more additional applications at 7 to 10 day intervals.

GRAPES, RASPBERRIES, BLACKBERRIES AND BLUEBERRIES: Apply at the rate of 5 to 10 lbs. per acre early in the season or whenever it is necessary to improve vine or cane vigor and growth. For best results, repeat at pre-bloom and 3 weeks later. Use sufficient water for good coverage.

CAUTION: DO NOT USE Folo Spray 20-20-20 where fruit color and maturity is delayed by a nitrogen application.

COTTON, SOYBEANS, PEANUTS, FIELD CORN, POPCORN, SUGAR BEETS, TABLE BEETS AND SWEET POTATOES: Apply at the rate of 5 to 10 lbs. per acre to improve plant vigor. For best results, make 1 to 5 applications depending on crop and season.

TOMATOES, PEPPERS, CUCUMBERS, MELONS AND SQUASH: Apply at the rate of 5 to 10 lbs. per acre. For best results, make first application when plants are 3 to 4 weeks old. Repeat at 7 to 10 day intervals with 5 to 7 applications per season.

POTATOES: Apply 5 to 10 lbs. per acre. Apply at initial tuber formation. For best results, apply when plants are young and continue at 7 to 10 day intervals through 30 day tuber setting period.

CELERY, LETTUCE, ENDIVE, BROCCOLI, CABBAGE, CAULIFLOWER, BRUSSEL SPROUTS, KALE AND SPINACH: Apply at 5 to 10 lbs. per acre when plants have been established for 3 weeks. For best results, use 3 to 6 sprays per season at 7 to 10 day intervals, depending on weather conditions and amount of growth desired.

CARROTS, PARSLEY, ASPARAGUS: Apply 10 to 15 lbs. per acre. Frequency of application depends on prevailing weather conditions and amount of growth desired.

BEANS, PEAS, SWEET CORN, LENTILS AND ONIONS: Apply at the rate of 5 to 10 lbs. per acre when plants are 3 to 4 weeks old. For best results, apply at 7 to 10 day intervals or as needed.

WHEAT, BARLEY, OATS, RYE, RICE, SUNFLOWER AND SUGAR CANE: Apply at the rate of 5 to 10 lbs. per acre throughout growing season. For best results, apply at tillering and again at early boot stage.

HOPS: Ground sprayers; apply 8 to 10 lbs. per acre in 100 to 200 gallons of water. Air application; apply 5 to 8 lbs. per acre in not less than 10 gallons of water. For best results on hops, apply when half way up the wire and again when they reach the top of the wire.

MINT: Apply 5 to 10 lbs. per acre when plants have 6-8" growth. For best results, make 1 to 5 applications during the growing season.

FORAGE AND HAY CROPS: ALFALFA, CLOVER AND GRASSES (such as orchard grass and Timothy): Apply at 5 to 10 lbs. per acre. For best results, apply at 7 to 10 day intervals depending on amount of growth desired.

SEED CROPS, (GRASS, ALFALFA, CLOVER, CARROTS, LETTUCE AND ASPARAGUS): Apply at 5 to 10 lbs. per acre. For best results, apply just prior to head formation and then at 7 to 10 day intervals.

COMMERCIAL NURSERY AND GREENHOUSE: As a direct foliar spray use 2 to 4 lbs. per 100 gallons or 4 tsp. per gallon of water. Greenhouse Cut Flowers, Flats, Small Pots: Use 2 lbs. per 100 gallons of water for intermittent feeding (every 2 weeks); for constant feeding (every watering) use 4 to 8 oz. per 100 gallons of water. Container Foliage Plants: Use up to 3½ lbs. per 100 gallons of water for intermittent feeding. For constant feeding use 8 to 12 oz. per 100 gallons of water. Orchids: For periodic feeding use 5 to 6 lbs. in 100 gallons of water; for constant feeding use 1 to 1½ lbs. per 100 gallons of water. Starter and Transplant: Dissolve 2 lbs. in 100 gallons of water using either as a foliar spray, or using 4 to 8 oz. of solution per plant. To prepare a slurry add ¼ lb. to a 3 gallon bucket of puddled soil.

Warning: This product contains a chemical known to the State of California to cause cancer, birth defects or other reproductive harm. Proposition 65, the Safe Drinking Water and Toxic Enforcement Act of 1986, requires notification of potential exposure to substances identified by the State of California as causing cancer, birth defects or other reproductive harm.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-jr.com>

DISCLAIMER OF WARRANTIES: Seller warrants that the chemical composition of this product conforms to the chemical description given on this label. THIS WARRANTY IS EXPRESSLY IN LIEU OF ALL OTHER WARRANTIES AND REPRESENTATIONS EXPRESSED, IMPLIED, OR STATUTORY, INCLUDING WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR USE. Timing, rate and method of application, weather and crop conditions, mixtures with chemicals not specifically recommended on this label or an accompanying written recommendation, are beyond the control of seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. Buyer further agrees in the event of damages arising from the use of this product to accept a replacement of the product or a refund of the purchase price of the product, at buyer's option, as full discharge of seller's liability. No one is authorized to make any other warranty, guarantee or directions concerning this product, and no such warranties, guarantees or directions shall be valid or binding upon seller. GHS_R2-15-15

NET WEIGHT: 25 POUNDS (11.34 Kg.)



J. R. SIMPLOT COMPANY

P.O. Box 70013, Boise ID 83707-0113 • (208) 336-2110

GUARANTOR

P.O. Box 198, Lathrop, CA 95330-0198