

TRACEMAKER BLEND

9% ZINC A Citrate / EDTA Chelate

PRECAUTIONS IN USING

Follow directions for foliar and soil applications as outlined on this label. Over-application of TRACEMAKER BLEND 9% ZINC may be phytotoxic or cause undesirable effects to foliage.

GENERAL INFORMATION

TRACEMAKER BLEND 9% ZINC is a liquid source of the micronutrient in a form available to plants. It may be mixed and applied with liquid fertilizers or water. It may be used as a foliar or soil applied spray.

Addition of this product to fertilizer solutions, especially concentrates, should not be done without first testing for compatibility on a small scale.

Foliar application with water should thoroughly cover all foliage - see directions for use. For best results apply in early morning or late evening. For aerial applications, use at least one gallon of water for each pint of this product applied to the crop.

TRACEMAKER BLEND 9% ZINC applied to the soil must be moved into the root zone to be absorbed by the plant. This may be placed in the root zone or moved into the zone by rainfall, irrigation or tillage.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

STORAGE: STORE IN COOL PLACE OUT OF DIRECT SUNLIGHT. MAY BE STORED IN UNHEATED FACILITY. SPILLS FROM BROKEN CONTAINERS CAN BE CLEANED UP BY ABSORBING ON CLAY OR OTHER SUITABLE ABSORBENT.

DISPOSAL: WASTES RESULTING FROM THE USE OF THIS PRODUCT MAY BE DISPOSED OF AT AN APPROVED WASTE DISPOSAL FACILITY.

CONTAINERS: TRIPLE RINSE (OR EQUIVALENT). THEN OFFER FOR RECYCLING OR RECONDITIONING. OR PUNCTURE AND DISPOSE OF IN A SANITARY LANDFILL OR BY OTHER PROCEDURES APPROVED BY STATE AND LOCAL AUTHORITIES.

NOTICE

Agri Trend, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Agri Trend, Inc. In no case shall Agri Trend, Inc. be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Agri Trend, Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.



AGRI TREND, INC.

P. O. Box 25597 • Yuma, Arizona 85367-0597

(520) 342-5852 • Fax: (520) 342-9346

AT-0071

GUARANTEED ANALYSIS 5.8-0-0 BY WEIGHT

Total Nitrogen	5.8%
5.8% Other Water Soluble Nitrogen	
Zinc (Zn)	9.0%
9.0% Chelated Zinc	
Derived from Zinc Sulfate, Citric Acid and Zinc EDTA (Ethylenediaminetetraacetate acid).	

NOTE: When mixing liquids with other chemicals, all chemical labels should be read in their entirety to ensure both chemical compatibility, and compatibility based on the pH.

CAUTION: KEEP OUT OF REACH OF CHILDREN

- Net Contents 1 Gallon (3.785 L); 10.5 pounds (4.767 Kg)
- Net Contents 5 Gallons (18.925 L); 52.5 pounds (23.835 Kg)

CONTAINS AMMONIA.
CAN CAUSE BURNS.
VAPOR HARMFUL.

HARMFUL IF SWALLOWED.

Avoid contact with eyes and clothing. Avoid prolonged contact with skin. Do not inhale vapors. Do not take internally.

FIRST AID

If inhaled: Remove to fresh air. If not breathing, give artificial respiration. If breathing is difficult, give oxygen. Call a physician. **If swallowed:** Drink plenty of water if conscious. Call a physician. **In case of contact:** Immediately flush eyes or skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. Call a physician. Wash clothing before reuse. Thoroughly clean shoes before reuse.

CONTENTS MAY BE UNDER PRESSURE
USE CAUTION WHEN OPENING.
WEAR GOGGLES AND GLOVES
WHEN HANDLING.

LOT NUMBER

For the Prevention and Correction
of Zinc Deficiencies in Crops

DIRECTIONS FOR USE APPLICATION RECOMMENDATIONS

FRUIT AND NUT CROPS

APPLES, CHERRIES, CITRUS, GRAPES, PEACHES, PEARS AND PLUMS: **Foliar:** 3/4-1 1/2 qts. in 100 gals. of water 3-4 weeks after full bloom. For severe deficiencies make 1 or 2 additional applications at 2-3 week intervals. **Soil:** 1-2 qts. per tree with liquid fertilizer or water in April or May.

ALMONDS, PECANS AND WALNUTS: **Foliar:** 1 qt. in 100 gals. of water at full leaf stage in the spring. Larger or older trees may receive repeat applications at 3 week intervals. **Soil:** 1 qt. per tree with liquid fertilizer or water in the spring.

VEGETABLES

PEPPERS, TOMATOES: **Foliar:** 1-2 qts. per acre in sufficient water to thoroughly cover foliage. Early and mid season. **Soil:** Band or sidedress 1-2 qts. per acre at planting or within a few weeks after planting.

BEANS, BEETS, BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CARROTS, CAULIFLOWER, CELERY, CUCUMBER, EGGPLANT, LETTUCE, MELONS, ONIONS, PEAS, POTATOES: **Foliar:** 1-1 1/2 qts. per acre in sufficient water to thoroughly cover foliage. Apply early to mid season. **Soil:** 2-4 qts. per acre in a band with liquid fertilizer or water at planting or 1-2 weeks after planting.

FIELD CROPS

ALFALFA: **Foliar:** 1-1 1/2 qts. per acre in sufficient water to thoroughly wet foliage when new growth is 4" to 5" high. **Soil:** Broadcast 2-4 qts. per acre with liquid fertilizer or water prior to planting.

MINT: **Foliar:** 1-1 1/2 qts. per acre in sufficient water to thoroughly wet foliage when new growth is 3" to 4" high. **Soil:** Broadcast 2-4 qts. per acre with liquid fertilizer or water prior to planting.

CANOLA, FLAX, CORN (FIELD AND SWEET), DRY BEANS, FIELD PEAS, HOPS, SAFFLOWER, SORGHUM, SOYBEANS, SUN-FLOWERS, WHEAT: **Foliar:** 1-1 1/2 qts. per acre in sufficient water to thoroughly cover foliage 4 weeks after planting. If necessary, repeat application in 3 weeks. **Soil:** Broadcast 2-4 qts. per acre with liquid fertilizer or water prior to or at planting.

COTTON, PEANUTS: **Foliar:** 1-1 1/2 qts. per acre in sufficient water when plant is over 6 inches tall. **Soil:** Incorporate 2-4 1/2 qts. per acre with liquid fertilizer or water prior to planting.

SUGAR BEETS: **Foliar:** 2-3 qts. per acre in sufficient water to thoroughly wet foliage after thinning. **Soil:** 2-4 qts. per acre in a band with liquid fertilizer or water prior to planting.

ORNAMENTALS

TREES AND SHRUBS: **Foliar:** 3/4-1 1/2 qts. in 100 gals. of water at full feed stage in the spring. Repeat applications at 3-4 week intervals if necessary. **Soil:** Spray 1 pt. per 1000 sq. ft. in the spring and water into the soil by sprinkler irrigation.

FLOWERING PLANTS, ORNAMENTAL PLANTS, ANNUALS AND OTHER ORNAMENTALS: **Foliar:** 2-3 fl. oz. per 1000 sq. ft. in sufficient water to thoroughly cover foliage and soil surface around plants. **Soil:** 3-4 fl. oz. per 1000 sq. ft. before planting or at planting. Work lightly into soil or water in with irrigation water.

Yuma Printing Co. 11/98