

NUTRI-AID

20-20-20

Water Soluble Fertilizer With Micronutrients



GUARANTEED ANALYSIS

Total Nitrogen (N).....	20.0%
3.9% Ammoniacal Nitrogen	
5.7% Nitrate Nitrogen	
10.4% Urea Nitrogen	
Available Phosphate (P ₂ O ₅).....	20.0%
Soluble Potash (K ₂ O).....	20.0%
Boron (B).....	0.02%
Copper (Cu).....	0.05%
Iron (Fe).....	0.10%
Manganese (Mn).....	0.05%
Zinc (Zn).....	0.05%

Derived from ammonium phosphate, urea, potassium nitrate, sodium borate, copper sulfate, iron sulfate, manganese sulfate and zinc sulfate.

**CAUTION: KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED—AVOID BREATHING DUST**

DIRECTIONS: 5 to 10 lbs. per 40-100 gallons of water.

Aerial Application: 3-10 lbs. per acre in sufficient water for good coverage.

Fill tank with water and slowly add **NUTRI-AID** while agitating. Continue to agitate until all **NUTRI-AID** is completely dissolved.

APPLICATION IN IRRIGATION WATER

Solid Set, Overhead, Drip Sprinkler or Pivotal Irrigation Systems: Dissolve **NUTRI-AID** in mix tank and apply 8-10 lbs. per acre in irrigation water. Use of anti-siphon or check valve is recommended to prevent solution from running back into primary water supply.

NUTRI-AID is generally compatible with pesticides and other fertilizers. Determine compatibility with a jar test before mixing products.

ASPARAGUS: Uses 5 to 10 lbs. per application. Make subsequent applications depending on weather and desired growth.

AVOCADOS: Use 10 to 20 lbs. per application in sufficient water to thoroughly wet foliage. Do not spray to run off.

BEANS, PEAS, LENTILS, RAPESEED (Canola): Use 5 to 10 lbs. per acre. Apply at first sign of bloom. Repeat 2 weeks later.

BERRIES: (Strawberries, raspberries, blackberries, blueberries.) Use 5 to 10 lbs. per acre in spring on new growth. Repeat at pre-bloom and 3 weeks later. Use sufficient water for good coverage.

CARROTS, BEETS, SWEET POTATOES, AND OTHER VEGETABLE ROOT CROPS: Use 5 to 10 lbs. per acre per application.

CITRUS INCLUDING ORANGE, LEMON, GRAPEFRUIT, TANGERINE: Use 10 to 20 lbs. per acre per application. In concentrate spray, use 10 to 12 lbs. per acre per application in 20 to 100 gallons of water. In dilute sprays, use 3 to 4 lbs. per 100 gallons of water and wet foliage completely.

COLE CROPS, LETTUCE, LEAFY VEGETABLES: Use 5 to 10 lbs. per acre. Apply right after thinning or transplanting. Repeat every 10 to 14 days.

CORN: (Field and Sweet): Use 5 to 10 lbs. per acre. Apply when corn is 6"-8" high and again prior to tasseling.

COTTON: Use as a broadcast spray of 5 to 10 lbs. per acre in sufficient water to evenly distribute material. Apply as a foliar spray 1 to 5 times per season for general maintenance.

CRANBERRIES: Use 5 to 10 lbs. per acre. Apply at popcorn stage. Repeat at hooking and 3 weeks later.

DECIDUOUS FRUITS (Apples, pears, cherries, plums, prunes, peaches, almonds, walnuts, filberts, pistachios and pecans): Apply in delayed dormant, pre-bloom, and post-bloom sprays at 3 to 5 lbs. per 100 gal. in dilute sprays. Apply 10 to 15 lbs. per acre in concentrate sprays. Avoid combining with material that will cause russetting.

FIELD CROPS (Mint, alfalfa, clover): Use 5 to 10 lbs. per acre. Apply when plants have 6"—8" new growth. Repeat application to new growth after each cutting.

GRAPES: Ground sprayers: Use 5 to 10 lbs. per acre in 50 to 100 gal. of water. Air Application: Use 5 to 8 lbs. per acre in 10 gal. of water. Apply when leaves emerge, pre-bloom and post-bloom until grapes are 1/4 size.

HOPS: Ground sprayers: Use 5 to 10 lbs. per acre in 100-200 gallons of water. Air Application: Use 5 to 8 lbs. per acre in no less than 10 gal. of water. Apply when hops are halfway up wire and again when they reach the top of the wire.

MELONS, CUCUMBERS AND OTHER CUCURBITS: Use 5 to 10 lbs. per acre per application. Apply two or more applications every 2 to 3 weeks beginning with pre-bloom.

ONIONS, GARLIC, TOMATOES, PEPPERS: Use 5 to 10 lbs. per acre per application. Begin application prior to bloom. Repeat every 2-3 weeks thereafter.

POTATOES: Use 5 to 10 lbs. per acre. Apply when plants are 4"—6" high (hooking stage). Subsequent applications should be at 2 to 3 week intervals.

SAFFLOWER, SUNFLOWER, AND OTHER OILSEED CROPS: Use 5 to 10 lbs. per acre per application.

SMALL GRAINS (Wheat, oats, barley, rye): Use 5 to 10 lbs. per acre in sufficient water to cover by air or ground. Apply at tillering and again at early boot stage.

SUGAR BEETS: Use 5 to 10 lbs. per acre. Apply when leaves are 10" across. Repeat application 2 to 4 weeks later.

CONDITIONS OF SALE AND LIMITED WARRANTY:

The directions for use of this product reflect the opinions of experts based on field uses and laboratory testing. Always follow label instructions. Manufacturer warrants that this product conforms to the product description on this label. Manufacturer or seller makes no other expressed or implied warranty. Neither manufacturer nor seller shall be liable for any injury or any application not specifically described on the label.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

MANUFACTURED BY: JH BIOTECH, INC. P.O. Box 3538 Ventura, CA 93006

NET WEIGHT: 25 LBS (11.25 Kg.) 50 LBS (22.7 Kg.)

Lot No.:

JH Biotech Inc.

0809-D-G-2
Copyright © JH Biotech, Inc.
All rights reserved.