

11.05 LBS. PER GALLON @ 68°F  
1235 GRAMS PER LITER

### APPLICATION RATES

Use a total of 2-3 gallons per season if needed:

**ALFALFA CLOVER** - 1 to 2 qts. per acre per application. For hay, apply at 6 to 8 inches of new growth and after each cutting. For seed, apply at bud stage and repeat at intervals of 7 to 10 days.

**APPLES, APRICOTS, CHERRIES, NECTARINES, PEARS, PLUMS, AND PRUNES** - 1 to 4 qts. per acre per application. Apply pre and /or post-bloom and repeat at intervals of 7 to 10 days.

**HOPS** - 1 to 2 qts. per acre.

**NUT CROPS (ALMONDS, FILBERTS, PECANS, PISTACHIOS, AND WALNUTS)** - 1 to 4 qts. per acre per application. Repeat every 7 to 10 days.

**ARTICHOKES** - 1 to 2 qts. per acre per application. Apply bud stage, repeat every 7 to 10 days.

**ASPARAGUS** - 1 to 2 qts. per acre per application. Apply just after start of new growth. Repeat every 7 to 10 days.

**AVOCADOS** - 1 to 2 qts. per acre per application.

**BEANS, SOYBEANS, PEAS, LENTILS,** - 1 to 2 qts. per acre. First application at bud stage. Repeat every 7 to 10 days.

**BLUEBERRIES, RASPBERRIES** - 1 to 2 qts. per acre pre bloom. Repeat every 7 to 10 days.

**CARROTS, BEETS, SWEET POTATOES AND OTHER VEGETABLE ROOT CROPS** - 1 to 3 qts. per acre when plants are three to four weeks old. Repeat every 7 to 10 days.

**CITRUS** - 2 to 4 qts. per acre per application pre-bloom. Then every 7 to 10 days.

**CORN** - 1 to 4 qts. per acre.


**COTTON** - 1 to 2 qts. per acre

**CRANBERRIES** - 1 to 2 qts. per acre. First application at hook stage, second application after fruit set.

**FIGS** - 1 to 3 qts. per acre

**GRAPES** - 1 to 4 qts. per acre. First application at the 4 to 6 leaf stage, repeat every 7 to 10 days as needed. Use enough dilution for full coverage.

Nutraplex® is a Registered Trademark of Western Nutrients Corporation.



# NUTRAPLEX® CAL MAG 5-0-0 Plus Boron

## GUARANTEED ANALYSIS

Total Nitrogen (N).....	5.6%
5.6% Nitrate Nitrogen	
Calcium (Ca).....	6.5%
Magnesium (Mg) .....	0.6%
Boron (B) .....	0.5%
Copper (Cu) .....	.05%
.05% Chelated Copper	
Iron (Fe) .....	.10%
.10% Chelated Iron	
Manganese (Mn) .....	.10%
.10% Chelated Manganese	
Zinc (Zn) .....	.20%
.20% Chelated Zinc	

Derived from Calcium Nitrate, Boric Acid, Magnesium Nitrate, Copper, Iron, Manganese, and Zinc Edta.  
ALSO CONTAINS NON - PLANT FOOD INGREDIENTS

### KEEP OUT OF REACH OF CHILDREN

WARRANTY: Western Nutrients Corp. makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on the Western Nutrients Corp. Label attached to the product container.

**WARNING: EXCESS AMOUNTS OF BORON (B) CAN CAUSE PLANT INJURY.**

MANUFACTURED BY  
WESTERN NUTRIENTS CORPORATION

245 Industrial Street, Bakersfield, Ca. 93307  
Phone (661) 327-9604/Fax (661) 327-1740  
E-mail: westernnutrients@lightspeed.net  
Website: <http://www.westernnutrients.net>

NET CONTENTS 5 GALLONS  
18.93 LITERS

### APPLICATION RATES

**LETTUCE, CELERY, AND LEAFY VEGETABLE CROPS** - 1 to 2 qts. per acre. Repeat every 15 days.  
**GRAPES** - 2 - 6 quarts per acre five times each season starting at first full leaf, then past bloom, early bunch set, berry size and at berry color and just before harvest.

**MELONS AND CUCUMBERS** - 1 to 2 qts. per acre. Make applications at intervals of 7 to 10 days starting just prior to bloom.

**MINT** - 1 to 2 qts. per acre.

**OLIVES** - 1 TO 2 qts. per acre.

**ONIONS AND GARLIC** - 1 to 2 qts. per acre.

**PEANUTS** - 1 to 2 qts. per acre. First application at early bloom stage, repeat at intervals of 7 to 10 days.

**POTATOES** - 1 to 2 qts. per acre per application. First application when plants are 4 to 12 inches high. Second application 3 weeks later, repeat every 10 days.

**SAFFLOWER AND SUNFLOWER** - 1 to 2 qts. per acre per application.

**STRAWBERRIES** - 1 to 2 qts. per acre per application.

**SUGAR BEETS** - 1 to 3 qts. per application. First application when plants are 6 to 8 inches across. Second application 3 to 4 weeks later, repeat as needed.

**TOMATOES AND PEPPERS** - 1 to 2 qts. per application.

**VEGETABLES** - 1 to 3 qts. per acre. First application at bud stage, repeat applications at intervals of 7 to 10 days.

**WHEAT, BARLEY, RICE, AND OTHER SMALL GRAINS** - 1 to 2 qts. per acre per application. First application at boot stage, repeat 7 to 10 days.

**SOIL** - 2 to 5 gallons per acre on all crops.

**DILUTION RATE:** Use sufficient water for thorough coverage. Maximum dilution 1 qt. per 10 gallons of water.

**GENERAL USE** - Nutraplex is compatible with most herbicide, pesticides, fungicides, and UAN solution. Use ample water for coverage. Can be used with all types of application equipment.