



GROUP 2 9 HERBICIDES

For control of weeds on specified noncrop use sites and forestry sites.

ACTIVE INGREDIENT:

Isopropylamine salt of imazapyr (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid)*	8.36%
Isopropylamine salt of glyphosate (N-(phosphonomethyl)glycine)	22.13%
OTHER INGREDIENTS:	69.51%
TOTAL	100.00%

* Equivalent to 6.82% (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid) or 0.637 pounds acid per gallon and 16.40% N-(phosphonomethyl)glycine acid or 1.531 pounds per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See next page for additional precautionary statements.

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or a doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going to treatment.	
FOR MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-866-944-8565.	

EPA REG. NO. 34704-989

EPA EST. NO. 34704-MS-001

NET CONTENTS 2½ GALS. (9.46 L)

042412 V1D 04G12

**Prep-It® HERBICIDE
EPA REG. NO. 34704-989**

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with eyes, skin, or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are made from any waterproof material. If you want more options, follow the instructions for **Category A** on an EPA chemical-resistant category selection chart.

Mixers, loaders, applicators and other handlers must wear:

- Long-sleeve shirt and long pants,
- Shoes plus socks,
- Chemical resistant gloves for all mixers and loaders, plus applicators using handheld equipment.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

ENGINEERING CONTROLS

Pilots must use an enclosed cockpit that meet the requirements listed in the Worker Protection Standards for agricultural pesticides in 40CFR 170.240(d)(6).

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands with plenty of soap and water before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to plants. Drift and run-off may be hazardous to plants in water adjacent to treated areas. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. See Directions for Use for additional precautions and requirements.

The use of treated waters on irrigated crops within 120 days of treatment is prohibited.

PHYSICAL AND CHEMICAL HAZARDS

Spray solutions of this product should be mixed, stored, and applied only in stainless steel, fiberglass, plastic, and plastic-lined steel containers.

DO NOT mix, store, or apply this product or spray solutions of this product in unlined steel (except stainless steel) containers or spray tanks.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not enter or allow others to enter treated areas until sprays have dried. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Prep-It® Herbicide must be used only in accordance with directions on the label. Keep containers closed to avoid spills and contamination.

Prep-It® HERBICIDE
EPA REG. NO. 34704-989

Prep-It Herbicide may be applied using helicopters, ground operated sprayers, and/or low-volume hand operated spray equipment such as backpack and pump-up sprayers.

Observe all cautions and limitations in the package labels of products used in combination with Prep-It Herbicide.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

The requirements in this box apply to use on trees being grown for sale or other commercial use, or for commercial seed production, or for production of timber or wood products, or for research purposes.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls,
- Shoes plus socks,
- Chemical-resistant gloves made of any waterproof material,
- Protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

DO NOT enter or allow others to enter treated areas until sprays have dried.

RESTRICTIONS

DO NOT use on food or feed crops. **DO NOT** use on Christmas trees. **DO NOT** treat irrigation ditches, or water used for crop irrigation or for domestic uses. Keep from contact with fertilizers, insecticides, fungicides, and seeds to prevent unintentional exposure of desirable vegetation to Prep-It Herbicide. **DO NOT** apply or drain or flush equipment on or near sensitive desirable plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. **DO NOT** side trim desirable vegetation with this product. Do not allow sprays to drift onto desirable plants.

Clean application equipment after using this product by thoroughly flushing with water.

PRODUCT INFORMATION

Prep-It Herbicide is an aqueous solution containing surfactant. It is mixed in water and generally applied as a postemergence spray for control of most annual and perennial grasses, broadleaf weeds, vines, and brambles and hardwood brush and trees for forestry site preparation.

Prep-It Herbicide may be applied on forestry sites that contain areas of temporary surface water caused by collection of water between planting beds, in equipment ruts, or in other depressions created by forestry management activities, except in the states of California and New York. It is permissible to treat drainage ditches, intermittent drainage, intermittently flooded low-lying sites, seasonally dry flood plains, and transitional areas between upland and lowland sites when no water is present, except in the states of California and New York. Only the edge of drainage ditches can be treated for drainage ditches that contain water. It is also permissible to treat marshes, swamps, and bogs after water has receded, as well as seasonally dry flood deltas, except in the states of California and New York. **DO NOT** make applications to natural or man-made bodies of water such as lakes, reservoirs, ponds, streams, rivers, and canals.

Prep-It® HERBICIDE
EPA REG. NO. 34704-989

Prep-It Herbicide is also recommended for control of undesirable vegetation along forest roads, non-irrigation ditchbanks, and the establishment and maintenance of wildlife openings except in the state of California. See use directions for **DIRECTED FOLIAR OR SPOT SPRAYS AND SITE PREPARATION TREATMENTS**.

SYMPTOMOLOGY:

Prep-It Herbicide is readily absorbed through the foliage and the roots and is translocated rapidly throughout the plant, with accumulation in the meristematic regions. Treated plants stop growing soon after spray application. Chlorosis first appears in the youngest leaf tissue. In perennials, the herbicide is translocated into the roots, thus preventing most resprouting. The foliage of most woody plants, brush and trees will normally display color change and necrosis within several weeks after application.

MIXING AND APPLICATION INSTRUCTIONS

WIND EROSION

Avoid treating powdery dry or light sandy soils when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation.

BROADCAST APPLICATIONS

Aerial Application Restrictions (Helicopter Only): Applicators must follow these requirements to avoid off-target drift movement:

1. Applicators are required to use a coarse or coarser droplet size (ASABE S572) or, if specifically using a spinning atomizer nozzle, applicators are required to use a volume mean diameter (VMD) of 385 microns or greater for release heights below 10 feet. Applicators are required to use a very coarse or coarser droplet size or, if specifically using a spinning atomizer nozzle, applicators are required to use a VMD of 475 microns or greater for release heights above 10 feet. Applicators must consider the effects of nozzle orientation and flight speed when determining droplet size.
2. Applicators are required to use upwind swath displacement.
3. The boom length must not exceed 90% of the rotor blade diameter to reduce spray drift.
4. Applications with wind speeds less than 3 mph and with wind speeds greater than 10 mph are prohibited.
5. Applications into temperature inversions are prohibited.

Applicators must follow the most restrictive use cautions to avoid drift hazards, including those found in this labeling as well as applicable state and local regulations and ordinances.

Helicopter Spray Equipment

Thoroughly mix the specified amount of Prep-It Herbicide in 5.0 to 30.0 gallons of water per acre and uniformly apply with properly calibrated aerial equipment. All precautions must be taken to minimize or eliminate spray drift. Applications must not be made under windy or gusty conditions. Use controlled droplet booms and nozzles configuration. A drift control agent may be added at the recommended label rate (Compadre® 1.0 pint per 100 gallons of water) except when applying with a **Microfoil boom, Thru-Valve Boom®** or other similar equipment. A foam reducing agent may also be added at the specified label rate, if needed.

IMPORTANT: DO NOT make applications by fixed wing aircraft. Maintain adequate buffer zones. Thoroughly clean application and mixing equipment, including landing gear, immediately after use. Prolonged exposure of this product to uncoated steel (except stainless steel) surfaces may result in corrosion and failure of the exposed part.

Ground Boom Application Restrictions (Broadcast): Use 5.0 or more gallons of water per acre. The actual minimum spray volume per acre is determined by the spray equipment used. Applicators must follow these requirements to avoid off-target drift movement:

1. Applicators are required to use a nozzle height below 4 feet above the ground or plant canopy and coarse or coarser droplet size (ASABE S572) or, if specifically using a spinning atomizer nozzle, applicators are required to use a volume mean diameter (VMD) of 385 microns or greater.
2. Applications with wind speeds greater than 10 mph are prohibited.
3. Applications into temperature inversions are prohibited.

Prep-It® HERBICIDE
EPA REG. NO. 34704-989

Ground Operated Spray Equipment

Thoroughly mix and apply the specified amount of Prep-It Herbicide in 5.0 to 60.0 gallons of water per acre. A drift control agent and a foam reducing agent (Compadre 1.0 pint per 100 gallons of water) may be added at the specified label rates. If desired, a spray pattern indicator may be added at the specified label rate. For best results, uniformly cover the foliage of the vegetation to be controlled with the spray solution.

IMPORTANT: DO NOT spray under windy or gusty conditions. Maintain adequate buffer zones. Clean application and mixing equipment after use by thoroughly flushing it with water.

DIRECTED FOLIAR OR SPOT SPRAY EQUIPMENT

In addition to broadcasting the mixed herbicide solution across an entire acre, the solution can also be directed to individual clumps of herbaceous and woody weeds or to spots within the acre. Backpack sprayers and ground-operated equipment with hoses are generally used for directed foliar or spot sprays. Specialized helicopter equipment can also be used for this purpose. When making directed or spot sprays application with backpack sprayers, ground operated equipment, helicopters, or similar equipment that permits directed application, thoroughly mix a solution of 5 to 10 percent by volume of Prep-It Herbicide.

To mix the spray solutions add the volume of Prep-It Herbicide indicated in the table below to the desired amount of water.

SPRAY SOLUTION MIXING GUIDE

Solution Volume	Prep-It Herbicide	
	Percentage of Total Solution Volume	
	5	10
1.0 gal	6.4 fl ozs	12.8 fl ozs
5.0 gals	2.0 pts	4.0 pts
10.0 gals	4.0 pts	8.0 pts
25.0 gals	10.0 pts	20.0 pts
100 gals	5.0 gals	10.0 gals

For best results, uniformly cover the foliage of the vegetation to be controlled with the spray solution.

IMPORTANT: DO NOT apply causing runoff from the treated foliage. Avoid direct application and drift to the foliage, thin bark and rooting zone of desired plant species as injury may occur. Even though the herbicide is directed to clumps and spots within an acre, and not broadcast.

DO NOT exceed 2.0 gallons of Prep-It Herbicide per acre.

SITE PREPARATION TREATMENTS

Prep-It Herbicide may be used at a rate of 1 gallon of product per acre to control labeled grasses, broadleaf weeds, vines and brambles, and woody brush and trees on forest sites in advance of regeneration for the following conifer crop species:

- | | |
|--|---|
| Loblolly Pine (<i>Pinus taeda</i>) | Shortleaf Pine (<i>Pinus echinata</i>) |
| Loblolly x Pitch Hybrid | Virginia Pine (<i>Pinus virginiana</i>) |
| Longleaf Pine (<i>Pinus palustris</i>) | Slash Pine (<i>Pinus elliotii</i>) |

Within 4 to 6 weeks of treatment, herbaceous weeds including grasses and woody vegetation will be controlled and may provide fuel to facilitate a site preparation burn, if desired, to enhance control of conifers or other species tolerant to the herbicide.

Apply 1.0 gallon of Prep-It Herbicide per acre in 5.0 to 30.0 gallons of total spray solution for helicopter applications or 5.0 to 60.0 gallons total spray solution for mechanical ground spray and backpack applications. Use higher spray volumes when controlling particularly dense or multi-layered canopies of hardwood stands, or difficult to control species.

Prep-It® HERBICIDE
EPA REG. NO. 34704-989

WEEDS CONTROLLED

When applied as directed Prep-It Herbicide will control, partially control, or suppress most woody brush, trees and herbaceous weeds, some of which are listed below. Degree of control and residual efficacy is both species and rate dependent. Prep-It Herbicide must be used only in accordance with the directions on this label.

GRASSES

The species of annual and perennial grasses controlled by Prep-It Herbicide include the following:

Annual Bluegrass	<i>Poa annua</i>	Orchardgrass	<i>Dactylis glomerata</i>
Bahiagrass	<i>Paspalum notatum</i>	Panicum	<i>Panicum</i> spp.
Barnyardgrass	<i>Echinochloa crus-galli</i>	Paragrass	<i>Brachiaria mutica</i>
Beardgrass	<i>Andropogon</i> spp.	Phragmites	<i>Phragmites australis</i>
Bermudagrass	<i>Cynodon dactylon</i>	Prairie cordgrass	<i>Spartina pectinata</i>
Big bluestem	<i>Andropogon gerardii</i>	Prairie threeawn	<i>Aristida oligantha</i>
Broadleaf signalgrass	<i>Brachiaria platyphylla</i>	Quackgrass	<i>Agropyron repens</i>
Canada bluegrass	<i>Poa compressa</i>	Reed canarygrass	<i>Phalaris arundinacea</i>
Cattail	<i>Typha</i> spp.	Saltgrass	<i>Distichlis s</i>
Cheat	<i>Bromus secalinus</i>	Sand dropseed	<i>Sporobolus cryptandrus</i>
Cogongrass	<i>Imperata cylindrica</i>	Sandbur	<i>Cenchrus</i> spp.
Crabgrass	<i>Digitaria</i> spp.	Smooth brome	<i>Bromus inermis</i>
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	Sprangletop	<i>Leptochloa</i> spp.
Dallisgrass	<i>Paspalum dilatatum</i>	Timothy	<i>Phleum pratense</i>
Downy brome	<i>Bromus tectorum</i>	Torpedograss	<i>Panicum repens</i>
Fall panicum	<i>Panicum dichotomiflorum</i>	Vaseygrass	<i>Paspalum urvillei</i>
Feathertop	<i>Pennisetum villosum</i>	Wild barley	<i>Hordeum</i> spp.
Fescue	<i>Festuca</i> spp.	Wild oats	<i>Avena fatua</i>
Foxtail	<i>Setaria</i> spp.	Wirestem muhly	<i>Muhlenbergia frondosa</i>
Giant reed	<i>Arundo donax</i>	Witchgrass	<i>Panicum capillare</i>
Goosegrass	<i>Eleusine indica</i>	Woolly cupgrass	<i>Eriochloa villosa</i>
Guineagrass	<i>Panicum maximum</i>		
Italian ryegrass	<i>Lolium multiflorum</i>		
Itchgrass	<i>Rottboellia exaltata</i>		
Johnsongrass	<i>Sorghum halepense</i>		
Junglerice	<i>Echinochloa colona</i>		
Kentucky bluegrass	<i>Poa pratensis</i>		
Lovegrass	<i>Eragrostis</i> spp.		

Prep-It® HERBICIDE
EPA REG. NO. 34704-989

BROADLEAF WEEDS

The species of annual and perennial broadleaf weeds controlled by Prep-It Herbicide include the following:

Arrowwood	<i>Pluchea sericea</i>	Mullein	<i>Verbascum</i> spp.
Broom snakeweed	<i>Gutierrezia sarothrae</i>	Nettleleaf goosefoot	<i>Chenopodium murale</i>
Bull thistle	<i>Cirsium vulgare</i>	Oxeye daisy	<i>Chrysanthemum</i> <i>leucanthemum</i>
Burclover	<i>Medicago</i> spp.		<i>Lepidium</i> spp.
Burdock	<i>Arctium</i> spp.	Pepperweed	<i>Amaranthus</i> spp.
Camphorweed	<i>Heterotheca subaxillaris</i>	Pigweed	<i>Plantago</i> spp.
Canada thistle	<i>Cirsium arvense</i>	Plantain	<i>Phytolacca americana</i>
Carolina geranium	<i>Geranium carolinianum</i>	Pokeweed	<i>Oenothera kunthiana</i>
Carpetweed	<i>Mollugo verticillata</i>	Primrose	<i>Tribulus terrestris</i>
Chickweed, mouseear	<i>Cerastium vulgatum</i>	Puncturevine	<i>Lythrum salicaria</i>
Clover	<i>Trifolium</i> spp.	Purple loosestrife	<i>Portulaca</i> spp.
Cocklebur	<i>Xanthium strumarium</i>	Purslane	<i>Richardia scabra</i>
Common chickweed	<i>Stellaria media</i>	Pusley, Florida	<i>Sisymbrium irio</i>
Common ragweed	<i>Ambrosia artemisiifolia</i>	Rocket, London	<i>Chondrilla juncea</i>
Cudweed	<i>Gnaphalium</i> spp.	Rush skeletonweed	<i>Centaurea repens</i>
Dandelion	<i>Taraxacum officinale</i>	Russian knapweed	<i>Salsola kali</i>
Desert camelthorn	<i>Alhagi pseudalhagi</i>	Russian thistle	<i>Atriplex</i> spp.
Diffuse knapweed	<i>Centaurea diffusa</i>	Saltbush	<i>Capsella bursa-pastoris</i>
Dock	<i>Rumex</i> spp.	Shepherdspurse	<i>Solanum elaeagnifolium</i>
Dogfennel	<i>Eupatorium capillifolium</i>	Silverleaf nightshade	<i>Polygonum</i> spp.
Fiddleneck, coast	<i>Amsinckia menziesii</i> var. <i>intermedia</i>	Smartweed	<i>Rumex</i> spp.
Filaree	<i>Erodium</i> spp.	Sorrell	<i>Sonchus</i> spp.
Fleabane	<i>Erigeron</i> spp.	Sowthistle	<i>Euphorbia</i> spp.
Giant ragweed	<i>Ambrosia trifida</i>	Spurge, annual	<i>Urtica dioica</i>
Goldenrod	<i>Solidago</i> spp.	Stinging nettle	<i>Helianthus</i> spp.
Grey rabbitbrush	<i>Chrysothamnus nauseosus</i>	Sunflower	<i>Melilotus</i> spp.
Henbit	<i>Lamium aplexicaule</i>	Sweet clover	<i>Descurainia pinnate</i>
Hoary vervain	<i>Verbena stricta</i>	Tansymustard, pinnate	<i>Cirsium texanum</i>
Horseweed	<i>Conyza canadensis</i>	Texas thistle	<i>Abutilon theophrasti</i>
Indian mustard	<i>Brassica juncea</i>	Velvetleaf	<i>Ambrosia psilostachya</i>
Japanese bamboo/knotweed	<i>Polygonum cuspidatum</i>	Western ragweed	<i>Daucus carota</i>
Knotweed, prostrate	<i>Polygonum aviculare</i>	Wild carrot	<i>Lactuca</i> spp.
Kochia	<i>Kochia scoparia</i>	Wild lettuce	<i>Pastinaca sativa</i>
Lambsquarters	<i>Chenopodium album</i>	Wild parsnip	<i>Brassica campestris</i>
Little mallow	<i>Malva parviflora</i>	Wild turnip	<i>Ambrosia grayi</i>
Milkweed	<i>Asclepias</i> spp.	Woollyleaf bursage	<i>Centaurea solstitialis</i>
Miner's lettuce	<i>Montia perfoliata</i>	Yellow starthistle	<i>Oxalis stricta</i>
		Yellow woodsorrel	

VINES AND BRAMBLES

The species of vines and brambles controlled by Prep-It Herbicide include the following:

Field bindweed	<i>Convolvulus arvensis</i>	Virginia creeper	<i>Parthenocissus</i> <i>quinquefolia</i>
Hedge bindweed	<i>Calystegia sequium</i>		<i>Polygonum convolvulus</i>
Honeysuckle	<i>Lonicera</i> spp.	Wild buckwheat	<i>Vitis</i> spp.
Morningglory	<i>Ipomoea</i> spp.	Wild grape	<i>Rosa</i> spp.
Poison ivy	<i>Rhus radicans</i>	Wild rose	Including Multiflora rose (<i>Rosa multiflora</i>)
Redvine	<i>Brunnichia cirrhosa</i>		Macartney rose (<i>Rosa</i> <i>bracteata</i>)
Trumpet creeper	<i>Campsis radicans</i>		

Prep-It® HERBICIDE
EPA REG. NO. 34704-989

WOODY BRUSH AND TREES

The species of woody brush and trees controlled by Prep-It Herbicide include the following:

Alder	(<i>Alnus</i> spp.)	Madrone, Pacific	<i>Arbutus menziesii</i>
American beech	(<i>Fagus grandifolia</i>)	Maple	<i>Acer</i> spp.
Ash	(<i>Fraxinus</i> spp.)	Melaleuca	<i>Melaleuca quinquenervia</i>
Aspen	(<i>Populus</i> spp.)	Mulberry ²	<i>Morus</i> spp.
Autumn olive	(<i>Elaeagnus umbellata</i>)	Oak	<i>Quercus</i> spp.
Bald cypress	(<i>Taxodium distichum</i>)	Persimmon ¹	<i>Diospyros virginiana</i>
Bigleaf maple	(<i>Acer macrophyllum</i>)	Poison oak	<i>Rhus diversiloba</i>
Birch	(<i>Betula</i> spp.)	Popcorn-tree	<i>Sapium sebiferum</i>
Black oak	(<i>Quercus kelloggii</i>)	Poplar	<i>Populus</i> spp.
Blackgum ¹	<i>Nyssa sylvatica</i>	Privet	<i>Ligustrum vulgare</i>
Boxelder	<i>Acer negundo</i>	Red alder	<i>Alnus rubra</i>
Brazilian peppertree	<i>Schinus terebinthifolius</i>	Red maple	<i>Acer rubrum</i>
Ceanothis	(<i>Ceanothis</i> spp.)	Saltcedar	<i>Tamarix pentandra</i>
Cherry ¹	<i>Prunus</i> spp.	Sassafras	<i>Sassafras albidum</i>
Chinaberry	<i>Melia azadarach</i>	Sourwood ¹	<i>Oxydendrum arboreum</i>
Chinese tallow-tree	<i>Sapium sebiferum</i>	Sumac	<i>Rhus</i> spp.
Chinquipin, giant	<i>Castanopsis chrysophylla</i>	Sweetgum	<i>Liquidambar styraciflua</i>
Cottonwood	<i>Populus trichocarpa</i> and <i>Populus deltoides</i>	Sycamore	<i>Plantus occidentalis</i>
Cypress	<i>Taxodium</i> spp.	Tan oak	<i>Lithocarpus densiflorus</i>
Dogwood	<i>Cornus</i> spp.	Titi	<i>Cyrilla racemiflora</i>
Eucalyptus	<i>Eucalyptus</i> spp.	Tree-of-heaven	<i>Ailanthus altissima</i>
Hawthorn	<i>Crataegus</i> spp.	Vaccinium spp.	<i>Vaccinium</i> spp. Including:
Hickory ¹	<i>Carya</i> spp.		Blueberry
Huckleberry	<i>Gaylussacia</i> spp.		(<i>Vaccinium</i> spp.)
Lyonia	<i>Lyonia</i> spp. Including:		Sparkleberry (<i>Vaccinium</i> <i>arboretum</i>)
	<i>Fetterbush</i>	Willow	<i>Salix</i> spp.
	(<i>Lyonia lucida</i> & <i>Staggerbush</i>	Yellow-poplar	<i>Liriodendron tulipifera</i>
	(<i>Lyonia mariana</i>)		

¹ Best control with applications prior to formation of fall leaf color.

² The degree of control may be species dependent.

Prep-It® HERBICIDE
EPA REG. NO. 34704-989

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: DO NOT store below 10 °F.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at www.acrecycle.org. If not recycled, then puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

For packages up to 5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For packages greater than 5 gallons and less than 56 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For refillable containers: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For help with any spill, leak, fire or exposure involving this material call day or night CHEMTREC 1-800-424-9300.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this

**Prep-It® HERBICIDE
EPA REG. NO. 34704-989**

product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS," AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Compadre and Prep-It are registered trademarks of Loveland Products, Inc.
Thru-Valve Boom is a registered trademark of Waldrum Specialties.

**FORMULATED FOR
LOVELAND PRODUCTS, INC.
P.O. BOX 1286, GREELEY, COLORADO 80632-1286**