

PLATFORM

13-4-8

Guaranteed Analysis

Total Nitrogen (N).....13.0%
3.60% Urea
9.40% Water Soluble Nitrogen*
Available Phosphoric Acid (P₂O₅).....4.0%
Soluble Potash (K₂O)..... 8.0%
Molybdenum (Mo).....0.05%

Derived From: urea-triazone solution, phosphoric acid, potassium carbonate, potassium hydroxide, sodium molybdate

*Slowly available nitrogen from urea-triazone solution

GENERAL INFORMATION:

Platform is a liquid foliar fertilizer, providing a source of Nitrogen, Phosphorus, Potassium and Molybdenum during primary growth stages. This material contains 1.41 lbs of Nitrogen (0.39 lbs as Urea, and 1.02 as Water Soluble Nitrogen), 0.44 lbs of P₂O₅, as Orthophosphoric Acid, 0.87 lbs of Potassium, and 0.005 lbs of Molybdenum, in each gallon. This product is for use in foliar sprays. It is intended to be a supplement to a fertility program. Always monitor crop nutrient needs and responses with a soil, leaf tissue and irrigation water analysis program.

DIRECTIONS FOR USE:

FOLIAR APPLICATIONS:

This material may be foliar applied. In a clean spray tank, add any dry powders and water and agitate. Add liquid fertilizers and this material last, continuing agitation. Always use enough water for uniform application and to prevent possible plant tissue damage to the intended crop from concentrated droplets. Contact your Source to Source representative for ultra-low volume rates. Apply during the cool part of the day, or when temperatures are below 85° F. Do not apply under windy conditions, or other conditions that promote fast drying of leaf surfaces. Do not store diluted material.

Foliar Spray Concentration Ratio:

Apply 2 quarts in 25-100 gallons of water.

PRECAUTIONS:

- Due to the dark color of this material, slight spotting of the harvested portion of the plant may occur with late season foliar applications. Where spotting may occur, (particularly on light green colored plant material or harvested material that is not washed post-harvest,) do not apply close to the targeted harvest date unless experience dictates otherwise. Apply with an appropriate adjuvant and spray dilution rate to avoid concentration of the spray solution during the drying period.

- Avoid applications that may cause this product to come into direct contact with the harvested portion of the plant if it will not be thoroughly washed before consumption.
- In crops where fruit maturity may be delayed by nitrogen applications, do not apply this product in late season sprays.
- This product is not recommended for applications within 30 days of harvest or at any time when added nitrogen will adversely affect crop quality or marketability.
- Avoid or reduce applications on Nitrogen sensitive varieties.
- Agitate prior to use.

COMPATIBILITY:

This product has a mildly alkaline pH and is compatible with many commonly applied nutrients and crop protection products. Contact a Source To Source representative for a list of compatible mixes. Jar test for compatibility prior to mixing with any pesticides, fertilizers or fertilizer blends. For unfamiliar mixes, an appropriate amount of the final mix should be applied to a small area of the intended crop to ensure plant compatibility prior to the planned application. User assumes full responsibility to ensure product, equipment and plant compatibilities.

STORAGE AND SHELF LIFE:

Store in a sealed container in cool, dry location, at temperatures above freezing. Triple rinse and dispose of containers according to local and federal regulations.

SUGGESTED APPLICATION RATES AND TIMINGS

TREE CROPS

<p>TREE NUTS (Almonds, Walnuts, Pecans, Pistachios)</p>	<p>Mature Trees: Apply 2 quarts per acre at pre-bloom, bloom, petal fall and nut sizing, for a total of 8 quarts per acre. Post harvest, apply an additional 2 quarts per acre.</p> <p>Non Bearing Trees: Apply 2 quarts per acre every 10-14 days as needed, during periods of canopy growth. Do not exceed a season maximum total of 10 quarts per acre.</p>
<p>STONEFRUIT (Apricots, Cherries, Nectarines, Peaches, Plums, Plumcots, Pluots, Prunes)</p>	<p>Mature Trees: Apply 2 quarts per acre at pre-bloom, bloom, petal fall and fruit sizing, for a total of 8 quarts per acre. Post harvest, apply an additional 2 quarts per acre.</p> <p>Non Bearing Trees: Apply 2 quarts per acre every 10-14 days as needed, during periods of canopy growth. Do not exceed a season maximum total of 10 quarts per acre.</p>
<p>CITRUS (Fingerlimes, Grapefruits, Lemons, Limes, Mandarins, Oranges, Tangelos)</p>	<p>Mature Trees: Apply 2 quarts per acre at featherleaf, bloom, petal fall and fruit sizing, for a total of 8 quarts per acre. Post harvest, apply an additional 2 quarts per acre.</p> <p>Non Bearing Trees: Apply 2 quarts per acre every 10-14 days as needed, during periods of canopy growth. Do not exceed a season maximum total of 10 quarts per acre.</p>

VINE CROPS

BLUEBERRIES	Mature Vines: Apply 2 quarts per acre at pre-bloom, bloom, petal fall and fruit sizing for a total of 8 quarts per acre. Post harvest, apply an additional 2 quarts per acre. Non Bearing Vines: Apply 2 quarts per acre every 10-14 days as needed, during periods of canopy growth. Do not exceed a season maximum total of 10 quarts per acre.
GRAPES	Mature Vines: Apply 2 quarts per acre starting at budbreak, in 10-14 day intervals up through fruit sizing as needed, for a total of 8 quarts per acre. Post harvest, apply an additional 2 quarts per acre. Non Bearing Vines: Apply 2 quarts per acre every 10-14 days as needed, during periods of canopy growth. Do not exceed a season maximum total of 10 quarts per acre.
KIWIS	Mature Vines: Apply 2 quarts per acre starting at budbreak, in 10-14 day intervals up through fruit sizing as needed, for a total of 8 quarts per acre. Post harvest, apply an additional 2 quarts per acre. Non Bearing Vines: Apply 2 quarts per acre every 10-14 days as needed, during periods of canopy growth. Do not exceed a season maximum total of 10 quarts per acre.

This material contains 1.41 lbs of Nitrogen (0.39 lbs as Urea, and 1.02 as Water Soluble Nitrogen.)

CAUTION: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. May cause eye and skin irritation.

NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. Buyer shall assume all risks of use including, but not limited to crop injury, ineffectiveness or other unintended consequences as may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of the seller. In no case shall Source to Source be liable for consequential, special or indirect damages resulting from the use or handling of this product.

Always refer to the label on the product before use.

Platform™ is a trademark product of Source to Source, Tulare CA

Information regarding the contents and levels of metals in this product is available by calling your Source To Source Representative at 866-727-4572.

This label is for use in California and Arizona. For use in other states please contact your Source to Source dealer, as labels may vary by state.

CAUTION: This material contains Molybdenum. The use of this material on any other crops other than those recommended may result in serious injury to crops and/or animals. Use only in accordance with directions.

NOT FOR USE IN ORGANIC CROP AND FOOD PRODUCTION IN CALIFORNIA.

10.88 lbs per gallon (1.3 kgs per liter) @ 68F°

- 5 gallons (18.92 liters)
- 250 gallons (946 liters)
- Bulk (_____ gallons)

Guaranteed by:

Source to Source

a TAP family company

3233 South "I" Street

Tulare, CA 93274

Toll Free: 866.727.4572

Ph: 559.686.4425

Fax: 559.686.5294