

BioFlora® Copper

FOR AGRICULTURE

5 Gallons (18.93 liters) • Net Weight 51.0 lb (23.09 kg)

Guaranteed Analysis:

Copper (Cu) 3.00%
3.00% Chelated Copper

Derivation :

Copper ethylene diamine tetraacetate (Cu EDTA)

Density:

10.20 lbs. per gallon; 1.22 kgs. per liter

Directions for Use:

Apply **BioFlora® Copper** alone or in combination with dry or liquid fertilizer for pre-plant, starter or side dress applications. For most effective treatment of row crops, make soil application in a band at planting time or as a side dressing shortly after planting.

For foliar applications dilute **BioFlora® Copper** with water as recommended, this product can be applied as a foliar spray. To ensure even distribution on leaf surface and increase effectiveness of foliar application, a small amount of wetting agent or detergent may be added.

KEEP AWAY FROM CHILDREN. NOT FOR HUMAN CONSUMPTION.

Note: Buyer assumes all risks of use, storage and handling of this material. Neither seller nor its agent make any warranty, expressed or implied, concerning this product, except in conformity with statements on the label.

BIOFLORA SYSTEMS

Manufactured and Guaranteed by:

GLOBAL ORGANICS, LLC

16121 W. EDDIE ALBERT WAY

GOODYEAR, AZ 85338

TELEPHONE: 623-932-1522 WEB: www.bioflora.com

Application Rates:

Corn, Sorghum, Beans, Cotton and Peppers

Soil Application: Apply 1.5 - 3 qts of **BioFlora® Copper** per acre in liquid starter fertilizer. Rate will depend on extent of copper deficiency and soil type.

Foliar Application: Dilute 10 oz. of **BioFlora® Copper** in 100 gallons of water and spray to thoroughly cover foliage. Apply 4 - 6 weeks after planting or when copper deficiency symptoms appear.

Fruit and Nut Crops; Including: Apples, Pears and Citrus

Soil Application: Apply 10 oz. - 3 qts of **BioFlora® Copper** per tree depending on soil conditions and size of tree. It is important to spray material uniformly around the tree and water in thoroughly for rapid response.

Foliar Application: Dilute 10 oz. of **BioFlora® Copper** per 100 gallons of water and spray to thoroughly cover foliage in early spring shortly after trees are fully leafed out. **Do not apply during bloom or just before harvest.**

Storage & Handling:

Store in original container in a cool dry place out of direct sunlight with ambient temperature above 32°F. In case of accidental exposure, flush with plenty of water. **Caution: Avoid contact with eyes. Wash thoroughly after handling.**

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

DISCLAIMER: Not for use in organic crop and organic food production in the State of California.

DISCLAIMER: This product does not contain microorganisms.

