



YaraVita™

**Bud Builder™
WP**



**Net weight
22lb 9.9 kg**

NOTICE

Before using this product, read the entire label

Bud Builder™ WP

0-6-0

A formulated powder mixture of magnesium and phosphorus with zinc and boron

Guaranteed analysis:

Available Phosphate (P₂O₅)	6%
Magnesium (Mg)	24%
0% water soluble Magnesium (Mg)	
Boron (B)	3%
Zinc (Zn)	10%

Derived from Zinc Phosphate, Magnesium Hydroxide, Sodium Borate

Contents made in Great Britain

Guaranteed & Manufactured by Yara UK Ltd., Manor Place, Wellington Road,

The Industrial Estate, Pocklington, York YO42 1DN, U.K.

Manufactured for Yara North America, Inc. 100 North Tampa Street, Suite 3200, Tampa,

FL 33602. Customer Service: 800-234-9376

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

This product is recommended for use in conjunction with a complete fertilization program based on soil or tissue analysis.

UN 3077

ENVIRONMENTALLY HAZARDOUS SUBSTANCE, SOLID,
N.O.S. (trizinc bis(orthophosphate))

**THIS PRODUCT IN NON-BULK PACKAGING IS
A MARINE POLLUTANT WHEN
TRANSPORTED BY VESSEL. HOWEVER, IT IS
NOT DOT REGULATED WHEN
TRANSPORTED BY HIGHWAY ONLY PER 49
CFR, 171.4(c)**

DANGER

Contains disodium octaborate

H360FD: May damage fertility.

May damage the unborn child.

H400: Very toxic to aquatic life.

H411: Toxic to aquatic life with long lasting effects.

P202: Do not handle until all safety precautions have been read and understood.

P280: Wear protective gloves/clothing and eye/face protection.

P273: Avoid release to the environment.

P308+P313: IF exposed or concerned: Get medical attention.

P391: Collect spillage.

P501: Dispose of contents and container according to local regulations.

F1673



15.10.14

WLU59931

YaraVita™ Bud Builder WP™

WARNING

This fertilizer contains boron which may be injurious to certain crops. Contact your county agent or field consultant for specific information.

APPLICATION RATES AND TIMINGS

Apples: 5-10 lbs/acre applied post-harvest before leaf fall. Also, 5 to 10 lbs/acre applied at bud burst.

Water rate: 50 to 100 gallons/acre.

Apricots: 10 lbs/acre applied post-harvest before leaf fall.

Repeat if necessary leaving 10 to 14 days between applications.

Also, 5 to 10 lbs/acre applied from bud burst to start of flowering. Water rate: 50 to 100 gallons/acre.

Blueberries: Apply 3-5 lbs/acre just before onset of leaf drop.

Repeat the application at bud separation the following fruiting season. Water rate: 20 gallons/acre.

Cherries: 5-10 lbs/acre applied post-harvest before leaf fall.

Repeat if necessary leaving 10 to 14 days between applications.

Also, 5-10 lbs/acre applied from bud burst to start of flowering. Water rate: 50 to 100 gallons/acre.

Conifers: 2 applications of 5-10 pounds/acre once there is new season leaf production and again in early autumn.

Water rate: 50 to 100 gallons/acre.

Grapevines: 1 to 2 applications of 5 lbs/acre applied pre-flowering at a 10-14 day interval. Also, 5-10 lbs/acre applied post-harvest before leaf fall. Water rate: 50 to 100 gallons/acre.

Nectarines: 10 lbs/acre applied post-harvest before leaf fall.

Repeat if necessary leaving 10 to 14 days between applications.

Also, 5 to 10 lbs/acre applied from bud burst to start of flowering. Water rate: 50 to 100 gallons/acre.

Nuts (deciduous): 5-10 lbs/acre applied post-harvest before leaf fall. Also, 5 to 10 lbs/acre applied from bud burst to start of flowering. Water rate: 50 to 100 gallons/acre.

Peaches: 5-10 lbs/acre applied post-harvest before leaf fall.

Repeat if necessary leaving 10 to 14 days between applications.

Also, 5-10 lbs/acre applied from bud burst to start of flowering. Water rate: 50 to 100 gallons/acre.

Pears: 5-10 lbs/acre applied post-harvest before leaf fall. Also, 5-10 lbs/acre applied at bud burst.

Water rate: 50 to 100 gallons/acre.

Plums: 5-10 lbs/acre applied post-harvest before leaf fall.

Repeat if necessary leaving 10 to 14 days between applications.

Also, 5-10 lbs/acre applied from bud burst to start of flowering. Water rate: 50 to 100 gallons/acre.

Strawberries (field grown): 3-5 lbs/acre applied from green bud to just prior to start of flowering. Also, 5 lbs/acre post-harvest, pre-leaf senescence. Water rate: 20 to 50 gallons/acre.

Extended recommendations available at

www.yara.us/agriculture/products/yaravita

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (Temperatures exceeding 40°C) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.