



1% Dursban® Granular Bait

For use on golf courses (turf only), road medians and industrial sites only.

For effective long lasting control of Chinch Bugs, Asian Roaches, Fire Ants, Fleas, Ticks, Foraging Ants, Mole Crickets, and other outdoor insects.

ACTIVE INGREDIENT:

Chlorpyrifos: [0,0,-diethyl-0-(3,5,6-trichloro-2-pyridyl) phosphorothioate].....	1.00%
INERT INGREDIENTS	99.00%
Total	100.00%

FOR PROFESSIONAL USE ONLY: mound treatment control of imported fire ants in lawns and recreational areas.

Controls Ants (including Carpenter, Argentine & Foraging), Army-worms (Beet, Fall, Yellow-striped), Centipedes, Chinch Bugs(2), Clover Mites, Cutworms, Crickets, Earwigs, Fiery Skipper, Fleas, Grasshoppers, Greenbugs, Leafhoppers, Millipedes, Mole Crickets (4), Roaches (Asian), Sod Webworms (Lawn Moth) (3), Sow Bugs, Springtails, Ticks (Brown Dog Tick) (1), Imported Fire Ants(5). (See numbered paragraphs for insect control)

Use for crawling insect control outside of industrial sites such as warehouses and food processing establishments.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

If inhaled	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-(800) 424-9300 for emergency treatment information.

NOTE TO PHYSICIAN

Chlorpyrifos is a cholinesterase inhibitor, Treat symptomatically. Atropine is an antidote by injection.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid contact with eyes, skin and clothing. Avoid breathing dust. Wash thoroughly after handling and before eating or smoking. Wash contaminated clothing before reuse. Keep away from food, feed-stuffs, and water supplies. Keep children and pets off treated areas until this material is washed into the soil and the grass is dry.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water, or other areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters. Use only as directed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
Storage: Store product in its original container in a cool, dry locked place out of reach of children.
Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned stay out of smoke.

EPA Reg. No. 10404-67
EPA Est. No. 10404-OH-04 (M). 10404-FL-1(S)
First letter of lot code indicates manufacturing site.

#069232

F1406
LESCO, Inc.
15885 Sprague Road
Strongsville, OH 44136-1799

SPECIFIC DIRECTIONS

The following numbered paragraphs correspond to numbers in parenthesis (x) on the front panel following the insect names. Insects not numbered may be controlled by following the GENERAL DIRECTIONS. Repeat treatments as needed for control.

(1) BROWN DOG TICKS, & FLEAS

Use 1 lb./435 sq ft (2.3 lb./1,000 sq ft) for area control of these insects, infesting golf courses (turf only), road medians and industrial sites where these are present and create a nuisance or a possible public health problem. DO NOT allow public use of treated area during applications.

(2) CHINCH BUGS

Apply 1 lb./435 sq ft (2.3 lb./1,000 sq ft or 100 lb./A) when insects appear and/or their damage is imminent. To prevent damage apply as soon as daily temperatures reach 80 F and repeat every 2 months.

(3) SOD WEBWORMS

For Sod Webworms, delay watering or mowing the treated area for 12 to 24 hours after treatment. Treat in the spring or when insects first appear troublesome. Additional treatments at 30 day intervals may be needed for new infestations. Apply at rate of 2.3 lb per 1,000 sq ft

(4) MOLE CRICKETS

Use 2.3 lbs. Per 1,000 sq. ft. Treat suspected problem areas when insects become a problem or as recommended by your local Ag. Extension Service Specialist.

The Best Time to Control Adult Mole Crickets

Jan Feb Mar Apr May Jun July Aug Sep Oct Nov Dec

1. _____
2. _____

**(5) IMPORTED FIRE ANT CONTROL
FOR PROFESSIONAL USE ONLY**

MOUND TREATMENT-For treatment of Fire Ant mounds occurring in lawns, turf and premise areas treat individual ant mounds measuring from 3 to 12 inches in diameter by applying 8 tablespoons (1.6 oz.) of granules evenly in a circle around the outer edges (1 to 2 feet from the center) of the mound. Finish the treatment by applying 2 tablespoons (0.4 oz.) sprinkled evenly over the center of the mound. After application of the granules, gently sprinkle the treated area with a minimum of 1 quart of water per 6 inches of mound diameter. Apply in cool weather 65 to 80 degrees F) or in early morning or late evening. Best results will be obtained if mound is not disturbed after treatment thus allowing ants to come and go and to be exposed to the insecticide. Repeat treatments as needed for additional control but not more than once every 7 days.

GENERAL DIRECTIONS FOR USE ON GOLF COURSES (TURF ONLY) ROAD MEDIANS AND INDUSTRIAL SITES. Apply 1% Dursban Granular uniformly to golf courses (turf only), using any hand or power operated type spreader designed for application of granules. Treat when insects first appear at the rate of 2.3 lb/1,000 sq ft (100 lb/Acre) for Surface insects. For best efficacy sprinkle the turf with water to move the granules into the zone of insect infestation. Repeat treatment as necessary. Observe Watering limitation in paragraph for No. (3) Sod Webworms.

RATE TO USE FOR SURFACE INSECTS

For control of Ants (as listed on front panel), Armyworms (Beet, Fall, Yellowstriped), Centipedes, Chinch Bugs, Clover Mites Cutworms, Crickets Earwigs, Fiery Skipper, Fleas, Grasshoppers, Greenbugs, Leafhoppers, Millipedes, Sod Webworms (Lawn Moth), Sowbugs, Springtails, and Ticks (Brown Dog Tick) use 2.3 lb/1,000 sq ft

PERIMETER AROUND INDUSTRIAL BUILDINGS/FOOD PROCESSING ESTABLISHMENTS

Use for crawling insect control outside of industrial plant sites such as warehouses and food processing establishments. To protect against insects entering the indoors, a band of granules may be placed around the outside of buildings to kill Ants (as listed on front panel), Ticks (Brown Dog Tick), Crickets, Earwigs, Fleas, Millipedes, roaches (Asian), Sowbugs, Springtails before they attempt to enter. Use 1 pound to treat 435 sq ft (as a band 5 ft. x 87 Ft. long) around the structure. Apply when insects have become troublesome or before they cause problems in the spring or early fall. Repeat one month later and then as needed for control but not more than once every 7 days.

LESCO ROTARY SPREADER APPROXIMATE SETTINGS

For Large Area Treatments

# Product/ 1,000 sq ft	#A/A	Effective Spreading Width	Third Hole	LESCO Calibration Gauge
1.0	0.44	6 ft.	OPEN	# 11
2.3	1.00	6 ft.	OPEN	# 14

***IMPORTANT:** These settings are only approximate. Age, condition of spreader, and operator speed can cause wide variation. Be sure to calibrate your spreader with each application.

NOTICE: LESCO, INC. warrants that this insecticide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this insecticide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

LESCO is a registered trademark of LESCO Technologies, LLC.
Dursban is a registered trademark of Dow Agrosciences, LLC.
G:REGULWPSTRONGSVAprvd Lbl - Word069232.doc Rev.8/7/02 WPBW

Net Weight: 40 lb (18.16 kg)

