

# TEAM\* Pro 0.86% + Turf Fertilizer

A Granular Fertilizer Containing Team Pro herbicide for the Pre-emergence Control of Crabgrass, Foxtail, Goosegrass, *Poa annua*, and Other Annual Grass and Broadleaf Weeds in Established Turfgrass and established landscape ornamentals.

**Active Ingredients:**  
 Benefin: N-butyl-N-ethyl- $\alpha, \alpha$ ,  $\alpha$ -trifluoro-2,6-dinitro-*p*-toluidine..... 0.43%  
 Trifluralin:  $\alpha, \alpha$ ,  $\alpha$ -trifluoro-2,6-dinitro-*N,N*-dipropyl-*p*-toluidine..... 0.43%  
**Inert Ingredients:**..... 99.14%  
 Total..... 100.00%  
 Contains 0.43 pounds active ingredient per 50 pound bag.

**Annual Grasses, Continued:**

<u>Common Name</u> goosegrass (silver crabgrass or crowfoot grass)	<u>Scientific Name</u> <i>Elymus indica</i>
barnyardgrass (watergrass)	<i>Echinochloa crus-galli</i>
green foxtail	<i>Setaria viridis</i>
yellow foxtail	<i>Setaria glauca</i>

**Broadleaf weeds:**

<u>Common Name</u> chickweed, common knotweed, prostrate lambsquarters, common pigweed purslane, common	<u>Scientific Name</u> <i>Stellaria media</i> <i>Polygonum aviculare</i> <i>Chenopodium album</i> <i>Amaranthus</i> spp. <i>Portulaca oleracea</i>
--	---

**Weeds suppressed:**

<u>Common Name</u> panicum, Texas pusley, Florida prostrate spurge spotted spurge creeping spurge creeping woodsorrel woodsorrel	<u>Scientific Name</u> <i>Panicum texanum</i> <i>Richardia scabra</i> <i>Euphorbia humistrata</i> <i>Euphorbia maculata</i> <i>Euphorbia serpens</i> <i>Oxalis corniculata</i> <i>Oxalis</i> spp.
---	--

**APPROVED USES:**

**Established Turfgrass**

**When to Apply:** For optimum results, this product must be applied one to two weeks prior to the anticipated germination of the target annual weeds. A single application will usually provide season-long weed control. However, a second or split application is recommended when the initial application is made well in advance of weed germination or when an extended period of weed control is required, such as areas with longer growing seasons or for lawn care operator programs.

**How to Apply:** Apply this product evenly over the turfgrass area using a properly calibrated drop type or rotary spreader. Calibrate the spreader prior to use according to the manufacturer's directions. Check frequently to make sure equipment is working properly and distributing granules uniformly. More uniform application may be achieved by spreading half the required amount of product over the area and then applying the remaining half in swaths at right angle to the first. Avoid skips and excessive overlaps as poor weed control or turfgrass injury may occur. Avoid use of spreaders that apply material in narrow concentrated bands. A splash board mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.

**Activation:** Optimum weed control performance will be obtained if treated areas receive rainfall or irrigation soon after application.

**Important Use Precautions:** Do not apply this product to golf course putting greens, dichondra, or newly sprigged areas of bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass. Any cultural practices that disturb the soil, such as aeration or verticutting, should be done prior to application. This product will not harm most nearby ornamentals.

**This product may be applied to the following established turfgrass species:**

**Established Cool Season Turfgrasses**

Bentgrass fairways  
Kentucky bluegrass  
Perennial ryegrass  
Fescues

**Established Warm Season Turfgrasses**

Bahiagrass  
Bermudagrass  
Centipedegrass  
Fescue, tall  
St. Augustinegrass  
Zoysiagrass

**BROADCAST APPLICATION RATES:**

Application Timing	Application Rate	Specific Use Directions
<b>Spring:</b> To control spring-to-summer germinating annual weeds. <b>Fall:</b> To control fall-germinating weeds.	174 to 349 lb/acre (4.00 to 8.01 lb/1000 ft <sup>2</sup> )	In cool season and warm season turfgrass areas, make a single application in the spring or fall, 1 to 2 weeks before expected time of weed germination.
<b>Split Application:</b> For extended control of spring and summer annual weeds.	174 lb/acre followed by 174 lb/acre (4.00 lb followed by 4.00 lb/1000 ft <sup>2</sup> )	In cool season turfgrass area and northern half of the warm season turfgrass area, apply the first application in spring 1 to 2 weeks before expected time of weed germination and make the second application 8 to 10 weeks later.
	233 lb/acre followed by 233 lb/acre (5.34 lb followed by 5.34 lb/1000 ft <sup>2</sup> )	In southern half of warm season turfgrass area, apply the first application in spring 1 to 2 weeks before the expected time of weed germination and make a second application 10 to 12 weeks later.

**Reseeding Precautions:** Team herbicide, which controls annual grass weeds, may also affect the establishment of new desirable turfgrass seedlings, if reseeding is attempted too soon after application. When reseeding, it is essential that proper cultural practices, such as soil cultivation, irrigation and fertilization be followed. Reseeding, following the last application of this product, should not be attempted until the applicable minimum time delay in the following table has been met:

Amount of Product Applied Per Acre (Pounds per 1000 sq ft)	Minimum Time Delay Following Last Application Before Reseeding (Weeks)
Single application of 174 lb/acre (4.00 lb/1000 sq ft)	8
Single application of 349 lb/acre (8.01 lb/1000 sq ft)	12 to 16
Split application of 174 lb/acre (4.00 lb/1000 sq ft)	8 to 10
Split application of 233 lb/acre (5.34 lb/1000 sq ft)	10 to 12

**SUGGESTED SPREADER SETTINGS**

Spreader	Ground Speed	Spreading Width	Recommended Rates Lb per 1000 sq ft		
			4.0	5.34	8.01
LESCO Rotary	3 mph	9 ft.	Calibration Gauge # 15	# 17	# 20
Cyclone® or Spyker®			4¼	4½	4¾

**IMPORTANT:** Spreader settings are only approximate. Age, condition of spreader, and operator speed can cause wide variation in application rate. Spreader should be calibrated prior to each application by the operator.

# 19-2-5

**GUARANTEED ANALYSIS**

Total Nitrogen (N)..... 19.00%  
 0.77% Ammoniacal Nitrogen  
 18.23% Urea Nitrogen\*  
 Available Phosphate (P<sub>2</sub>O<sub>5</sub>)..... 2.00%  
 Soluble Potash (K<sub>2</sub>O)..... 5.00%  
 Sulfur (S) Total..... 1.75%  
 1.75% Free Sulfur (S)  
 Derived From: Polymer Plus Sulfur Coated Urea, Urea, Ammonium Phosphate, Muriate of Potash.  
 Chlorine (Cl) Max..... 4.00%  
 \*5.13% Slowly Available Nitrogen from LESCO® Poly Plus® Sulfur Coated Urea.  
 Made in U.S.A. F1406

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**Storage:** Store in original container only. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Notice:** Read the entire label. Use only according to label directions. Before buying this product, read Limited Warranty and Disclaimer Section elsewhere on this label. In case of emergency endangering health or the environment involving this product, call collect 517-636-4400. Do Not Ship or Store with Food, Feeds, Drugs, or Clothing.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing dust and contact with eyes, skin, or clothing. The granular formulation may cause skin sensitization reactions in certain individuals. Use protective clothing such as coveralls or a long-sleeved shirt and long pants, shoes and socks, and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**FIRST AID**

**If in eyes:** Flush eyes with plenty of water. Call a physician if irritation persists.

**If swallowed:** Do not induce vomiting. Call a physician or Poison Control Center. Drink promptly a large quantity of milk, egg white, gelatin solution, or, if these are not available, large quantities of water. Avoid alcohol.

**If on skin:** Wash with plenty of soap and water. Get medical attention if irritation develops.

**If inhaled:** If breathing difficulty occurs, move individual to fresh air and get medical attention. If not breathing, provide cardiopulmonary resuscitation assistance (mouth-to-mouth) and get medical attention immediately.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sites. Do not contaminate water when disposing of equipment washwaters.

**General Information:** This fertilizer product contains Team\* Pro, a pre-emergence herbicide for control of annual grasses and broadleaf weeds in established turfgrasses and established landscape ornamentals.

**Weeds Controlled**

**Annual Grasses:**

<u>Common Name</u> annual bluegrass crabgrass, smooth crabgrass, large or hairy	<u>Scientific Name</u> <i>Poa annua</i> <i>Digitaria ischaemum</i> <i>Digitaria sanguinalis</i>
--	--

**#080387**

# TEAM\* Pro 0.86% + Turf Fertilizer

## Established Ornamental Plants in Landscape Settings:

This product is recommended for pre-emergence control of annual grasses and certain broadleaf weeds in a wide variety of established ornamental trees, shrubs and flowers.

**Application timing:** Apply one to two weeks prior to the expected germination time of target annual weeds or to clean cultivated weed free areas. Do not apply this product to plantings when the foliage is wet.

**Soil Preparation:** This product can be surface applied around established plants or on ornamental beds. Apply to a smooth soil surface (no large clods) to assure uniform distribution. Clean-cultivate and hoe to remove existing weeds before applying this product.

**Application Equipment and Application Techniques:** See General Information section.

**Activation:** Optimum weed control performance will be obtained when followed by overhead irrigation or rainfall, within a few hours after application.

## BROADCAST APPLICATION RATES:

Season of Application	Pounds per 1000 sq ft
Spring Application (Summer Annuals)	4.00 to 8.01 lb/1000 sq ft
Fall Application (Winter Annuals)	4.00 to 8.01 lb/1000 sq ft

In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10-12 weeks following the initial application.

This product may be used for annual weed control in landscape settings on the following established ornamental plant species:

## ORNAMENTAL TREES:

### Common Name

almond  
apple, crabapple  
apricot  
ash, white  
baldcypress  
birch, European white  
blackgum  
cherry  
chestnut, Chinese  
cottonwood  
dogwood, flowering  
dogwood, kousa  
douglas fir  
fir, balsam  
hemlock, Canada  
honey locust  
larch, Japanese  
locust, black  
maple, Norway  
maple, red  
maple, silver  
maple, sugar  
oak, pin  
oak, scarlet  
peach  
pine, Austrian  
pine, eastern white  
pine, Japanese black  
pine, loblolly  
pine, red  
pine, scotch  
planetree, London  
plum  
redbud, eastern  
spruce, Colorado  
spruce, Norway  
spruce, white  
sweetgum  
sycamore  
tuliptree  
walnut, black

### Scientific Name

*Prunus dulcis*  
*Malus* spp.  
*Prunus armeniaca*  
*Fraxinus americana*  
*Taxodium distichum*  
*Betula pendula*  
*Nyssa sylvatica*  
*Prunus* spp.  
*Castanea mollissima*  
*Populus deltoides*  
*Cornus florida*  
*Cornus kousa*  
*Pseudotsuga menziesii*  
*Abies balsamea*  
*Tsuga canadensis*  
*Gleditsia triacanthos*  
*Larix kaempferi*  
*Robinia pseudoacacia*  
*Acer platanoides*  
*Acer rubrum*  
*Acer saccharinum*  
*Acer saccharum*  
*Quercus palustris*  
*Quercus coccinea*  
*Prunus persica*  
*Pinus nigra*  
*Pinus strobus*  
*Pinus thunbergiana*  
*Pinus taeda*  
*Pinus resinosa*  
*Pinus sylvestris*  
*Platanus acerifolia*  
*Prunus* spp.  
*Cercis canadensis*  
*Picea pungens*  
*Picea abies*  
*Picea glauca*  
*Liquidambar styraciflua*  
*Platanus occidentalis*  
*Liriodendron tulipifera*  
*Juglans nigra*

## ORNAMENTAL SHRUBS:

### Common Name

andromeda, Japanese  
arborvitae, American  
azalea  
barberry, Japanese  
barberry, mento  
boxwood, common  
boxwood, littleleaf  
camellia, Japanese  
camellia, sasanqua  
cherry laurel, American  
cinquefoil  
cleysera, Japanese  
cotoneaster, cranberry  
cotoneaster, zabel  
deutzia  
elaegnus, silverberry  
euonymus, spreading  
euonymus, winged  
euonymus, wintercreeper  
firethorn  
forsythia  
guava, pineapple  
holly  
honeysuckle  
indiahawthorn  
juniper  
laurel, mountain  
lilac, common  
mock orange  
pittosporum, Japanese  
privet  
redcedar, eastern  
rhododendron  
spiraea, vanhouttei  
viburnum  
weigela

### Scientific Name

*Pieris japonica*  
*Thuja occidentalis*  
*Rhododendron* spp.  
*Berberis thunbergii*  
*Berberis mentorensis*  
*Buxus sempervirens*  
*Buxus microphylla*  
*Camellia japonica*  
*Camellia sasanqua*  
*Prunus caroliniana*  
*Potentilla* spp.  
*Cleyera japonica*  
*Coloneaster apiculata*  
*Coloneaster zabelii*  
*Deutzia* spp.  
*Elaeagnus pungens*  
*Euonymus kiautschovica*  
*Euonymus alatus*  
*Euonymus fortunei*  
*Pyracantha* spp.  
*Forsythia* spp.  
*Feijoa sellowiana*  
*Ilex* spp.  
*Lonicera* spp.  
*Raphiolepis indica*  
*Juniperus* spp.  
*Kalmia latifolia*  
*Syringia vulgaris*  
*Philadelphus* spp.  
*Pittosporum tobira*  
*Ligustrum* spp.  
*Juniperus virginiana*  
*Rhododendron* spp.  
*Spiraea vanhouttei*  
*Viburnum* spp.  
*Weigela* spp.

## ORNAMENTAL SHRUBS, CONTINUED:

### Common Name

willow  
yew, anglojap  
yew, Japanese  
yewpine

### Scientific Name

*Salix* spp.  
*Taxus media*  
*Taxus cuspidata*  
*Podocarpus macrophyllus*

## ORNAMENTAL GROUND COVER PLANTINGS:

### Common Name

aaronsbeard  
bellflower, adriatic  
bellflower, poscharsky  
ceanothus  
coreopsis  
cotoneaster  
coyote brush  
crown vetch  
daisy, trailing African  
fern, asparagus  
gazania  
germander  
ice plant, largeleaf  
ivy, Algerian  
ivy, English  
lily-of-the-nile  
lilyturf, bigblue  
marigold  
myoporum  
plumbago, dwarf  
rockrose  
rosemary  
rupturewort  
snow-in-summer  
speedwell  
st. johnswort  
stonecrop (sedum)  
strawberry, beach  
thrift  
verbena  
wirevine, creeping  
yarrow, wolly

### Scientific Name

*Hypericum calycinum*  
*Campanula elatines*  
*Campanula poscharskyana*  
*Ceanothus* spp.  
*Coreopsis* spp.  
*Cotoneaster* spp.  
*Baccharis pilularis*  
*Coronilla vana*  
*Osteospermum fruticosum*  
*Asparagus densiflorus*  
*Gazania* spp.  
*Teucrium chamaedrys*  
*Carpobrotus edulis*  
*Helix canariensis*  
*Hedera helix*  
*Agapanthus* spp.  
*Liriope muscari*  
*Tagetes* spp.  
*Myoporum laetum*  
*Ceratostigma umbaginoides*  
*Cistus* spp.  
*Rosmarinus officinalis*  
*Herniaria glabra*  
*Cerastium tomentosum*  
*Veronica* spp.  
*Hypericum coris*  
*Sedum* spp.  
*Fragaria chiloensis*  
*Armeria maritima*  
*Verbena* spp.  
*Muehlenbeckia axillaris*  
*Achillea tomentosa*

## Roses and Other Established Flowers:

African daisy  
aster (perennial)  
balsam  
blackeyed susan  
calendula  
carnation  
centaurea, velvet  
chrysanthemum  
coreopsis  
cornflower  
cosmos  
dahlia  
dianthus  
dusty miller  
floss flower  
forget-me-not  
four o'clock  
gaillardia  
gladiolus  
golden glow  
impatiens  
ixora  
lobelia  
lupine

marigold  
marigold, cape  
morninggloory  
nasturtium  
nicotiana  
petunia  
phlox  
pincushion flower  
poppy, California  
portulaca  
rose  
salvia  
shasta daisy  
snapdragon  
snow-on-the-mountain  
stock  
sunflower  
sweet alyssum  
sweet pea  
sweet sultan  
sweet william  
vinca  
yarrow  
zinnia

## WARRANTY DISCLAIMER

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

## INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

## LIMITATION OF REMEDIES

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

- Refund of purchase price paid by buyer or user for product bought, or
- Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses. The terms of the "Warranty Disclaimer" above and this "Limitation of Remedies" cannot be varied by any written or verbal statements of agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the "Warranty Disclaimer" or this "Limitation of Remedies" in any manner.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn. RA Dept 15885 Sprague Road, Strongsville, OH 44136 referring to the item number found on this bag.

Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.regulatory-info-a1.com>

Team is a registered trademark of Dow Agrosiences LLC. LESCO and Poly Plus are registered trademarks of LESCO Technologies, LLC. Cyclone and Spyker are registered trademarks of Spyker Spreaders, LLC.

G:\REGULWP\STRONGSVL\Labels in Progress\Team\080387.doc Rev. 11/14/02 BWB (073197)

**Net Weight: 50 lb (22.7 kg)**

EPA Reg. No. 62719-289-10404

EPA Est. No. 10404-OH-04

# #080387

F1406  
Distributed by: LESCO, Inc.  
15885 Sprague Road  
Strongsville, OH 44136-1799

