



Sanmite[®]

75 WP



Miticide/Insecticide

A wettable powder for commercial use on ornamental plants grown in greenhouses and outdoors.

PACKAGING

Unit Size:

4 x 1 oz. pouch

Units Per Case: 4

Boxes Per Master Case: 4

STOCK #95181

(See next page
for label information)

Distributed by:

Scotts-Sierra Crop Protection Company

14111 Scottslawn Road • Marysville, OH 43041

If you have any questions regarding the use of this product, call TOLL FREE 1-800-492-8255



Miticide/Insecticide

ACTIVE INGREDIENT:

[2-tert-butyl-5-(4-tert-butylbenzylthio)-4-chloropyridazin-3(2H)-one] 75%

OTHER INGREDIENTS 25%

TOTAL 100.0%

Plus water-soluble packaging

KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

If inhaled: Move person to fresh air. If not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth. Call a poison control center or doctor for further treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING. May be fatal if inhaled. Do not breathe dust or spray mist. For handling activities, use dust/mist filtering respirator (MSHA/NIOSH approval numbers prefix TC-21C), or a NIOSH approved respirator with a N, P, R, or HE prefilter. Wear long-sleeved shirt and long pants, socks and shoes and waterproof gloves. Harmful if swallowed or absorbed through skin. Avoid contact with skin. Remove contaminated clothing and wash before reuse. Causes moderate eye irritation. Do not get in eyes or on clothing. Wear goggles, face shield, or safety glasses. Wash thoroughly with soap and water after handling.

EPA Reg. No. 7969-106-58185

EPA Est. No. 67545-AZ-1

NET WEIGHT: .25 Lb.

(4 x 1 oz. water-soluble bags per pouch)

©2002, The Scotts Company. World rights reserved.

Sanmite® is a registered trademark of Nissan Chemical Industries, Ltd.



Scotts-Sierra Crop Protection Company
14111 Scottslawn Road
Marysville, OH 43041
1-800-492-8255

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Protective eyewear
- Shoes plus socks
- For handling activities, use dust/mist filtering respirator (MSHA/NIOSH approval numbers prefix TC-21C), or a NIOSH approved respirator with a N, P, R, or HE prefilter.
- Chemical-resistant headgear for overhead exposure

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not re-use them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high-water mark. Keep out of lakes, ponds, or streams. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target area. Drift or runoff from treated areas may be hazardous to fish in adjacent sites. This product is toxic to bees. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area. Application early in the morning or at dusk is suggested.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation. For end use only. Do not repackage or reformulate without manufacturer's written approval.

All applicable directions, restrictions, precautions and Disclaimer and Limitation of Liability are to be followed. This labeling must be in the user's possession during application.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear
- For handling activities during handgun applications with direct overhead exposures, wear either a respirator with an organic vapor removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-21C).
- For all other exposures, wear a dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C).

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place away from heat or open flame. This package contains water-soluble bags inside a foil liner (overwrap). Do not remove the water-soluble bags from the overwrap except for immediate use.

If all the water-soluble bags are not used, carefully reseal the overwrap. The water-soluble bags may break if they are exposed to moisture, handled excessively, or handled with wet hands or wet gloves.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal:

- **Water-soluble packaging:** The outer case and inner overwrap packaging of the water-soluble bag should be incinerated or disposed of in a sanitary landfill, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. Do not re-use the empty packaging.

In Case of Emergency

In case of large-scale spillage regarding this product, call:
CHEMTREC 800-424-9300

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment.
- Your local poison control center (hospital).

Steps to be taken in case material is released or spilled:

Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal. Remove contaminated clothing, and wash affected skin areas with water. Wash clothing before re-use. Keep the spill out of all sewers and open bodies of water.

I. GENERAL INFORMATION

Sanmite® 75WP miticide/insecticide is intended for control of mites and whiteflies on ornamental plants, flowers, and foliage crops. **Sanmite® 75WP** provides excellent knock-down and residual control. A good evaluation of performance can generally be made 4-7 days after treatment.

Crop Tolerance

All crops listed in **Table 2. Plant Species Tested for Tolerance to Sanmite® 75WP** are tolerant to **Sanmite® 75WP**.

Mode of Action

Sanmite® 75WP works primarily through contact action. Treat plants when pests are immature or at a susceptible stage and populations are building, before crop damage occurs.

Resistance Management

Using **Sanmite® 75WP** in successive miticide applications is not recommended. Use **Sanmite® 75WP** as part of a sound resistance management program that includes rotation with other treatments having different modes of action.

Spray Coverage

Apply **Sanmite® 75WP** in sufficient water to ensure thorough coverage of foliage. Thorough coverage is required for optimum control. To achieve adequate coverage, use proper spray pressure, nozzles, nozzle spacing, and volume per acre. Consult spray nozzle and accessory guide for information pertaining to proper equipment calibration.

Cleaning Spray Equipment

Clean spray equipment thoroughly using a strong detergent or commercial sprayer cleaner according to the manufacturer's directions before and after applying this product.

II. APPLICATION INSTRUCTIONS

Apply **Sanmite® 75WP** at rates recommended in **Table 1**.

Application Rates. Avoid drift to all other crops and nontarget areas.

Table 1. Application Rates

Pest	Rate per 100 gallons of water
Broad mite European Red Mite Southern red mite Tumid mite Two-spotted spider mite	4 bags (4 ounces)
Whiteflies	4-6 bags (4-6 ounces)

Table 2. Plant Species Tested for Tolerance to Sanmite® 75WP miticide/insecticide *

Common Name	Scientific Name
Ageratum, Blue Blazer	<i>Ageratum houstonianum</i>
Aluminum Plant	<i>Pilea cadierei</i>
Alyssum	<i>Lobularia maritima</i>
Anthurium	<i>Anthurium spp.</i>
Arborvitae, American	<i>Thuja occidentalis, smaragd</i>
Aster, Rainbow	<i>Aster sp.</i>
, Solidago	
Azalea	<i>Rhododendron sp.</i>
Barberry, Japanese Red Leaf	<i>Berberis thunbergii (atropurpureum)</i>
Begonia	<i>Begonia semperflorens</i>
Boxwood, Japanese	<i>Buxus japonica</i>
Caladium	<i>Caladium sp.</i>
Camellia	<i>Camellia japonica</i>
Carnation, Pallas Londerga	<i>Dianthus caryophyllus</i>
, Pink Candy	
Celosia, Dwarf Mixed	<i>Celosia argenta</i>
Chamaedorea Palm	<i>Chamaedorea elegans</i>
Chrysanthemum	<i>Chrysanthemum spp.</i>
Christmas Cactus	<i>Schlumbergera bridgesii</i>
Cimmaron	<i>Medicago sativae</i>
Cinquefoil	<i>Potentilla fruticosa 'May white'</i>
Coleus, Scarlet Wizard	<i>Coleus hybridus</i>
Cosmos	<i>Cosmos sp.</i>
Croton, Pictum	<i>Codiaeum variegatum</i>
Cyclamen, Red	<i>Cyclamen persicum</i>
, White	
Dahlia	<i>Dahlia spp.</i>
Daylily	<i>Hemerocallis spp.</i>
Dianthus, Pink	<i>Dianthus spp.</i>
, Telstar Lavender	
, Telstar White	
Dieffenbachia, Dumb cane	<i>Dieffenbachia sp.</i>
Dogwood, Cornelian Cherry	<i>Cornus spp.</i>
Dusty Miller	<i>Centaurea cineraria</i>
Dracaena	<i>Dracaena marginata</i>
Elm	<i>Ulmus spp.</i>
Euonymus	<i>Euonymus spp.</i>
False Cypress	<i>Chamaecyparis pisifera</i>
Fern, Pteris	<i>Pteris biaurita</i>
, Asparagus	<i>Asparagus setaceus</i>
, Maidenhair	<i>Adiantum sp.</i>
Fir, Fraser	<i>Abies fraseri</i>
, Noble	<i>Abies porcera</i>
Fire Thorn	<i>Pyracantha coccinea</i>
Fuchsia	<i>Fuchsia sp.</i>
Gardenia, August Beauty	<i>Gardenia jasminoides</i>
Geranium, Scarlet Orbit	<i>Geranium sp.</i>
Gerbera Daisy	<i>Gerbera sp.</i>
Gloxinia	<i>Sinningia speciosa</i>
Gold Dust Plant	<i>Aucuba japonica</i>
Goldfish plant	<i>Alloplectus nummularia</i>
Hibiscus	<i>Hibiscus spp.</i>
Holly, Chinese	<i>Ilex cornuta</i>
, Burford	<i>Ilex cornuta 'Burfordii'</i>
, Japanese	<i>Ilex crenata</i>
Honeysuckle	<i>Lonicera spp.</i>
Hyacinth, Common	<i>Hyacinthus orientalis</i>
Hydrangea	<i>Hydrangea spp.</i>
Hydrangea-vine	<i>Schizophragma hydrangea</i>
Impatiens, New Guinea hybrids	<i>Impatiens wallerana</i>
, Celsia high energy	
, Sunshine	
Iris, Miniature	<i>Iris spp.</i>

Common Name	Scientific Name
Ivy, Cascade	<i>Hedera helix</i>
, English	
, Spade	
Juniper	<i>Juniperus spp.</i>
Kalanchoe	<i>Kalanchoe sp.</i>
Lilac	<i>Syringa patula</i>
Lily, Easter	<i>Lilium longiflorum</i>
, Calla	<i>Zantedeschia sp.</i>
, Peace	<i>Spathiphyllum sp.</i>
, Mauna Loa	<i>Spathiphyllum sp.</i>
, Orange pixie	<i>Lilium longiflorum</i>
Lobelia	<i>Lobelia spp.</i>
Magnolia	<i>Magnolia spp.</i>
Mandevilla, Pink	<i>Mandevilla sp.</i>
Marigold	<i>Tagetes erecta</i>
Mock Orange	<i>Philadelphus coronarius</i>
Muscari, (Grape Hyacinth)	<i>Muscari spp.</i>
Palm, Parlor (Neantha Bella)	<i>Chamaedorea elegans</i>
Pansy	<i>Viola wittrockiana</i>
Pear, Bradford	<i>Pyrus calleryana 'Bradford'</i>
Petunia, Harmony Boy	<i>Petunia hybrida</i>
, White Cascade	
, White Madness	
Phlox, Summer	<i>Phlox paniculata</i>
Photinia, Red Tip	<i>Photinia x fraseri</i>
Piggyback plant	<i>Tolmiea menziesii</i>
Pinks (Dianthus)	<i>Dianthus spp.</i>
Pink Splash	<i>Hypoestes phyllostachya</i>
Pittosporum	<i>Pittosporum spp.</i>
Poinsettia	<i>Euphorbia pulcherrima</i>
Ponytail plant	<i>Beaucarnea recurvata</i>
Poppy	<i>Papaver spp.</i>
Pothos	<i>Epipremnum aureum</i>
Prayer plant	<i>Maranta leuconeura</i>
Primrose	<i>Primula sp.</i>
Privet	<i>Ligustrum x vicaryi</i>
Redvein Enkianthus	<i>Enkianthus spp.</i>
Rose	<i>Rosa spp.</i>
Rose moss	<i>Portulaca grandiflora</i>
Rhododendron, English Roseum	<i>Rhododendron sp.</i>
Salvia	<i>Salvia splendens</i>
Schefflera	<i>Schefflera actinophylla</i>
Snapdragon	<i>Antirrhinum spp.</i>
Spirea	<i>Spirea spp.</i>
Spruce, Norway Nest	<i>Picea abies 'nidiformis'</i>
Sunflower, Miniature	<i>Helianthus annuus</i>
Syngonium	<i>Syngonium podophyllum</i>
Tobacco, Ornamental	<i>Nicotiana spp.</i>
Trumpetcreeper	<i>Campsis grandiflora</i>
Tulips	<i>Tulipa spp.</i>
Viburnum, Snowball Bush	<i>Viburnum spp.</i>
Verbena, Blaze	<i>Verbena hybrida</i>
, Garden	
, Lemon	
, Scarlet Romance	
Vinca, Little Blanche	<i>Vinca spp.</i>
, Periwinkle	
Violet, African	<i>Saintpaulia ionantha</i>
Wandering Jew	<i>Tradescantia albiflora</i>
Wisteria	<i>Wisteria spp.</i>
Zinnia, Scarlet Flame	<i>Zinnia elegans</i>
, Giant Cactus	
, Lilliput	
, Dreamland	

III. ADDITIVES

In general, no additives are necessary for effective use of **Sanmite® 75WP** miticide/insecticide. However, in situations where local conditions such as hard water are a problem, adjuvants or wetting agents may be used to achieve thorough spray coverage.

Do not place water-soluble bags directly into dormant or summer-spray-type oils. PVA pouches are water soluble, not oil soluble. Do not use with nutritional sprays containing boron. Boron will prevent the bags from dissolving in water. Rinse the tank thoroughly before adding any material in PVA bags.

IV. MIXING ORDER

- 1) **Water:** Begin by agitating a thoroughly clean sprayer tank half full of clean water.
- 2) **Products in PVA bags:** Determine the number of water-soluble bags to be used based on **Table 2**. Place the water-soluble bags into the mixing tank. The water-soluble bags dissolve in water and the contents will disperse. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
- 3) **Water-dispersible products:** (dry flowables, wettable powders, suspension concentrates, or suspo-emulsions)
- 4) **Water-soluble products**
- 5) **Emulsifiable concentrates**
- 6) **Water-soluble additives**
- 7) Remaining quantity water

Maintain constant agitation during application. For more information, refer to section **V. Tank Mixing Application**.

V. GENERAL TANK MIXING INFORMATION

No tank mixes are specifically recommended with this product. The phytotoxic potential of **Sanmite® 75WP** has been assessed on a wide variety of common ornamental plants with no phytotoxicity observed. However, all plant species and their varieties and cultivars have not been tested with possible tank mix combinations, sequential pesticide treatments, and adjuvants and surfactants. Local conditions can also influence crop tolerance and may not match those under which Scotts has conducted testing. Therefore, before using **Sanmite® 75WP**, test the product on a sample of the crop to be treated to ensure that a phytotoxic response will not occur as a result of applications.

VI. General Restrictions and Limitations - All Crops

- Maximum seasonal use rate: Do not exceed 21.34 ounces of **Sanmite® 75WP** miticide/insecticide per acre, per year.
- Restricted Entry Interval (REI): 12 hours
- Do not enter a treated greenhouse or a treated indoor area without protective equipment for 12 hours unless one of the following items is completed:
 - 10 air exchanges
 - 2 hours of system ventilation
 - 4 hours of ventilation using vents, windows, or other passive ventilation
 - All required PPE is worn.
- Sequential Treatment: Do not use **Sanmite® 75WP** in successive miticide applications. Use **Sanmite® 75WP** in rotation with other treatments having different modes of action.
- Do not apply this product through any type of irrigation system.
- Do not apply this product aerially.
- Do not use **Sanmite® 75WP** with nutritional sprays that contain boron.
- Do not apply this product as a smoke, mist, fog, or aerosol.
- Do not repackage or reformulate without manufacturer's written approval. For end use only.

Crops:

This product can be used on the following crops:

Ornamental plants Foliage crops

Look inside for complete **Restrictions and Limitations** and **Application Instructions**.

Pests:

This product controls the following pests:

Common Name	Scientific Name
Broad mite	<i>Polyphagotarsonemus latus</i>
European red mite	<i>Panonychus ulmi</i>
Southern red mite	<i>Oligonychus ilicis</i>
Tumid mite	<i>Tetranychus tumidus</i>
Two-spotted spider mite	<i>Tetranychus urticae</i>
Whitefly, Silverleaf , Greenhouse	<i>Bemisia argentifolii</i> <i>Trialeurodes vaporariorum</i>

IMPORTANT NOTICE: DISCLAIMER AND LIMITATION OF LIABILITY

This product has been researched to provide necessary data to support its use on turf and ornamentals as described on the label. However, it is understood that tests may not have been carried out on all varieties and under all growing conditions on all plants listed on the label. The user should always follow the label directions and exercise his judgment and caution when using this product on a given variety until familiar with the results under his growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS, OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT IF NOT USED IN ACCORDANCE WITH DIRECTIONS OR ESTABLISHED SAFE PRACTICE. The exclusive remedy of the user or Buyer, and the limit of liability of the Scotts Company or its affiliates, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by a duly authorized representative of The Scotts Company.

