

# STOLLER® HELP (ReZist & Stablizer)

## Specimen Label

**STOLLER ReZist**

0-0-0

U.S. Patent No. 5,997,600

F1601

**ReZist**



Manufactured by Stoller Enterprises, Inc.  
4001 W. Sam Houston Parkway North, Suite 100  
Houston Texas 77043 USA

Phone: 1-... or 71... | Fax: 713-461-4467

E- Mail: [stoller@stollerusa.com](mailto:stoller@stollerusa.com) | Website: [www.StollerUSA.com](http://www.StollerUSA.com)

**Z- REZIST**

# STOLLER ReZist

## Guaranteed Analysis

0-0-0

Copper (Cu)	1.75%
1.75% Chelated Copper	
Manganese (Mn)	1.75%
1.75% Chelated Manganese	
Zinc (Zn)	1.75%
1.75% Chelated Zinc	

Derived from Copper Sulfate, Manganese Oxide and Zinc Oxide, chelated with Ethanol, 2- amino-2- hydroxy-1,2,3- propanetricarboxylate  
MAXIMUM Chlorine (Cl) content is less than 0.1%

## PRECAUTIONARY STATEMENT CAUTION

### Hazards to Humans and Domestic Animals

Harmful if absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes or clothing.

### Personal Protective Equipment

Applicators and other handlers must wear long- sleeved shirt and long pants, waterproof gloves and shoes plus socks. Wash clothing after use with a detergent and hot water.

### User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. If contaminated, remove clothing and wash thoroughly and put on clean clothing.

### Environmental Hazards

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not re- use empty container. Store ReZist™ in a cool place and out of direct sunlight. Dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities by burning. If burned, stay out of smoke.

## GENERAL INFORMATION

Nutrients are essential for the health of plants. ReZist™ contains plant nutrients that are important in plants to maintain their health and vitality under adverse conditions. ReZist™ can be used in combination with fungicides and bactericides to maintain plant tolerance to abiotic stress.

## DIRECTION FOR USE

Tank mix combination with fungicides or bactericides can be prepared by adding 1/2 of the required amount of water to the mix tank. With the agitator running, add the ReZist™ and keep the agitator running and then add the fungicide or bactericide. Maintain agitation until all of the mixture has been applied. Do not let the spray mixture stand overnight in the spray tank. If the compatibility of the tank mixture is not known, before mixing a commercial application conduct a jar test by mixing the proposed mixture in the same ratio in quart jar and letting it stand for 5 minutes. If the combination remains mixed or can be re- mixed readily, it is physically compatible. Use spray pressure and nozzles to produce a fine spray to provide thorough coverage of the crop. Begin treatment when plants are young or new growth begins.

### Fruit, Nut and Ornamental Trees:

**Deciduous:** Apply ReZist™ at 2 quarts per acre (4.6 liter/ hectare). If fungicides are applied, use 1/2 rates with each spray application throughout the season.

**Non- Deciduous:** Apply ReZist™ at 2 quarts per acre (4.7 liter/ hectare) when new shoot growth is observed. Repeat spray in 21 to 28 days. Apply in combination with fungicide or bactericide sprays for ease in application.

### Field, Vegetables and Vine Crops:

Apply ReZist™ and at 1 print/ acre (1.2 liter/ hectare). Begin when seedlings are young or new growth starts with perennial crops. Continue sprays every 21 to 28 days or with the normal fungicide spray schedule used. If disease pressure requires more frequent application of fungicides, increase the amount to 1 quart per acre of ReZist™ as long as these conditions exist.

### Greenhouse and Nursery Crops:

Apply 0.5% solution of ReZist™ as a wetting spray. Use as a tank mix with the fungicide sprays. Spray 7 days before transplanting or harvest.

### Turf and Golf Course Grasses:

Apply 0.5% solution of ReZist™ as a fine setting spray. Combine with fungicide sprays. Do not use more frequently than 21 days.

Information regarding the contents and levels of metals in this product is available on the internet at [http:// www.aapfco.org/ metals.htm](http://www.aapfco.org/metals.htm).

**NOTE:** The information contained herein is true and accurate to the best of our knowledge and belief, but is presented without guarantee since field conditions and use are beyond our control. Neither the manufacturer nor the seller make any warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use of this material when such use is contrary to label instruction. Read and follow the label directions carefully.

# STOLLER® HELP (ReZist & Stablizer)

## Specimen Label

F1601

# STABILIZER



Manufactured by Stoller Enterprises, Inc.  
4001 W. Sam Houston Parkway North, Suite 100  
Houston Texas 77043 USA

Phone: 1-... or 71... | Fax: 713-461-4467

E- Mail: [stoller@stollerusa.com](mailto:stoller@stollerusa.com) | Website: [www.StollerUSA.com](http://www.StollerUSA.com)

Z- STABILIZ

# Stoller Stabilizer

## Guaranteed Analysis

Calcium(Ca):..... 8.0%

Boron (B)..... 0.5%

Derived from calcium chloride and boric acid.

**Net Weight:** 11.45 lb./ gallon

**Freezing Temperature:** 200F

## GENERAL INFORMATION

**STABILIZER** is designed to be sprayed on plant foliage to regulate localized ethylene production within the plant. It will also guard against calcium replacement in cell walls. when used as recommended, **STABILIZER** will reduce internal damage to storage tissue.

## CAUTION: KEEP OUT OF REACH OF CHILDREN

**WARNING:** Harmful if swallowed. Avoid breathing spray mist. May cause irritation of nose, throat and/ or skin. Avoid contact with eyes, skin and clothing. If swallowed: Induce vomiting by sticking finger down throat. Never give anything to an unconscious person. Get medical attention. **If in eyes:** Wash eyes with plenty of clear water for at least 15 minutes. **If on skin:** wash with plenty of soap and water. Get medical attention.

**CONDITIONS OF SALE:** 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than the officers of Seller, is authorized to make any warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product limited to replacement of product or refund of purchase price. Information regarding the contents and levels of metals in this product is available on the internet at [http:// www.aafpco.org/ metals](http://www.aafpco.org/metals).

## DIRECTIONS FOR USE:

**STABILIZER** is designed to be used along with **ReZist™** in order to improve plant resistance to many negative growth factors.

**STABILIZER** should always be tank- mixed with **ReZist™** at the time of application.

## ALWAYS USE EQUAL AMOUNTS OF REZIST AND STABILIZER FOR EACH SPRAY APPLICATION.

Begin treatments when plants are young or when stem appearance is first observed.

**Field Crops:** Foliar spray one quart **STABILIZER** per acre (2.4 liters/ hectare) when plants have 3 to 4 leaves. Do not use less than a 1% solution. Tank mix an equal amount of **ReZist™** with **STABILIZER** for greater effectiveness.

**Vegetable Crops and Potatoes:** Foliar spray one quart of **STABILIZER** per acre (2.4 liters/ hectare) when plants have 2 to 4 leaves. Do not use less than a 1% solution. Repeat sprays on a 14 day schedule. Tank mix an equal amount of **ReZist™** with **STABILIZER** for greater effectiveness.

**Vegetable Crops and Potatoes:** Foliar spray one quart of **STABILIZER** per acre (2.4 liters/ hectare) when plants have 2 to 4 leaves. Do not use less than a 1% solution. Repeat sprays on a 14 day schedule. Tank mix an equal amount of **ReZist™** with **STABILIZER** for greater effectiveness.

**Trees and Ornamental Crops:** Foliar spray a 1% solution in enough water for thorough coverage. Begin when problems are first observed. Repeat applications 21 days after the first one. Add an equal amount of **ReZist™** to the spray tank for increased effectiveness.

## DO NOT MIX WITH FERTILIZER, HERBICIDES OR PLANT GROWTH REGULATORS.

Repeat as needed.

## WARNING

This product contains **Boron** and the use of this product on any crop other than those recommended by this label may result in serious injury to the crop.