

PULL HERE TO OPEN ►



CruiserMaxx[®]

Rice

syngenta[®]

Insecticide with Fungicides

A seed treatment product for protection against damage from, or control of, listed insects and diseases on rice (dry-seeded)

Active Ingredient:

Thiamethoxam ¹	26.40%
Fludioxonil ²	0.28%
Azoxystrobin ³	1.32%
Mefenoxam ⁴	1.65%
Other Ingredients:	70.35%
Total:	100.00%

¹CAS No. 153719-23-4

²CAS No. 131341-86-1

³CAS No. 131860-33-8

⁴CAS No. 70630-17-0, 69516-34-3

CruiserMaxx Rice contains 2.57 pounds thiamethoxam, 0.027 pounds fludioxonil, 0.13 pounds azoxystrobin and 0.16 pounds mefenoxam per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use in booklet.

EPA Reg. No. 100-1369 EPA Est. 100-NE-001

SCP 1369A-L1A 0211
335395

15 gallons
Net Contents

TM

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<p>HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372</p>	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Causes moderate eye irritation. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing vapor or spray mist. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse.

Personal Protective Equipment (PPE)

Applicators and Other Handlers Must Wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material - Category A (e.g., natural rubber ≥ 14 mils)
- Shoes plus socks

continued...

PRECAUTIONARY STATEMENTS *(continued)*

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This product is toxic to wildlife, freshwater and estuarine/marine fish and highly toxic to aquatic invertebrates. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water or rinsate.

This product contains active ingredients that have high potential for runoff for several months or more following application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains these active ingredients. Azoxystrobin degradation products and thiamethoxam have properties and characteristics associated with chemicals detected in ground water. These chemicals may leach into the ground water if used in areas where soils are permeable, particularly where the water table is shallow.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC, or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use in hopper box, planter box, slurry box, or other farmer-applies applications. Apply CruiserMaxx Rice in commercial seed treatment facilities only.

CruiserMaxx Rice may be applied by closed or open system seed treatment application processes. **DO NOT** apply more than 75 gallons of CruiserMaxx Rice per 8-hour day for seed treatments utilizing an open system. There is no volume limitation for closed system applications.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR INSECT AND/OR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigor and poor quality, except for the purpose of curative control of existing disease pests, may result in reduced germination and/or reduction of seed and seedling vigor. Treat a small quantity of seed using equipment similar to that planned for treating the total seed lot. Conduct germination tests on a small portion of seed before committing the total seed lot to a selected seed treatment. Due to seed quality, crop or variety sensitivity, and seed storage conditions beyond the control of Syngenta, no claims are made to guarantee the germination of carry-over seed or propagating material for all crop seed.

USE INFORMATION

CruiserMaxx Rice is a premix of the active ingredients thiamethoxam insecticide and fludioxonil, mefenoxam and azoxystrobin fungicides. Fludioxonil, mefenoxam and azoxystrobin are the active ingredients in Maxim®, Apron XL® and Dynasty® brand fungicides, respectively.

CruiserMaxx Rice is a seed treatment product that protects against damage from certain early-season insects, and controls or suppresses certain soil-borne and seed-borne diseases of dry-seeded rice.

Thiamethoxam is a systemic seed treatment insecticide belonging to the neonicotinoid class of chemistry. Thiamethoxam controls certain chewing and sucking insects through contact and ingestion.

Fludioxonil is active against *Rhizoctonia* and *Fusarium*.

Mefenoxam fungicide is active against *Pythium*.

Azoxystrobin is active against *Rhizoctonia*, *Pythium*, and some soil-borne and seed-borne diseases such as *Pyricularia grisea* (rice blast).

CruiserMaxx Rice is a selective seed treatment insecticide and fungicide, and its use is compatible with integrated pest management programs.

MIXING PROCEDURES

Important: Thoroughly shake the container of CruiserMaxx Rice prior to use.

Apply CruiserMaxx Rice as a water-based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of insect or disease control. Thoroughly mix the specified amount of CruiserMaxx Rice into the required amount of water for the slurry treater and dilution rate to be used. The typical density of CruiserMaxx Rice is 9.68 pounds per gallon. Consult the manufacturer of the application equipment you plan to use for suitability for this application and for instructions on operation and calibration of the equipment.

- Use an EPA-approved dye or colorant that imparts an unnatural color to the seed as stated in 40 CFR 153.155 (c).
- Allow seed to dry before bagging.
- Store away from feed and foodstuffs.

SEED BAG LABEL REQUIREMENTS

The Federal Seed Act requires that bags containing treated seeds shall be labeled with the following statements:

- This seed has been treated with thiamethoxam insecticide and fludioxonil, mefenoxam and azoxystrobin fungicides.
- Do not use for feed, food or oil purposes.
- User is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.

In addition, the U.S. Environmental Protection Agency requires the following statements on bags containing seeds treated with CruiserMaxx Rice:

- Store away from food and feedstuffs.
- Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- Do not contaminate water bodies when disposing of planting equipment wash waters.
- Dispose of seed packaging in accordance with local requirements.
- In the event of crop failure or harvest of a crop grown from CruiserMaxx Rice treated seed, the field may be replanted immediately to barley, canola, corn, cotton, cucurbit vegetables, fruiting vegetables, legume vegetables, mint (peppermint and spearmint), oil seed crops (black mustard seed, borage seed, crambe seed, field mustard seed, flax seed, Indian mustard seed, Indian rapeseed seed, Rapeseed seed, and safflower seed), rice, root vegetables, sorghum, strawberry, sunflowers, tobacco, tuberous and corm vegetables and wheat. For any other crop, the minimum plant back interval is 120 days from the date the CruiserMaxx Rice treated seed was planted. A cover crop other than the crops listed above that is planted for erosion control or soil improvement may be planted sooner than the 120 day interval; however, the crop may not be grazed or harvested for food or feed.
- Do not allow children, pets, or livestock to have access to treated seed.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Dispose of all excess treated seed. Leftover treated seed may be doublesown around the headland or buried away from water sources in accordance with local requirements.
- Do not make any soil or foliar application of products containing thiamethoxam to crops grown from seed treated with CruiserMaxx Rice.
- Do not use at a rate that will result in more than 0.266 lb. thiamethoxam per acre (120.66 grams a.i./A) per season.

CROP USE DIRECTIONS

Resistance Management

Cruiser Maxx Rice contains thiamethoxam, a Group 4A insecticide; fludioxonil, a Group 12 fungicide; mefenoxam, a Group 4 fungicide; and azoxystrobin, a Group 11 fungicide.

Some insect pests are known to develop resistance to products after repeated use. Because resistance development cannot be predicted, the use of this product should conform to sound resistance management strategies established for the crop and use area. Syngenta encourages responsible product stewardship to ensure effective long-term control of the insects on this label.

Note: The fungicide mefenoxam contained in this product could be subject to development of insensitive strains of fungi or may be ineffective against naturally occurring strains of fungi. Development of insensitivity or natural tolerance cannot be predicted. Therefore, Syngenta cannot assume liability for crop damage resulting from insensitive or tolerant strains of fungi. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible insensitive or tolerant strains of fungi which may occur.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

CruiserMaxx Rice contains a Group 4A insecticide (thiamethoxam, belonging to the neonicotinoid class of chemistry). Insect biotypes with acquired or inherent resistance to Group 4A insecticides may eventually dominate the insect population if Group 4A insecticides are used repeatedly as the predominant method of control for targeted species. This may result in partial or total loss of control of those species by CruiserMaxx Rice or other Group 4A insecticides.

In order to maintain susceptibility to this class of chemistry:

- Avoid using Group 4A insecticides exclusively for season long control of insect species with more than one generation per crop season.
- For insect species with successive or overlapping generations, apply a Group 4A insecticide using a "treatment window" approach. A treatment window is a period of time as defined by the stage of crop development and/or the biology of the pests of concern. Within the treatment window, depending on the length of residual activity, there may either be single or consecutive applications (seed treatment, soil, foliar, unless otherwise stated) of the Group 4A insecticides. Following a treatment window of Group 4A insecticides, rotate to a treatment window of effective products with a different mode of action before making additional applications of Group 4A insecticides.
- A treatment window rotation, along with other IPM practices for the crop and use area, is considered an effective strategy for preventing or delaying a pest's ability to develop resistance to this class of chemistry.
- If resistance is suspected, do not reapply any Group 4A insecticides.

Other Insect Resistance Management (IRM) practices include:

- Incorporating IPM techniques into your insect control program.
- Monitoring treated insect populations for loss of field efficacy.
- Using tank-mixtures or premixes with insecticides from a different target site of action group as long as the involved products are all registered for the same crop outlet and effective rates are applied.

For additional information on Insect Resistance Management:

- Contact your local extension specialist, certified crop advisor and/or product manufacturer for additional insect resistance management recommendations.
- Visit the Insecticide Resistance Action Committee (IRAC) on the web at: <http://www.irac-online.org/>.

Use Restrictions

CruiserMaxx Rice may be applied by closed or open system seed treatment application processes. **DO NOT** apply more than 75 gallons of Cruiser per 8-hour day for seed treatments utilizing an open system. There is no volume limitation for closed system applications.

Rice (Dry-Seeded Use Only)

When applied according to the following CruiserMaxx Rice Rate Table, CruiserMaxx Rice provides early season insect protection against injury by grape colaspis, rice water weevil, chinch bugs, aphids and thrips. CruiserMaxx Rice also provides early season disease protection against soil and seed-borne fungi including *Pythium*, *Rhizoctonia*, *Fusarium* and seed-borne *Pyricularia grisea* (rice blast).

CruiserMaxx Rice Rate Table

Crop	Rate of CruiserMaxx Rice	
	fl oz per 100 pounds seed*	Total mg active ingredients per seed (based on 21,000 seeds/pound)
Rice	7.0	0.0337 (0.030 thiamethoxam 0.0003 fludioxonil 0.0015 azoxystrobin 0.0019 mefenoxam)

*Each fluid ounce of CruiserMaxx Rice contains 9.1 grams of thiamethoxam, 0.10 grams fludioxonil, 0.46 grams of azoxystrobin and 0.57 grams mefenoxam.

CruiserMaxx Rice is not approved for use in water seeded rice production.

Do not plant or sow CruiserMaxx Rice treated rice seed by aerial application equipment.

Do not use treated fields for the aquaculture of edible fish and crustacean.

Do not exceed 120 lb. seed/acre seeding rate.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment.

Pesticide Storage

Store in a cool, dry place. Do not store above 90°F for extended periods.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling (<5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Container Handling (>5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other side and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Apron XL®, ApronMAXX®, Cruiser®, Dynasty®, Maxim®, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company

©2011 Syngenta

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1369A-L1A 0211
335395



Insecticide with Fungicides

A seed treatment product for protection against damage from, or control of, listed insects and diseases on rice (dry-seeded)

Active Ingredient:	
Thiamethoxam ¹	26.40%
Fludioxonil ²	0.28%
Azoxystrobin ³	1.32%
Mefenoxam ⁴	1.65%
<hr/>	
Other Ingredients:	70.35%
Total:	100.00%

- ¹ CAS No. 153719-23-4
- ² CAS No. 131341-86-1
- ³ CAS No. 131860-33-8
- ⁴ CAS No. 70630-17-0, 69516-34-3

CruiserMaxx Rice contains 2.57 pounds thiamethoxam, 0.027 pounds fludioxonil, 0.13 pounds azoxystrobin and 0.16 pounds mefenoxam per gallon.

EPA Reg. No. 100-1369
EPA Est. 100-NE-001

Cruiser® and the Syngenta logo are trademarks of a Syngenta Group Company

©2011 Syngenta

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

**SCP 1369A-L1A 0211
335395**

15 gallons

Net Contents

KEEP OUT OF REACH OF CHILDREN. CAUTION

See additional precautionary statements and directions for use in booklet.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

HOT LINE NUMBER: For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call **1-800-888-8372**.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Causes moderate eye irritation. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing vapor or spray mist. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse.

Environmental Hazards: This product is toxic to wildlife, freshwater and estuarine/marine fish and highly toxic to aquatic invertebrates.

Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water or rinsate. This product contains active ingredients that have high potential for runoff for several months or more following application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains these active ingredients. Azoxystrobin degradation products and thiamethoxam have properties and characteristics associated with chemicals detected in ground water. These chemicals may leach into the ground water if used in areas where soils are permeable, particularly where the water table is shallow.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment.

Pesticide Storage: Store in a cool, dry place. Do not store above 90°F for extended periods. For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling: Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¹/₄ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other side and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.



BAR CODE # IS
(01) 0 07 02941 42644
LAST DIGIT IS CHECK DIGIT
UCC/EAN 128