

STOLLER® XYLEX 6-18-6

Specimen Label

A liquid foliar nutrient with
micronutrients

XYLEX



Manufactured for Stoller Enterprises, Inc.
4001 W. Sam Houston Parkway North, Suite 100
Houston Texas 77043 USA
Phone: 1-800-539-5283 or 713-461-1493 | Fax: 713-461-4467
E- Mail: stoller@stollerusa.com | Website: www.StollerUSA.com

Stoller XYLEX 6-18-6

Directions for Use

GUARANTEED ANALYSIS

6-18-6

Total nitrogen (N).....	6.00%
12% Urea Nitrogen	
Available phosphate (P2O5).....	18.00%
Soluble potash (K2O):.....	6.00%
Boron (B):.....	0.02%
Cobalt (Co).....	0.0005%
Copper (Cu).....	0.05%
0.10% water soluble Copper (Cu)	
Iron (Fe).....	0.10%
0.40% water soluble Iron (Fe)	
Manganese (Mn).....	0.05%
00.40% water soluble Manganese (Mn)	
Molybdenum (Mo):.....	0.0005%
Zinc (Zn).....	0.05%
0.60% water soluble Zinc (Zn)	

Derived from urea, potassium chloride, calcium monocarbamide, monohydrogen chloride and N +Calcium.
Freezing temperatures less the 20°F.

Derived from urea, phosphoric acid, potassium chloride, sodium borate, cobalt sulfate, copper sulfate, ferrous sulfate, manganese sulfate, sodium molybdate and zinc sulfate.

Chlorine (Cl) maximum.....0.5%

Non- plant food ingredients:

Sodium gluconate.....4.5%

Purpose: source of complexing agent and

General Information

Xylex foliar nutrient compound is to be used as a supplement to soil-applied nutrients and is particularly beneficial under conditions where soil nutrients are not readily available.

Xylex is a complete liquid nutrient compound and when used in a spray program will prevent nutrient deficiencies.

Xylex will assure adequate nutrients at the critical periods of flower and fruit development. **Xylex** will aid root growth, crop yield and quality.

Xylex has an acidity of about 1.5 pH and can be used as a buffering aid with most fungicides, herbicides and insecticides susceptible to alkaline hydrolysis.

Xylex can be applied by ground or aerial application or by injecting into a sprinkler irrigation system.

Follow this mixing procedure: 1. Water, 2. **Xylex**, 3. Pesticide

CAUTION:

Keep Out Of Reach Of Children

WARNING: This product contains boron and the use of this product on any crop other than those recommended on the label may result in serious injury to the crop.

This product is only to be used on soils which respond to molybdenum.

Crops high in molybdenum are toxic to grazing animals (ruminants). Avoid use with basic copper fungicides.

Conditions Of Sale: 1. Seller warrants this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an officer of Seller is authorized to make any warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

Xylex foliar nutrient and buffering aid is formulated as a soluble concentrate and should be applied with a minimum dilution of 1 to 10 parts water for 1 spray application. Add the recommended amount to the water in the partially filled spray tank and while agitating add the remaining water required and other spray materials. Unless indicated differently use sufficient water to provide thoroughly wetting of the leaves.

Vegetables: Apply 1 to 2 quarts per acre at weekly intervals. Start when crops are a few inches high. Continue sprays up to 2 weeks before final harvest. Use a minimum of 20 gallons/ acre of spray volume in ground sprayer and 5 gallons/ acre in aerial spray equipment.

Tree Fruit and Nuts: Apply 1 to 3 quarts per acre in concentrate or dilute spray equipment using 50 to 400 gallons of water per acre. Application can begin at delayed dormancy and be continued every 2 to 3 weeks during the growing season.

Citrus: Apply 1 to 2 gallons per acre in concentrate or dilute spray equipment beginning at pre- bloom. Four sprays per year are recommended.

Field Crops: Apply 1 to 2 quarts per acre when plant leaves are large enough to absorb the foliar nutrient. Spray again just prior to bloom and repeat at 10 to 14- day intervals until crop is mature.

Recommendations For Use

Xylex may be used on the following crops: Alfalfa, Almonds, Apples, Artichokes, Asparagus, Avocados, Barley, All beans, Broccoli, Cabbage, Carrots, Cauliflower, Celery, Citrus, Corn, Cotton, Grapes, Hops, Lettuce, Melons, Milo, Nectarines, Peaches, Peanuts, Peas, Pecans, Peppers, Pistachios, Potatoes, Rice, Soybean, Sorghum, Strawberries, Sugar beets, Sugar cane, Sunflowers, Sweet corn, Tobacco, Tomatoes, Turnips, Walnuts, Watermelons and Wheat.

WARNING: Harmful if swallowed. Avoid breathing spray mist. May cause irritation of nose, throat, and/ or skin. Avoid contact with eyes, skin and clothing. **IF SWALLOWED:** Induce vomiting by sticking finger down throat. Never give anything by mouth to an unconscious person. Get medical attention. **IF IN EYES:** Wash eyes with plenty of clear water for at least 15 minutes. Get medical attention.