

# Fungicide VII



## FEATURES & USE TIPS:

- Systemic fungicide that prevents and controls dollar spot (including benzimidazole tolerant strains), brown patch, anthracnose and rust.
- New DG PRO™ dispersible carrier - minimizes mower pickup after irrigation.
- Contains triadimefon, the active ingredient used in Bayleton
- Convenient, dry-flowable formula.
- Can be applied with rotary or drop spreader.
- Small particle size - ideal for use on golf course greens and low-cut bentgrass fairways.
- Root absorbed for consistent uptake into the turf.

**PRODUCT CODE:**  
**AGC8573**

Date: 072902  
Supersedes: 000000

## PRODUCT CHARACTERISTICS:

**Product Type:**  
Granular Fungicide

**Particle Size:**  
SGN: 100

**Net Weight:**  
29.18 lbs.

**Coverage:**  
22,000 sq. ft.

## PRODUCT SPECIFICATIONS:

**Active Ingredient:** 0.59% Triadimefon

Andersons Golf Products and DGPro are trademarks of The Andersons Agriservices, Inc.

*To order, contact your local Andersons Golf Products distributor.*

*For more information, contact Andersons at our website:  
[www.andersonsgolfproducts.com](http://www.andersonsgolfproducts.com)  
or call 1-800-225-2639.*

*PRODUCT SHEET*



ACTIVE INGREDIENT

1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-butanone . . . . . 0.59%

OTHER INGREDIENTS . . . . . 99.41%

Total . . . . . 100.00%

EPA Reg. No. 9198-187 EPA Est. No. 538-OH-1S, 9198-OH-1M, 9198-OH-2B, 9198-AL-001A Underlined letter is first letter used in run code on bag

Recommended for Use by Professional Turfgrass Managers

KEEP OUT OF THE REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION, causes eye irritation. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. If in eyes, flush with plenty of water. Get medical attention if irritation persists. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock. Keep children and pets off treated area until material has been washed off the foliage and the area is completely dry.

Environmental Hazards: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

For use on turfgrass areas consisting of Kentucky bluegrass, annual bluegrass, bentgrass, perennial ryegrass, fine fescue, bermudagrass or mixtures where these grasses predominate.

TO PREVENT DOLLAR SPOT, ANTHRACNOSE AND RUST apply to moist or dry foliage at the RECOMMENDED RATE when environmental conditions favor disease activity.

Repeat the application at 2 week intervals throughout the expected disease period.

TO CONTROL ACTIVE DOLLAR SPOT, ANTHRACNOSE AND RUST apply to moist or dry foliage at two week intervals. Use the DOUBLE RATE for light to moderate infestations and the HEAVY RATE for severe infestations. When the disease activity is under control, apply at the RECOMMENDED RATE at two week intervals.

TO PREVENT BROWN PATCH apply to moist or dry foliage at the DOUBLE RATE when environmental conditions favor disease activity. Repeat the application at 2 week intervals throughout the expected disease period.

TO CONTROL ACTIVE BROWN PATCH apply to moist or dry foliage at the HEAVY RATE every two weeks. When the disease activity is under control, apply at the DOUBLE RATE at two week intervals.

To avoid product pick-up lightly irrigate treated areas soon after application.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing. Protect from water and extreme heat.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY

Manufacturer warrants that this product conforms to

the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

## LEGAL RIGHTS

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION. (RVnot040302)

### Suggested Spreader Settings

To provide proper distribution, calibrate your spreader before application.

29.18 lbs treats 22,000 square feet at the RECOMMENDED RATE (1.32 lbs/1,000 square feet)

29.18 lbs treats 11,000 square feet at the DOUBLE RATE (2.65 lbs/1,000 square feet)

29.18 lbs treats 5,500 square feet at the HEAVY RATE (5.30 lbs/1,000 square feet)

SPREADER	GROUND SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS		
			RECOMMENDED RATE	DOUBLE RATE	HEAVY RATE
Andersons SS-2 Drop	3 mph	overlap wheels	2 3/4	4	5
Andersons Rotaries: AccuPro/SR-2000 (Cone 9)	3 mph	10 feet	H 1/2	J	L 1/2
Scotts Rotaries: R-8, R-8A (Cone 9)	3 mph	10 feet	H	I	K
Vicon (03 Series-Standard Spout)	4.5 mph	15 feet	11	13	18
Lely Models WTR, WFR, HR, 1250. (PTO at 450 rpm)	4.5 mph	27 feet	3 1/2 III-A	4 1/4 III-A	5 1/2 III-A

Note: These spreader settings were established using standard equipment available from the spreader manufacturer at swath widths and speeds typically used within the industry. It is recommended that all spreader equipment be calibrated at the time of application to achieve the desired application rate.

SPECIMEN  
LABEL

**AGC8573**

Product of USA

**Manufactured by:**

The Andersons Lawn Fertilizer Division, Inc.  
P. O. Box 119, Maumee, OH 43537  
1-800-225-2639

Visit us on the web at: [www.AndersonsGolfProducts.com](http://www.AndersonsGolfProducts.com)