

Contec[®]DG

Dispersing Granule Technology with
0.103% DIMENSION[®] Turf Herbicide



24-0-10

FEATURES & USE TIPS:

- Patented dispersing granule formula
- Contains Dimension turf herbicide
- 90% of the nitrogen is from MUtech-L

PRODUCT SPECIFICATIONS:

N - (24%):	2.40% Ammoniacal Nitrogen 1.86% Urea Nitrogen 12.93% Water Insoluble Nitrogen 6.81% Other Water Soluble Nitrogen* *6.81% slowly available nitrogen from methylene ureas and ureaform
K - as K₂O - (10%):	Potassium sulfate
S - (6.15%):	Free Sulfur
Active Ingredient:	Dithiopyr (0.103%)

PRODUCT CODE:

AGC2424WDGDM5

Date: 022208
Supersedes: 000000

PRODUCT CHARACTERISTICS:

Product Type:

Combination dispersible granular nutrient and Dimension[®] pre-emergent herbicide

Nitrogen Type:

MUtech-L[™]

Particle Size:

SGN: 150

Net Weight:

50 lbs.

Coverage:

9,000 sq. ft.
@ Medium Rate

Andersons Golf Products and ContecDG[®] are registered trademarks of The Andersons, Inc.; MUtech is a trademark of The Andersons, Inc.; Dimension is a registered trademark of Dow AgroSciences, L.L.C.

To order, contact your local Andersons Golf Products distributor.

For more information, contact Andersons at our websites: www.andersonsgolfproducts.com or www.contecdg.com or call 1-800-225-2639.

Following is a specimen label that may at some point be revised or otherwise become out of date. It is intended as a guide in providing general information regarding the use of this product. Always read and follow the label on the product container.

PRODUCT SHEET

KEEP OUT OF REACH OF CHILDREN CAUTION PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Cause eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

FIRST AID

If in eyes: Hold eye open and flush slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor for treatment advice. Have person sip glass of water to swallow. Do not induce vomiting unless told by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8851 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic life. Do not apply to streams, rivers, ponds, lakes, or other aquatic areas. Do not apply to wetlands or other sensitive areas. Do not apply to areas adjacent to any body of water. Do not apply to areas where the water may be used for drinking or other domestic purposes. Do not apply to areas where the water may be used for irrigation. Do not apply to areas where the water may be used for other purposes. Do not apply to areas where the water may be used for other purposes. Do not apply to areas where the water may be used for other purposes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Read the entire label before using this product. Use only according to label instructions. Not for use on or in any animal, bird, or fish. Do not use on or in any animal, bird, or fish. Do not use on or in any animal, bird, or fish.

New York Use Restrictions: In New York State this product may only be used by professional applicators and at no more than 1.0 pounds per 1,000 square feet (equal to 0.5 pounds active ingredient per acre) per year and is prohibited from use in Nassau and Suffolk Counties.

GENERAL INFORMATION

This product is a herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf courses, lawns, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions.

USE PRECAUTIONS

Turfgrass Safety: This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding. Seeding or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on turfs that have been heavily weakened by weather, pests, disease, chemical, or mechanical treatments, may increase the chances of turf injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

Cool-Season Grasses

- Bahamas creeping (Agrostis palustris)
- Bluegrass, Kentucky (Poa pratensis)
- Fescue, fine* (Festuca arvensis)
- Fescue, tall (Festuca arvensis)
- Poa annua (Poa annua)
- Poa perenne (Poa perenne)

Warm-Season Grasses

- Bermudagrass (Cynodon dactylon)

SPECIMEN LABEL

This product will also control (at the Program 2 rate) or suppress (at the Program 3 rate) the following weeds when applied prior to their emergence: bluegrass (annual), Poa annua, bittersweet, chickweed, vent, geranium (Carolina), pansy, plantain, and sheepsfoot.

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with crop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions by adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A minimum application rate can be made by spreading the remaining half at a right angle to the previous direction.

Avoid steaming, sifting, or other equipment malfunctions by properly adjusting and maintaining equipment. Avoid equipment malfunctions by applying uniform distribution of granules.

When split fertilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emergence.

TABLE 1. Recommended Application Use Rates* (Lbs. of this product/1,000 sq. ft.)

PROGRAM USE (Turfgrass or)	1 (Lbs. of this product/1,000 sq. ft.)		2 (Lbs. of this product/1,000 sq. ft.)		3 (Lbs. of this product/1,000 sq. ft.)	
	Preemergence (high)	Postemergence (low)	Preemergence (high)	Postemergence (low)	Preemergence (high)	Postemergence (low)
Region						
North	2.8	4.0	4.0	5.5		
Transition	4.0	5.5	8.4	11.0**		
South	5.5	8.4	11.0**			
Coastal South	8.4	11.0**				

**DO NOT apply more than 11 lbs. per application, and no more than 33 lbs. per year. Postemergence control is limited; see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below.

North: All other areas.
Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, NY, & RI.

South: AL, AR, AZ, CA, GA, LA, MS, NC, NN, NV, OK, SC, TN, & TX.

Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX.

**May require split or sequential applications for turf control.

Fall Applications: This product can also be applied in the fall in areas where the turfgrass is dormant. Apply this product in the early fall of the year. The fall application should be followed by an appropriate spring application to provide season-long control.

Turf for Improved Control: For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application. When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result. Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control.

CONTROL OF OTHER GRASS AND BROADLEAF WEEDS

Spring Applications: Used as directed for crabgrass control in the spring, this product will also control (at the Program 3 rate) or suppress (at the Program 2 rate) the following weeds when applied prior to their emergence: bahamagrass; bluegrass (annual); roughstalk, cowgrass (annual & perennial); smutgrass; searocket; Kentucky bluegrass (annual & perennial); annual ryegrass; perennial ryegrass; bermudagrass (annual & perennial); sturge (prostrate & spotted). Also controlled at the Program 2 rate.

Fall Applications: Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring.

SUGGESTED SPREADER SETTINGS

These suggested spreader settings are not intended to replace calibration. Please calibrate your spreader before applying product.
50 lbs. covers 17,000 sq. ft. (0.29 acre) at 174 lbs./1,000 sq. ft.
50 lbs. covers 15,000 sq. ft. (0.27 acre) at 174 lbs./1,000 sq. ft.
50 lbs. covers 9,000 sq. ft. (0.16 acre) at 240 lbs./1,000 sq. ft.
50 lbs. covers 5,000 sq. ft. (0.14 acre) at 360 lbs./1,000 sq. ft.

SPREADER	GROUND SPEED (ft./min.)	SPREADER SETTINGS			
		LOW RATE (0.1 lb./ft.²)	MEDIUM RATE (0.25 lb./ft.²)	HIGH RATE (0.5 lb./ft.²)	AREA
Andersen-Koehn 200, 395-200 (cone 9)	3 mph	11 feet	M	N 1/2	O 1/2
Andersen-Koehn 1000	3 mph	11 feet	M	N 1/2	O 1/2
Scotch & R. 9A (cone 9)	3 mph	11 feet	J 1/4	K 1/2	P 1/2
Lady Maki's mowers, etc. 850	4.5 mph	34 feet	4.34	6.12	7.34
P70 at 450 rpm (Patent lit.)	4.5 mph	20 feet	16	20	24
Zoom (33 Series)					31

This product will also control (at the Program 3 rate) or suppress (at the Program 2 rate) the following weeds when applied prior to their emergence: bluegrass (annual), Poa annua, bittersweet, chickweed, vent, geranium (Carolina), pansy, plantain, and sheepsfoot.

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with crop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions by adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A minimum application rate can be made by spreading the remaining half at a right angle to the previous direction.

Avoid steaming, sifting, or other equipment malfunctions by properly adjusting and maintaining equipment. Avoid equipment malfunctions by applying uniform distribution of granules.

When split fertilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emergence.

TABLE 1. Recommended Application Use Rates* (Lbs. of this product/1,000 sq. ft.)

PROGRAM USE (Turfgrass or)	1 (Lbs. of this product/1,000 sq. ft.)		2 (Lbs. of this product/1,000 sq. ft.)		3 (Lbs. of this product/1,000 sq. ft.)	
	Preemergence (high)	Postemergence (low)	Preemergence (high)	Postemergence (low)	Preemergence (high)	Postemergence (low)
Region						
North	2.8	4.0	4.0	5.5		
Transition	4.0	5.5	8.4	11.0**		
South	5.5	8.4	11.0**			
Coastal South	8.4	11.0**				

**DO NOT apply more than 11 lbs. per application, and no more than 33 lbs. per year. Postemergence control is limited; see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below.

North: All other areas.
Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, NY, & RI.

South: AL, AR, AZ, CA, GA, LA, MS, NC, NN, NV, OK, SC, TN, & TX.

Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX.

**May require split or sequential applications for turf control.

Fall Applications: This product can also be applied in the fall in areas where the turfgrass is dormant. Apply this product in the early fall of the year. The fall application should be followed by an appropriate spring application to provide season-long control.

Turf for Improved Control: For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application. When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result. Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control.

CONTROL OF OTHER GRASS AND BROADLEAF WEEDS

Spring Applications: Used as directed for crabgrass control in the spring, this product will also control (at the Program 3 rate) or suppress (at the Program 2 rate) the following weeds when applied prior to their emergence: bahamagrass; bluegrass (annual); roughstalk, cowgrass (annual & perennial); smutgrass; searocket; Kentucky bluegrass (annual & perennial); annual ryegrass; perennial ryegrass; bermudagrass (annual & perennial); sturge (prostrate & spotted). Also controlled at the Program 2 rate.


Fall Applications: Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring.

SUGGESTED SPREADER SETTINGS

These suggested spreader settings are not intended to replace calibration. Please calibrate your spreader before applying product.
50 lbs. covers 17,000 sq. ft. (0.29 acre) at 174 lbs./1,000 sq. ft.
50 lbs. covers 15,000 sq. ft. (0.27 acre) at 174 lbs./1,000 sq. ft.
50 lbs. covers 9,000 sq. ft. (0.16 acre) at 240 lbs./1,000 sq. ft.
50 lbs. covers 5,000 sq. ft. (0.14 acre) at 360 lbs./1,000 sq. ft.

SPREADER	GROUND SPEED (ft./min.)	SPREADER SETTINGS			
		LOW RATE (0.1 lb./ft.²)	MEDIUM RATE (0.25 lb./ft.²)	HIGH RATE (0.5 lb./ft.²)	AREA
Andersen-Koehn 200, 395-200 (cone 9)	3 mph	11 feet	M	N 1/2	O 1/2
Andersen-Koehn 1000	3 mph	11 feet	M	N 1/2	O 1/2
Scotch & R. 9A (cone 9)	3 mph	11 feet	J 1/4	K 1/2	P 1/2
Lady Maki's mowers, etc. 850	4.5 mph	34 feet	4.34	6.12	7.34
P70 at 450 rpm (Patent lit.)	4.5 mph	20 feet	16	20	24
Zoom (33 Series)					31

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info.epa.gov>



Dispersing Granule Technology

with
0.103% DIMENSION®
Turf Herbicide
24-0-10

This product is a herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs.

GUARANTEED ANALYSIS	ACTIVE INGREDIENT
Total Nitrogen (N) 24.00%	Dithiopyr, 3,5-pyridinedicarbonyl acid 2-[(difluoromethyl)-(4-(2-methylpropyl)-6-(difluoromethyl))-5,5-dimethyl ester] 0.103%
2.46% Ammoniacal Nitrogen	
1.86% Urea Nitrogen	
6.81% Other Water Soluble Nitrogen	
12.59% Water Insoluble Nitrogen	
Soluble Potash (K ₂ O) 10.00%	TOTAL 100.00%
Sulfur (S) 6.15%	EPA Reg. No. 9198-120
6.15% Free Sulfur (S)	EPA Est. No. 9198-OH-TM, 9198-OH-EB, 9198-AL-001A
0.00% Ammonium Sulfate	(Underlined letter is first letter used in run code on end of bag.)
0.81% slowly available nitrogen from methyl urea and urea	
	FDZ

Visit our website: AndersonsGolfProducts.com

SGN 150

Net Wt 50 Lbs (22.68 kg)