

FIFRA 24(c) Special Local Need Label



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Opensight[®]
EPA Reg. No. 62719-597

24(c) Special Local Need Registration SLN OR-110017

**For Control of Herbaceous Broadleaf Weeds and Woody Plants in Forest
Plantations**

(For Distribution and Use Only in the State of Oregon)

Expiration Date: This label is valid until December 31, 2021, or until otherwise amended, withdrawn, canceled, or suspended.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read this SLN labeling and the label affixed to the container for Opensight™ specialty herbicide before applying. All applicable use directions, precautions and restrictions on this SLN labeling and the label affixed to the container must be followed.

Active Ingredients:

Potassium salt of 2-pyridine carboxylic acid, 4-amino-3,6-dichloro-	62.13%
Metsulfuron methyl.....	9.45%
Other Ingredients	<u>28.42%</u>
Total	100.0%

Acid Equivalent: aminopyralid (2-pyridine carboxylic acid, 4-amino-3,6-dichloro-) - 52.5%

Contains 0.62 pound potassium salt of aminopyralid active ingredient (0.525 pound acid equivalent) and 0.0945 pound metsulfuron methyl per pound of product

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Directions for Use

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Forest Plantations

Opensight™ specialty herbicide may be applied to forest plantations, as an aerial or ground broadcast application for site preparation, or as a spot application or directed spray application for conifer release (see Application Methods section) to control herbaceous broadleaf weeds and woody plants. Opensight may be applied alone or in tank-mix combinations with labeled rates of other herbicides provided: (1) the tank mix product is labeled for the timing and method of application for the use site to be treated and (2) mixing is not prohibited by the label of the registered tank mixed products. Use as directed in the Directions of Use section of the tank-mix partner.

Follow mixing instructions under the Application Methods section below. Unless otherwise directed, use an agricultural labeled crop oil concentrate, methylated seed oil, or non-ionic surfactant for all treatments. Apply surfactants at a minimum rate of 0.25% v/v (1 quart per 100 gallons spray solution) or at the manufacturers recommended rate.

Application Methods

For best control from broadcast and directed spray applications of Opensight, use a spray volume which will provide thorough plant coverage. Recommended spray volumes are usually 10 to 30 gallons per acre by air, or 10 to 100 gallons per acre by ground.

Forest Site Preparation Directions (aerial or ground broadcast, or spot applications)

Use up to 3.3 oz per acre of Opensight and apply in a suggested total spray volume of 10 to 30 gallons per acre. Tank mixtures with other herbicides registered for forest use, such as Accord XRT II, may improve control of woody brush, grasses, and herbaceous plants. When tank mixtures of herbicides are used for forest site preparation, labels for all products in the mixture must be followed; use the most restrictive label directions, including the longest recommended waiting period before planting.

Note: Conifers planted sooner than 2 months west of the Cascade Mountains and 4 months east of the Cascade Mountains after treatment with Opensight may be injured.

Directed Spray Applications for Conifer Release (including spot application)

NOTE: To prevent injury to or loss of desirable trees, including conifers, avoid spray contact with the foliage, especially small trees and their terminal buds, and avoid application in the root zone especially when making spot treatments.

Use Opensight to release conifers from competing invasive and broadleaf weeds such as thistles, maretail, bracken fern, wild carrot, mustards, knapweeds, tansy ragwort, and sensitive woody plants such as blackberry, buckbrush, snowberry, locust, redbud, and wild roses. The spray mixture should be directed onto foliage of competitive plants using knapsack or backpack sprayers with flat fan nozzles or equivalent any time after plants have reached full leaf size, but before autumn coloration. The majority of treated hardwoods should be less than 6 feet in height to ensure adequate spray coverage. Care should be taken to direct spray away from contact with conifer foliage, particularly foliage of desirable conifers.

Spot Application for site preparation or conifer release: Spot applications may be made at rates equivalent to the broadcast-applied rates of Opensight at 2 to 3.3 oz. per acre per annual growing season depending on the target species. Spray volume should be sufficient to thoroughly and uniformly wet weed foliage. Repeat treatments may be made, but the total amount of Opensight applied must not exceed 3.3 oz per acre per year. To prevent misapplication, spot treatments should be applied using calibrated application systems. Apply the spray mixture using a calibrated boom, boomless nozzle and, hand-held, or backpack sprayers.

Spot applications may be made at a rate of up to 0.22 lb acid equivalent aminopyralid (6.6 oz of Opensight) per acre; however, not more than 50% of an acre may be treated.

Restrictions:

- For use under this SLN label do not apply this product through any type of irrigation system.
- Read and follow all directions, restriction and precautions for manure management on the section 3 label for Opensight.
- Do not harvest hay or forage treated with Opensight.
- Do not treat more than 50% of an acre using spot treatment.
- Do not apply more than a total of 0.11 lb acid equivalent aminopyralid per acre (3.3 oz per acre of Opensight) per annual growing season as a result of broadcast, spot or repeat applications.
- Buffer Requirement Near Aquatic Areas: For herbicide applications in forestland, refer to the Oregon Forest Practices Act to determine if a buffer (no spray zone) or another type of requirement is mandated, <http://www.oregon.gov/ODF/herbicideinfo.shtml>

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R372-021

Accepted: 01/03/17

Replaces: R372-002