

Product Bulletin



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Blackhawk®

EPA Reg. No. 62719-523

Special 2(ee) Recommendation†

(For Distribution and Use Only in the states of Idaho, Minnesota, North Dakota, Oregon, and Washington)

Control of Potato Psyllid

ATTENTION

†This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Blackhawk® Naturalyte® insect control before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Blackhawk according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Blackhawk.

Directions for Use

Refer to product label for Blackhawk for General Use Precautions, Mixing Directions and Application Directions.

Blackhawk should be used as part of an integrated program to manage potato psyllid.

Apply 3.5 oz ounces of Blackhawk per acre to potatoes for control of potato psyllid. Begin at first signs of infestations. Due to the occurrence of multiple generations in a growing season, repeated applications may be required. Under moderate to high pest pressure, do not extend application interval beyond 7 to 10 days. Follow resistance management recommendations on the product label.

Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area. Refer to specific crop sections of the product label for additional information and restrictions.

®Trademark of Dow AgroSciences LLC

R373-007

Issued: 08/15/16

Replaces R373-003