



Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CALIFORNIA

Caparol® 4L

**EPA Reg. No. 100-620
EPA SLN No. CA-980017**

For Control Of Certain Weeds In Transplanted Celery

Active Ingredient:

Prometryn: 2,4-bis(isopropylamino)-6-(methylthio)-s-triazine	44.4%
Other Ingredients	55.6%
Total:	100.0%

Caparol 4L contains 4 lbs. active ingredient per gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, and precautions including statements pertaining to the Worker Protection Standards, on the EPA-registered Caparol 4L label.

This label must be in the possession of the user at the time of pesticide application.

To amend Caparol 4L label to include pre-transplant applications:

Single Pre-Transplant Application

On coarse-textured soils and soils low in organic matter, make one application of 2 to 4 pts. per acre of Caparol 4L in a minimum of 20 gallons of water. On fine-textured soils and soils high in organic matter, make one application of 3.2 to 4 pts. per acre of Caparol 4L in a minimum of 20 gallons of water. Make the application up to 21 days prior to transplanting. Make the application to a weed-free bed and do not disturb the treated area between application and

transplanting. Note: A post-transplant application will not be allowed if the pre-transplant application was made at the maximum allowable use rate (4 pts. per acre).

Split Pre-Transplant and Post-Transplant Application:

On coarse-textured soils and soils low in organic matter, the lower rate range of 2 to 4 pts. per acre of Caparol 4L in a minimum of 20 gallons of water is recommended. The maximum post-transplant application allowable on coarse-textured soils and soils low in organic matter is 3.2 pts. per acre.

On fine-textured soils and soils high in organic matter, the higher rate range of 3.2 to 4 pts. per acre of Caparol 4L in a minimum of 20 gallons of water is recommended.

Make the pre-transplant application up to 21 days prior to transplanting. Make the application to a weed-free bed and do not disturb the treated area between application and transplanting. A second application of Caparol 4L may be made during the 2 to 6 week period after transplanting and before weeds are two inches tall. The total amount of Caparol 4L applied to the crop (pre-transplant plus post-transplant applications) cannot exceed the maximum allowable use rate (4 pts. per acre).

Post Transplant Application

As indicated on the EPA-registered Caparol 4L label make one application of 2 to 3.2 pts. per acre of Caparol 4L on coarse-textured soils and soils low in organic matter in a minimum of 20 gallons of water. Make one application of 3.2 to 4 pts. per acre of Caparol 4L on fine-textured soils and soils high in organic matter in a minimum of 20 gallons of water. Application may be made over-the-top to celery during the 2 to 6 week period after transplanting and before weeds are two inches tall.

NOTE: To avoid possible illegal residues, do not make a Caparol 4L application within 45 days of harvest.

Rotational Crops: The following crops may be seeded five months after applying no more than 4 pts. of Caparol 4L per acre on celery: cabbage, celery, corn, onions, peas, and red beets.

Caparol® 4L is a trademark of Syngenta

24(c) registrant:
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419-8300

EPA SLN No: CA-980017

Label Code: CA0620052AA0204