



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

LUNA® TRANQUILITY

Reduced Application Rate for use in potato

EPA Reg. No. 264-1085

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF ARIZONA, ARKANSAS, CALIFORNIA, COLORADO, CONNECTICUT, FLORIDA, GEORGIA, IDAHO, ILLINOIS, INDIANA, IOWA, KENTUCKY, MAINE, MARYLAND, MICHIGAN, MINNESOTA, MISSISSIPPI, MISSOURI, MONTANA, NEBRASKA, NEW JERSEY, NEW MEXICO, NORTH CAROLINA, NORTH DAKOTA, OHIO, OKLAHOMA, OREGON, PENNSYLVANIA, RHODE ISLAND, SOUTH CAROLINA, TENNESSEE, TEXAS, UTAH, VERMONT, VIRGINIA, WASHINGTON, WEST VIRGINIA, WISCONSIN

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. It is a violation of federal law to use this product in a manner inconsistent with its label. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

POTATO		
Disease Control	Product Rate	Use Information
White mold (<i>Sclerotinia sclerotiorum</i>)	11.2 fl oz/acre	Begin fungicide applications preventatively. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the shorter intervals.
Early blight (<i>Alternaria solani</i>) Brown leaf spot (<i>Alternaria alternata</i>) Botrytis leaf spot (<i>Botrytis cinerea</i>) Black dot (<i>Colletotrichum coccodes</i>) Powdery mildew (<i>Erysiphe cichoracearum</i>) (<i>Leveillula taurica</i>) (<i>Erysiphe</i> spp.)	8.0 to 11.2 fl oz/acre	Begin fungicide applications preventatively. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the shorter intervals.

Restrictions: Do not apply more than 43.6 fl oz of LUNA®TRANQUILITY per acre per season. If aerial applications are made do not apply more than 33.6 fl oz of LUNA®TRANQUILITY per season. Do not apply LUNA®TRANQUILITY within 7 days of harvest. To limit the potential for development of disease resistance to these fungicide classes, do not make more than 2 sequential applications of LUNA®TRANQUILITY or any Group 7 or Group 9 containing fungicide before rotating with a fungicide from a different Group.

Use of this product according to this bulletin has not been reviewed or endorsed by the State of Indiana.

**For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)**

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*